

Complex Analysis HW 4

Ryan Chen

April 1, 2023

Note. Write $\operatorname{cis} t := \cos t + i \sin t$.

P81.3. Set

$$g(z) := \frac{z^3}{27} f(z)$$

so that

$$|g(z)| = \frac{|z|^3}{27} |f(z)|$$

Then on $|z| = 3$,

$$|g(z)| \leq \frac{3^3}{27} \cdot 1 = 1$$

and on $|z| = 1$,

$$|g(z)| \leq \frac{1^3}{27} \cdot 27 = 1$$

Thus by the maximum principle, on $1 \leq |z| \leq 3$,

$$|g(z)| \leq 1 \implies |f(z)| \leq \frac{27}{|z|^3}$$

P94.2 From $u(z) = z^2 + \bar{z}^2$, set

$$g(z) := \partial u(z) = 2z$$

$$G(z) := \int g(z) dz = z^2$$

$$f(z) := 2G(z) = 2z^2$$

We see f is entire and check that

$$\frac{1}{2} \left(f(z) + \overline{f(\bar{z})} \right) = \frac{1}{2} (2z^2 + 2\bar{z}^2) = z^2 + \bar{z}^2 = u(z)$$

P94.4 Pf. With u harmonic, there exists an entire function f such that $\operatorname{Re} f = u$. Then e^f is entire and

$$|e^f| = e^u < e^c$$

so by Liouville's theorem e^f is constant, hence f is constant. In turn, u is constant.

P96.4 Set $u(x, y) := y$ and $v(x, y) := \cosh \pi x \sin \pi y$.

$$v_{xx} = \pi^2 \cosh \pi x \sin \pi y, \quad v_{yy} = -\pi^2 \cosh \pi x \sin \pi y \implies v_{xx} + v_{yy} = 0$$

Thus v is harmonic. Also, $v|_{y=0} = v|_{y=1} = 0$. Then the function $w := u + v$ is a sum of harmonic functions hence harmonic, and w satisfies the boundary conditions $w|_{y=0} = 0$ and $w|_{y=1} = 1$.

P107 Pf. Fix z in the unbounded component of α^c . The winding number is

$$n(z, \alpha) = \frac{1}{2\pi i} \int_{\alpha} \frac{dw}{w - z}$$

Since α is bounded and closed, $\frac{1}{|w - z|}$ is continuous in w on α , so it attains a maximum at some $w_0 \in \alpha$.

$$|n(z, \alpha)| \leq \frac{1}{2\pi} \left| \int_{\alpha} \frac{dw}{w - z} \right| \leq \frac{\text{length}(\alpha)}{2\pi|w_0 - z|} \xrightarrow{z \rightarrow \infty} 0$$

so that $\lim_{z \rightarrow \infty} |n(z, \alpha)| = 0$. Thus $|n(z, \alpha)| < 1$ for large enough $|z|$. Now $n(z, \alpha) \in \mathbb{Z}$, so $|n(z, \alpha)| = 0$ hence $n(z, \alpha) = 0$ for large enough $|z|$. Also $n(z, \alpha)$ must be constant on components of α^c , thus $n(z, \alpha) = 0$ on the unbounded component of α^c .

P114.1 The Laurent expansions are:

(i) For $0 < |z| < 2$,

$$\frac{1}{z(z^2 + 4)} = \frac{1}{4z(1 - (-z^2/4))} = \frac{1}{4z} \sum_{k \geq 0} \frac{(-1)^k z^{2k}}{4^k} = \sum_{k \geq 0} \frac{(-1)^k z^{2k-1}}{4^{k+1}}$$

(ii) For $|z| > 2$,

$$\frac{1}{z(z^2 + 4)} = \frac{1}{z^3(1 - (-4/z^2))} = \frac{1}{z^3} \sum_{k \geq 0} \frac{(-1)^k 4^k}{z^{2k}} = \sum_{k \geq 0} \frac{(-1)^k 4^k}{z^{2k+3}}$$

(iii) For $0 < |z + 2i| < 2$,

$$\frac{1}{z(z^2 + 4)} = \frac{1}{z(z - 2i)(z + 2i)} = \frac{1}{z(z - 2i)} \sum_{k \geq 0} (z + 2i)^k$$

P119.1

(i) The function

$$f(z) = \frac{1}{(z - 4)(z^3 - 1)}$$

has singularities at $z = 4$ and $z = \text{cis } \frac{2\pi k}{3}$, $0 \leq k \leq 2$, i.e. $z = 4, 1, -\frac{1}{2} + \frac{\sqrt{3}}{2}i, -\frac{1}{2} - \frac{\sqrt{3}}{2}i$. Considering the circle $\alpha : |z - 2| = \frac{5}{2}$ has leftmost endpoint $-\frac{1}{2}$, the singularities that lie inside α are $z = 4, 1$. We now find

$$(z - 4)f(z) = \frac{1}{z^3 - 1} \xrightarrow{z \rightarrow 4} \frac{1}{64 - 1} = \frac{1}{63}$$

so that $\text{Res}(f, 4) = \frac{1}{63}$, and

$$(z - 1)f(z) = \frac{1}{(z - 4)(z^2 + z + 1)} \xrightarrow{z \rightarrow 1} \frac{1}{-3(1 + 1 + 1)} = -\frac{1}{9}$$

so that $\text{Res}(f, 1) = -\frac{1}{9}$. Thus

$$\int_{\alpha} f(z) dz = 2\pi i [\text{Res}(f, 4) + \text{Res}(f, 1)] = 2\pi i \left[\frac{1}{63} - \frac{1}{9} \right] = 2\pi i \left[-\frac{6}{63} \right] = -\frac{4\pi i}{21}$$

(ii) The function

$$g(z) = \frac{3z^2 + 2}{(z-1)(z^2 + 9)}$$

has singularities at $z = 1, \pm 3i$ which all lie within the circle $\beta : |z| = 5$. We find

$$(z-1)g(z) = \frac{3z^2 + 2}{z^2 + 9} \xrightarrow{z \rightarrow 1} \frac{3+2}{1+9} = \frac{5}{10} = \frac{1}{2}$$

so that $\text{Res}(g, 1) = \frac{1}{2}$, and

$$\begin{aligned} (z \mp 3i)g(z) &= \frac{3z^2 + 2}{(z-1)(z \pm 3i)} \xrightarrow{z \rightarrow \pm 3i} \frac{3(-9) + 2}{(\pm 3i - 1)(\pm 6i)} = \frac{-25}{-18 \mp 6i} \\ &= \frac{25}{18 \pm 6i} \frac{18 \mp 6i}{18 \mp 6i} = \frac{450 \mp 150i}{324 + 36} = \frac{450 \mp 150i}{360} = \frac{15 \mp 5i}{12} \end{aligned}$$

so that $\text{Res}(g, \pm 3i) = \frac{15 \mp 5i}{12}$. Thus

$$\begin{aligned} \int_{\beta} g(z) dz &= 2\pi i [\text{Res}(g, 1) + \text{Res}(g, 3i) + \text{Res}(g, -3i)] = 2\pi i \left[\frac{1}{2} + \frac{15-5i}{12} + \frac{15+5i}{12} \right] \\ &= 2\pi i \left[\frac{1}{2} + \frac{30}{12} \right] = 2\pi i \left[\frac{1}{2} + \frac{5}{2} \right] = 6\pi i \end{aligned}$$

P119.2

(i) The function

$$f(z) = \frac{1}{\cosh 2z}$$

has singularities at (write $w = 2z = x + iy$)

$$\cosh 2z = 0 \implies \cosh x \cos y + i \sinh x \sin y = 0 \implies \cosh x \cos y = 0, \sinh x \sin y = 0$$

The first equation gives

$$\cos y = 0 \implies y = \frac{\pi}{2} + \pi n, \quad n \in \mathbb{Z}$$

The second equation gives $x = 0$ or $y = \pi n$, so it must be the case $x = 0$ and $y = \frac{\pi}{2} + \pi n$.

$$w = i \left(\frac{\pi}{2} + \pi n \right) \implies z = \frac{w}{2} = \left(\frac{\pi}{4} + \frac{\pi}{2} n \right) i$$

The only singularities that lie within the circle $\gamma : |z| = 1$ are for $n = 0, -1$, i.e. $z = \pm \frac{\pi}{4}i$. Label the points as z_1 and z_2 .

$$\text{Res}(f, z_j) = \lim_{z \rightarrow z_j} (z - z_j) f(z) = \lim_{z \rightarrow z_j} \frac{z - z_j}{\cosh 2z}$$

Using L'Hopital's rule,

$$\text{Res}(f, z_j) = \lim_{z \rightarrow z_j} (z - z_j) f(z) \frac{1}{2 \sinh 2z} = \frac{1}{2 \sinh 2z_j} = \frac{1}{2 \sinh(\pm \frac{\pi}{2}i)} = \frac{1}{2i \sin(\pm \frac{\pi}{2})} = -\frac{i}{2(\pm 1)} = \mp \frac{i}{2}$$

Thus

$$\int_{\gamma} f(z) dz = 2\pi i \left[\text{Res}\left(f, \frac{\pi}{4}i\right) + \text{Res}\left(f, -\frac{\pi}{4}i\right) \right] = 2\pi i \left[-\frac{i}{2} + \frac{i}{2} \right] = 0$$

(ii) The function

$$g(z) = \cot z = \frac{\cos z}{\sin z}$$

has singularities at

$$\sin z = 0 \implies z = k\pi, \quad k \in \mathbb{Z}$$

The only singularity that lies within the circle $\gamma : |z| = 1$ is for $k = 0$, i.e. $z = 0$.

$$\text{Res}(g, 0) = \lim_{z \rightarrow 0} zg(z) = \lim_{z \rightarrow 0} \frac{z \cos z}{\sin z}$$

Using L'Hopital's rule,

$$\text{Res}(g, 0) = \lim_{z \rightarrow 0} \frac{\cos z - z \sin z}{\cos z} = \frac{1 - 0}{1} = 1$$

Thus

$$\int_{\gamma} g(z) dz = 2\pi i \text{Res}(g, 0) = 2\pi i$$

P127.4 Pf. Set

$$I := \int_0^{\infty} f(x) dx, \quad f(z) := e^{iz^2}$$

Fix large $R > 0$ and define the contours

$$\alpha := [0, R] + C_R - \gamma_R, \quad C_R : z = Re^{it}, \quad 0 \leq t \leq \frac{\pi}{4}, \quad \gamma_R : z = te^{i\pi/4}, \quad 0 \leq t \leq R$$

Since f is analytic on the region enclosed by α ,

$$0 = \int_{\alpha} f(z) dz = \int_{[0, R]} f(z) dz + \int_{C_R} f(z) dz - \int_{\gamma_R} f(z) dz \quad (127.4.1)$$

We compute the third integral. On γ_R ,

$$z = te^{i\pi/4}, \quad 0 \leq t \leq R \implies z^2 = t^2 e^{i\pi/2} = it^2 \implies f(z) = e^{-it^2}$$

Using the Gaussian integral,

$$\int_{\gamma_R} f(z) dz = e^{i\pi/4} \int_0^R e^{-t^2} dt \xrightarrow{R \rightarrow \infty} e^{i\pi/4} \int_0^{\infty} e^{-t^2} dt = e^{i\pi/4} \frac{\sqrt{\pi}}{2}$$

We compute the second integral. On C_R ,

$$z = Re^{it}, \quad 0 \leq t \leq \frac{\pi}{4}, \implies z^2 = R^2 e^{i2t} = R^2 \cos 2t + iR^2 \sin 2t \implies iz^2 = iR^2 \cos 2t - R^2 \sin 2t \implies |f(z)| = e^{-R^2 \sin 2t}$$

Now using the fact $\sin t \geq \frac{2t}{\pi}$ for $0 \leq t \leq \frac{\pi}{2}$, we have $\sin 2t \geq \frac{4t}{\pi}$ for $0 \leq t \leq \frac{\pi}{4}$, hence $|f(z)| \leq e^{-R^2 4t/\pi}$.

$$\left| \int_{C_R} f(z) dz \right| \leq \int_{C_R} |f(z)| |dz| \leq \int_0^{\pi/4} e^{-R^2 4t/\pi} R dt = -\frac{R\pi}{4R^2} e^{-4R^2 t/\pi} \Big|_0^{\pi/4} = -\frac{\pi}{4R} (e^{-R^2} - 1) \xrightarrow{R \rightarrow \infty} 0(0-1) = 0$$

Putting together the calculations, as we take $R \rightarrow \infty$ in equation (127.4.1),

$$0 = I - e^{i\pi/4} \frac{\sqrt{\pi}}{2} \implies I = e^{i\pi/4} \frac{\sqrt{\pi}}{2} = \left(\frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) \frac{\sqrt{\pi}}{2} \implies \int_0^{\infty} \cos(x^2) dx = \text{Im } I = \frac{\sqrt{\pi}}{2\sqrt{2}}$$

P131.1 Set $D := \{|z| < 1\}$, so that $\partial D = \{|z| = 1\}$.

(i) Write

$$h(z) = f(z) + g(z), \quad f(z) = 5z^4, \quad g(z) = z^2 + 2z - i$$

On ∂D ,

$$|f(z)| = 5|z|^4 = 5, \quad |g(z)| \leq |z|^2 + 2|z| + |i| = 4 \implies |f(z)| > |g(z)|$$

In D , f has a root at $z = 0$ of order 4, so by Rouché's theorem h has 4 roots in D .

(ii) Write

$$h(z) = g(z) + f(z), \quad g(z) = z^3, \quad f(z) = 8z^2 - 8z + 2$$

On ∂D , using the reverse triangle inequality,

$$|f(z)| \geq |8|z|^2 - 8|z| - |2|| = |8 - 8 - 2| = 2, \quad |g(z)| = |z|^3 = 1 \implies |f(z)| > |g(z)|$$

The roots of f are

$$0 = f(z) = 8z^2 - 8z + 2 = (4z - 2)(2z - 1) \implies z = \frac{1}{2}, \frac{1}{2} \in D$$

so by Rouché's theorem h has 2 roots in D .

(iii) Write

$$h(z) = g(z) + f(z), \quad g(z) = e^z, \quad f(z) = 3z$$

On ∂D , $|z| = 1$, so write $z = \text{cis } t$.

$$|f(z)| = 3|z| = 3, \quad |g(z)| = |e^z| = e^{\cos t} \leq e^1 < 3 \implies |f(z)| > |g(z)|$$

In D , f has a root at $z = 0$ of order 1, so by Rouché's theorem h has 1 root in D .