

Scientific Computing HW 7

Ryan Chen

March 17, 2023

Problem 1.

- (a) Multiplying the BVP by -1 and integrating it, we find $k(x)u' = M$ for some constant M , i.e. $u' = \frac{M}{k(x)}$. The solution is then

$$u(x) = u_a + \int_a^x \frac{M}{k(s)} ds$$

If $x \leq c$ then

$$u(x) = u_a + \int_a^x \frac{M}{k_1} ds = u_a + \frac{M}{k_1}(x - a)$$

If $x > c$ then

$$u(x) = u_a + \int_a^c \frac{M}{k(s)} ds + \int_c^x \frac{M}{k(s)} ds = u_a + \int_a^c \frac{M}{k_1} ds + \int_c^x \frac{M}{k_2} ds = u_a + \frac{M}{k_1}(c - a) + \frac{M}{k_2}(x - c)$$


Apply BCs.

$$u_b = u(b) = u_a + \frac{M}{k_1}(c - a) + \frac{M}{k_2}(b - c) = u_a + M \left[\frac{c - a}{k_1} + \frac{b - c}{k_2} \right] \implies M = \frac{u_b - u_a}{\frac{c - a}{k_1} + \frac{b - c}{k_2}}$$

In summary, the solution is

$$u(x) = \begin{cases} u_a + \frac{M}{k_1}(x - a), & x \leq c \\ u_a + \frac{M}{k_1}(c - a) + \frac{M}{k_2}(x - c), & x > c \end{cases} \quad \text{where} \quad M = \frac{u_b - u_a}{\frac{c - a}{k_1} + \frac{b - c}{k_2}}$$

- (b) Given the parameters, it is enough to solve for the values of the 9 mesh points shown below.



The finite difference scheme is


$$L_h u_P = -\frac{1}{h^2} [k_w u_W + k_e u_E - (k_e + k_w) u_P] = 0 \implies -(k_w + k_e) u_P + k_w u_W + k_e u_E = 0$$

Applying the scheme to each mesh point, we obtain a linear system.

	1	2	3	4	5	6	7	8	9			
1	$-2k_1$	k_1								u_1		$-k_1 u_a$
2	k_1	$-2k_1$	k_1							u_2		0
3		k_1	$-2k_1$	k_1						u_3		0
4			k_1	$-k_1 - k_2$	k_2					u_4		0
5				k_2	$-2k_2$	k_2				u_5	$=$	0
6					k_2	$-2k_2$	k_2			u_6		0
7						k_2	$-2k_2$	k_2		u_7		0
8							k_2	$-2k_2$	k_2	u_8		0
9								k_2	$-2k_2$	u_9		$-k_2 u_b$

We solve it and plot the numerical solution u along with the exact solution from part (a). In this case the solutions agree exactly.




Code: <https://github.com/RokettoJanpu/Scientific-Computing-2/blob/main/hw7q1.ipynb>



Problem 2. Code:

<https://github.com/RokettoJanpu/Scientific-Computing-2/blob/main/hw7q2.ipynb>

Below are meshes for the L-shape, the pentagon with a smaller pentagon removed, and the semicircle with two smaller circles removed.



Problem 3.

- (a) Multiply the BVP by -1 and multiply by w .

$$w(x)u''(x) = -w(x)f(x)$$

Integrate on $[0, 1]$. Integrating the LHS by parts, we repeatedly differentiate w and integrate u .

$$\begin{aligned} \int_0^1 w(x)u''(x)dx &= w(x)u'(x) \Big|_0^1 - w'(x)u(x) \Big|_0^1 + \int_0^1 w''(x)u(x)dx \\ &= w(1)u'(1) - w(0)u'(0) - w'(1)u(1) + w'(0)u(0) + \int_0^1 w''(x)u(x)dx \\ &= -w'(1) + w'(0) + \int_0^1 w''(x)u(x)dx \end{aligned}$$

We obtain an integral equation for u .

$$-w'(1) + w'(0) + \int_0^1 w''(x)u(x)dx = - \int_0^1 w(x)f(x)dx \implies \int_0^1 w''(x)u(x)dx = - \int_0^1 w(x)f(x)dx + w'(1) - w'(0)$$

- (b) Fix $1 \leq i \leq N$. The stiffness matrix entries are

$$A_{ij} = \int_0^1 \varphi'_i(x)\varphi'_j(x)dx$$

First compute

$$\varphi'_i(x) = \begin{cases} 0, & x \leq x_{i-1} \text{ or } x \geq x_{i+1} \\ \frac{1}{x_i - x_{i-1}}, & x_{i-1} < x \leq x_i \\ -\frac{1}{x_{i+1} - x_i}, & x_i < x < x_{i+1} \end{cases}$$

We examine cases for the value of j .

- If $j \leq i - 2$ or $j \geq i + 2$ then $\varphi'_i\varphi'_j = 0$ hence $A_{ij} = 0$.
- If $j = i - 1$ then

$$\varphi'_i(x)\varphi'_j(x) = \begin{cases} 0, & x \leq x_j \text{ or } x \geq x_i \\ -\frac{1}{(x_i - x_{i-1})(x_{j+1} - x_j)}, & x_j < x < x_i \end{cases}$$

hence

$$A_{ij} = -\frac{x_i - x_j}{(x_i - x_{i-1})(x_{j+1} - x_j)}$$

- If $j = i + 1$ then

$$\varphi'_i(x)\varphi'_j(x) = \begin{cases} 0, & x \leq x_i \text{ or } x \geq x_j \\ -\frac{1}{(x_{i+1} - x_i)(x_j - x_{j-1})}, & x_i < x < x_j \end{cases}$$

hence

$$A_{ij} = -\frac{x_j - x_i}{(x_{i+1} - x_i)(x_j - x_{j-1})}$$

- If $j = i$ then

$$\varphi'_i(x)\varphi'_j(x) = \begin{cases} 0, & x \leq x_{i-1} \text{ or } x \geq x_{i+1} \\ \frac{1}{(x_i - x_{i-1})^2}, & x_{i-1} < x \leq x_i \\ \frac{1}{(x_{i+1} - x_i)^2}, & x_i < x < x_{i+1} \end{cases}$$

hence

$$A_{ij} = \frac{x_i - x_{i-1}}{(x_i - x_{i-1})^2} + \frac{x_{i+1} - x_i}{(x_{i+1} - x_i)^2} = \frac{1}{x_i - x_{i-1}} + \frac{1}{x_{i+1} - x_i}$$

In summary,

$$A_{ij} = \begin{cases} 0, & j \leq i-2 \text{ or } j \geq i+2 \\ -\frac{x_i - x_j}{(x_i - x_{i-1})(x_{j+1} - x_j)}, & j = i-1 \\ -\frac{x_j - x_i}{(x_{i+1} - x_i)(x_j - x_{j-1})}, & j = i+1 \\ \frac{1}{x_i - x_{i-1}} + \frac{1}{x_{i+1} - x_i}, & j = i \end{cases}$$

Problem 4.

- (a) **Pf.** The matrix G is given by

$$G = A^{-1} \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}, \quad A := \begin{bmatrix} 1 & 1 & 1 \\ x_1 & x_2 & x_3 \\ y_1 & y_2 & y_3 \end{bmatrix}$$

We find A^{-1} by its adjugate. By cofactor expansion over the first row,

$$D := \det(A) = \begin{vmatrix} x_2 & x_3 \\ y_2 & y_3 \end{vmatrix} - \begin{vmatrix} x_1 & x_3 \\ y_1 & y_3 \end{vmatrix} + \begin{vmatrix} x_1 & x_2 \\ y_1 & y_2 \end{vmatrix} = x_2y_3 - x_3y_2 - x_1y_3 + x_3y_1 + x_1y_2 - x_2y_1$$

The matrix of cofactors is

$$\text{cof}(A) = \begin{bmatrix} x_2y_3 - x_3y_2 & -x_1y_3 + x_3y_1 & x_1y_2 - x_2y_1 \\ -y_3 + y_2 & y_3 - y_1 & -y_2 + y_1 \\ x_3 - x_2 & -x_3 + x_1 & x_2 - x_1 \end{bmatrix}$$

The adjugate of A is

$$\text{adj}(A) = \text{cof}(A)^T = \begin{bmatrix} x_2y_3 - x_3y_2 & -y_3 + y_2 & x_3 - x_2 \\ -x_1y_3 + x_3y_1 & y_3 - y_1 & -x_3 + x_1 \\ x_1y_2 - x_2y_1 & -y_2 + y_1 & x_2 - x_1 \end{bmatrix}$$

Thus

$$G = \frac{1}{D} \text{adj}(A) \begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 1 \end{bmatrix} = \frac{1}{D} \begin{bmatrix} y_2 - y_3 & x_3 - x_2 \\ y_3 - y_1 & x_1 - x_3 \\ y_1 - y_2 & x_2 - x_1 \end{bmatrix}$$

Fix an even permutation (i, j, k) of $1, 2, 3$ (i.e. one of $(1, 2, 3)$, $(2, 3, 1)$, $(3, 1, 2)$).

$$\eta_i(x, y) = \frac{\begin{vmatrix} 1 & x & y \\ 1 & x_j & y_j \\ 1 & x_k & y_k \end{vmatrix}}{\begin{vmatrix} 1 & x_i & y_i \\ 1 & x_j & y_j \\ 1 & x_k & y_k \end{vmatrix}}$$

Since the permutation (i, j, k) is even, the denominator is $\det(A^T) = \det(A) = D$. By cofactor expansion over the first row, the numerator is $x_j y_k - x_k y_j - (y_k - y_j)x + (x_k - x_j)y$, so that

$$\partial_x \eta_i(x, y) = \frac{1}{D}(y_j - y_k), \quad \partial_y \eta_i(x, y) = \frac{1}{D}(x_k - x_j)$$

Thus

$$\begin{bmatrix} \partial_x \eta_1(x, y) & \partial_y \eta_1(x, y) \\ \partial_x \eta_2(x, y) & \partial_y \eta_2(x, y) \\ \partial_x \eta_3(x, y) & \partial_y \eta_3(x, y) \end{bmatrix} = \frac{1}{D} \begin{bmatrix} y_2 - y_3 & x_3 - x_2 \\ y_3 - y_1 & x_1 - x_3 \\ y_1 - y_2 & x_2 - x_1 \end{bmatrix} = G$$

(b) Using the functions η_j from (a) and the fact $\nabla \eta_j$ is the j th row of G ,

$$\begin{aligned} u(x, y) &= \sum_{j=1}^3 u_j \eta_j(x, y) \implies \nabla u(x, y) = \sum_{j=1}^3 u_j \nabla \eta_j(x, y) = \frac{1}{D} \left(u_1 \begin{bmatrix} y_2 - y_3 \\ x_3 - x_2 \end{bmatrix} + u_2 \begin{bmatrix} y_3 - y_1 \\ x_1 - x_3 \end{bmatrix} + u_3 \begin{bmatrix} y_1 - y_2 \\ x_2 - x_1 \end{bmatrix} \right) \\ &\implies \nabla u(x, y) = (x_2 y_3 - x_3 y_2 - x_1 y_3 + x_3 y_1 + x_1 y_2 - x_2 y_1)^{-1} \begin{bmatrix} u_1(y_2 - y_3) + u_2(y_3 - y_1) + u_3(y_1 - y_2) \\ u_1(x_3 - x_2) + u_2(x_1 - x_3) + u_3(x_2 - x_1) \end{bmatrix} \end{aligned}$$