

HW

Ryan Chen

May 18, 2023

Problem 1

(a) In this part we use the fact

$$\int_{-\infty}^{\infty} \exp(-ax^2 + bx) dx = \left(\frac{\pi}{a}\right)^{1/2} \exp\left[\frac{b^2}{4a}\right]$$

Take the Fourier transform of the PDE in x , using the fact $\partial_x^n \psi = (i\xi)^n \psi$.

$$\hat{\psi}_t = \frac{i}{2}(i\xi)^2 \hat{\psi} = -\frac{i}{2}\xi^2 \hat{\psi} \implies \hat{\psi}(\xi, t) = \hat{\psi}_0(\xi) \exp\left[-\frac{i}{2}\xi^2 t\right]$$

Take the inverse Fourier transform.

$$\psi(x, t) = \frac{1}{(2\pi)^{1/2}} \int_{-\infty}^{\infty} \hat{\psi}_0(\xi) \exp\left[ix\xi - \frac{i}{2}t\xi^2\right] d\xi$$

Take the Fourier transform of the initial condition.

$$\begin{aligned} \hat{\psi}_0(\xi) &= \frac{1}{(2\pi)^{1/2}} \frac{1}{(2\pi\sigma_0^2)^{1/4}} \int_{-\infty}^{\infty} \exp\left[-\frac{x^2}{4\sigma_0^2} + ik_0x - i\xi x\right] dx \\ &= \frac{1}{(2\pi)^{3/4}\sigma_0^{1/2}} \int_{-\infty}^{\infty} \exp\left[-\frac{x^2}{4\sigma_0^2} + i(k_0 - \xi)x\right] dx \\ &= \frac{1}{(2\pi)^{3/4}\sigma_0^{1/2}} \pi^{1/2} 2\sigma_0 \exp[-(\xi - k_0)^2 \sigma_0^2] \\ &= \frac{2^{1/4}\sigma_0^{1/2}}{\pi^{1/4}} \exp[-(\xi - k_0)^2 \sigma_0^2] \end{aligned}$$

Then

$$\psi(x, t) = \frac{1}{(2\pi)^{1/2}} \frac{2^{1/4}\sigma_0^{1/2}}{\pi^{1/4}} \int_{-\infty}^{\infty} \exp\left[-\sigma_0^2(\xi - k_0)^2 - \frac{i}{2}t\xi^2 + ix\xi\right] d\xi$$

Rewrite the argument of exp as

$$-\sigma_0^2(\xi - k_0)^2 - \frac{i}{2}t\xi^2 + ix\xi = -\sigma_0^2(\xi^2 + k_0^2 - 2k_0\xi) - \frac{i}{2}t\xi^2 + ix\xi = -\left(\sigma_0^2 + \frac{i}{2}t\right)\xi^2 + (ix + 2\sigma_0^2 k_0)\xi - \sigma_0^2 k_0^2$$

so that

$$\begin{aligned}
\psi(x, t) &= \frac{\sigma_0^{1/2}}{2^{1/4}\pi^{3/4}} \int_{-\infty}^{\infty} \exp\left[-\left(\sigma_0^2 + \frac{i}{2}t\right)\xi^2 + (ix + 2\sigma_0^2 k_0)\xi - \sigma_0^2 k_0^2\right] d\xi \\
&= \frac{\sigma_0^{1/2}}{2^{1/4}\pi^{3/4}} e^{-\sigma_0^2 k_0^2} \left(\frac{\pi}{\sigma_0^2 + \frac{i}{2}t}\right)^{1/2} \exp\left[\frac{-x^2 + 4\sigma_0^4 k_0^2 + 4i\sigma_0^2 k_0 x}{4(\sigma_0^2 + \frac{i}{2}t)}\right] \\
&= \frac{\sigma_0^{1/2} e^{-\sigma_0^2 k_0^2}}{2^{1/4}\pi^{1/4}} \left(\sigma_0^2 + \frac{i}{2}t\right)^{-1/2} \exp\left[\frac{-x^2 + 4\sigma_0^4 k_0^2 + 4i\sigma_0^2 k_0 x}{4(\sigma_0^2 + \frac{i}{2}t)}\right]
\end{aligned}$$

(b) Discretize the PDE in space with stepsize h and use central differences.

$$\psi'_j(t) = \frac{i}{2h^2}[\psi_{j+1}(t) + \psi_{j-1}(t) - 2\psi_j(t)]$$

Let v be such that $v(x_j, t) = \psi_j(t)$.

$$v_t(x, t) = \frac{i}{2h^2}[v(x+h, t) + v(x-h, t) - 2v(x, t)]$$

Taylor expand.

$$\begin{aligned}
v(x+h, t) &= v + hv_x + \frac{1}{2}h^2v_{xx} + \frac{1}{6}h^3v_{xxx} + \frac{1}{24}h^4v_{xxxx} + \frac{1}{120}h^5v_{xxxxx} + O(h^6) \\
v(x-h, t) &= v - hv_x + \frac{1}{2}h^2v_{xx} - \frac{1}{6}h^3v_{xxx} + \frac{1}{24}h^4v_{xxxx} - \frac{1}{120}h^5v_{xxxxx} + O(h^6)
\end{aligned}$$

Plug in.

$$v_t = \frac{i}{2h^2} \left[h^2v_{xx} + \frac{1}{12}h^4v_{xxxx} + O(h^6) \right] = \frac{i}{2}v_{xx} + \frac{i}{24}h^2v_{xxxx} + O(h^4)$$

We obtain the (third order) modified equation.

$$v_t - \frac{i}{2}v_{xx} = \frac{i}{24}h^2v_{xxxx}$$

The Fourier transform of the RHS term is

$$\frac{i}{24}h^2\xi^4\hat{v}$$

so its corresponding term within the solution $\hat{v}(\xi, t)$ in Fourier space is

$$\exp\left[\frac{i}{24}h^2\xi^4t\right]$$

Thus the modified equation introduces artificial Fourier modes which do not decay over time.

Problem 3

(a) Fix a test function $v \in H_0^1(\Omega)$. Multiply the PDE by v and integrate over Ω .

$$\int_{\Omega} \epsilon v \Delta u dx = \int_{\Omega} (u^3 - u)v dx$$

Using Green's first identity and the fact $v = 0$ on $\partial\Omega$, the LHS is

$$\int_{\Omega} \epsilon v \Delta u dx = - \int_{\Omega} \epsilon \nabla u \cdot \nabla v dx + \int_{\partial\Omega} \epsilon v \frac{\partial u}{\partial n} ds = - \int_{\Omega} \epsilon \nabla u \cdot \nabla v dx$$

Then we obtain the weak formulation.

$$-\int_{\Omega} \epsilon \nabla u \cdot \nabla v dx = \int_{\Omega} (u^3 - u) v dx \implies \int_{\Omega} \epsilon \nabla u \cdot \nabla v dx - \int_{\Omega} (u - u^3) v dx = 0$$

(b) e

(c) Code for problem 3:

<https://github.com/RokettoJanpu/Scientific-Computing-2/blob/main/FINAL%20q3.ipynb>

Below is a mesh of Ω .

