

# Scientific Computing HW 5

Ryan Chen

October 1, 2024

1. Throughout we use the Matlab syntax for matrices and submatrices.

```
(a)  for  $i = 1, \dots, n$  do
       $M[i, i:] \leftarrow M[i, i:] / M[i, i]$ 
      for  $j \neq i$  do
           $M[j, i:] \leftarrow M[j, i:] - M[j, i] * M[i, i:]$ 
      end for
  end for
```

The flop count in the reassignment of  $M[j, i:]$  is  $2(2n - i + 1)$  due to it being a row vector with  $2n - i + 1$  entries. The flop count of the whole algorithm is

$$\begin{aligned} W(n) &= \sum_{i=1}^n 2(2n - i + 1)(n - 1) \\ &\sim (2n - 1) \int_0^n (2n - x + 1) dx \\ &= 2(n - 1) \left( 2nx - \frac{1}{2}x^2 + x \right) \Big|_0^n \\ &= 2(n - 1) \left( 2n^2 - \frac{1}{2}n^2 + n \right) \\ &= 3n^3 + O(n^2) \end{aligned}$$

(b) We can view the problem in four parts: The LU decomposition, finding  $L^{-1}$ , finding  $U^{-1}$ , and finding  $U^{-1}L^{-1}$ .

- The LU decomposition flop count is  $W_1(n) = \frac{2}{3}n^3 + O(n^2)$ .
- The idea of finding  $L^{-1}$ : From  $LL^{-1} = I$ , we can find the  $k$ th column of  $L^{-1}$  by solving  $Lx = e_k$ . Since  $L$  is lower triangular, we do this by forward substitution.

```
 $L^{-1} \leftarrow 0_{n \times n}$ 
for  $k = 1, \dots, n$  do
     $x \leftarrow 0_{n \times 1}$ 
     $x[k] \leftarrow 1$ 
    for  $j = k + 1, \dots, n$  do
         $S \leftarrow -L[j, k]$ 
        for  $i = k + 1, \dots, j - 1$  do
             $S \leftarrow S - L[j, i] * x[i]$ 
        end for
         $x[j] \leftarrow S$ 
    end for
     $L^{-1}[:, k] \leftarrow x$ 
end for
```

The flop count in the reassignment of  $S$  is 2, so the flop count in the  $i$  loop is  $2(j-1-k)$ . The flop count of the algorithm is

$$\begin{aligned}
W_2(n) &= \sum_{k=1}^n \sum_{j=k+1}^n 2(j-1-k) \\
&\sim 2 \int_0^n \int_x^n (y-1-x) dy dx \\
&= 2 \int_0^n \left( \frac{1}{2}y^2 - y - xy \right) \Big|_x^n dx \\
&= 2 \int_0^n \left( \frac{1}{2}n^2 - n - nx + \frac{1}{2}x^2 + x \right) dx \\
&= 2 \left( \frac{1}{2}n^2x - nx - \frac{1}{2}nx^2 + \frac{1}{6}x^3 + \frac{1}{2}x^2 \right) \Big|_0^n \\
&= 2 \left( \frac{1}{2}n^3 - n^2 - \frac{1}{2}n^3 + \frac{1}{6}n^3 + \frac{1}{2}n^2 \right) \\
&= \frac{1}{3}n^3 + O(n^2)
\end{aligned}$$

- Similarly, we can find the  $k$ th column of  $U^{-1}$  by solving  $Ux = e_k$ . Since  $U$  is upper triangular, we do this by backward substitution.

```

 $U^{-1} \leftarrow 0_{n \times n}$ 
for  $k = 1, \dots, n$  do
   $x \leftarrow 0_{n \times 1}$ 
  for  $j = k-1, \dots, 1$  (decrement  $j$  after each iteration) do
     $S \leftarrow 0$ 
    for  $i = j+1, \dots, k$  do
       $S \leftarrow S - U[j, i] * x[i]$ 
    end for
     $x[j] \leftarrow S/U[j, j]$ 
  end for
   $U^{-1}[:, k] \leftarrow x$ 
end for

```

The flop count in the reassignment of  $S$  is 2, so the flop count in the  $i$  loop is  $2(k-j)$ . Using WolframAlpha, the flop count of the algorithm is

$$\begin{aligned}
W_3(n) &= \sum_{k=1}^n \sum_{j=1}^{k-1} (2(k-j) + 1) \\
&\sim \int_0^n \int_0^x (2x - 2y + 1) dy dx \\
&= \int_0^n (2xy - y^2 + y) \Big|_0^x dx \\
&= \int_0^n (x^2 + x) dx \\
&= \frac{1}{3}n^3 + O(n^2)
\end{aligned}$$

- Note that  $U^{-1}L^{-1}$  is a product of an upper triangular matrix and a lower triangular matrix.

The positions of the zeros allows us to drop some terms when computing entries,

$$(A^{-1})_{ij} = \sum_{k=1}^n (U^{-1})_{ik} (L^{-1})_{kj} = \sum_{k=\max(i,j)}^n (U^{-1})_{ik} (L^{-1})_{kj}$$

```

A-1 ← 0n×n
for i = 1, ..., n do
  for j = 1, ..., n do
    S ← 0
    for k = max(i, j), ..., n do
      S ← S + U-1[i, k] * L-1[k, j]
    end for
    A-1[i, j] ← S
  end for
end for

```

The flop count in the reassignment of  $S$  is 2, so the flop count in the  $k$  loop is  $2(n - \max(i, j) + 1)$ . Using WolframAlpha, the flop count of the algorithm is

$$\begin{aligned}
W_4(n) &= \sum_{i=1}^n \sum_{j=1}^n (2(n - \max(i, j)) + 1) \\
&\sim \int_0^n \int_0^n (2n - 2\max(x, y) + 1) dy dx \\
&= 2n^3 - 2 \underbrace{\int_0^n \int_0^n \max(x, y) dy dx}_{=: I(x)} + n^2
\end{aligned}$$

To rewrite  $I(x)$ , consider

$$\max(x, y) = \begin{cases} x, & y \leq x \\ y, & y > x \end{cases}$$

Then compute

$$\begin{aligned}
I(x) &= \int_0^x x dy = \int_x^n y dy \\
&= x^2 + \frac{1}{2}(n^2 - x^2) \\
&= \frac{1}{2}(n^2 + x^2)
\end{aligned}$$

Thus

$$\begin{aligned}
W_4(n) &\sim 2n^3 - \int_0^n (n^2 + x^2) dx + n^2 \\
&= 2n^3 - n^3 - \frac{1}{3}n^3 + n^2 \\
&= \frac{2}{3}n^3 + O(n^2)
\end{aligned}$$

In total, the flop count of finding  $A^{-1}$  via its LU decomposition is

$$W(n) = W_1(n) + W_2(n) + W_3(n) + W_4(n) = 2n^3 + O(n^2)$$

2. (a) i. Fix  $A, B \in \mathcal{L}$  and let  $C := AB$ . For  $i < j$ ,

$$\begin{aligned} c_{ij} &= \sum_{k=1}^n a_{ik} b_{kj} \\ &= \sum_{k=1}^i a_{ik} b_{kj} + \sum_{k=i+1}^n a_{ik} b_{kj} \\ &= 0 + 0 = 0 \end{aligned} \quad b_{kj} = 0 \text{ for } k < j \text{ and } a_{ik} = 0 \text{ for } k > i$$

Thus  $C$  is lower triangular. For all  $i$ ,

$$\begin{aligned} c_{ii} &= \sum_{k=1}^n a_{ik} b_{ki} \\ &= \sum_{k=1}^{i-1} a_{ik} b_{ki} + a_{ii} b_{ii} + \sum_{k=i+1}^n a_{ik} b_{ki} \\ &= 0 + a_{ii} b_{ii} + 0 \end{aligned} \quad \begin{aligned} b_{ki} &= 0 \text{ for } k < i \text{ and } a_{ik} = 0 \text{ for } k > i \\ b_{ii}, a_{ii} &> 0 \end{aligned}$$

Thus  $C$  has positive diagonal entries. We conclude  $C \in \mathcal{L}$ .

- ii. Fix  $A \in L$ . Since  $A$  is lower triangular,  $\det A$  is the product of its diagonal entries, which are positive, so  $\det A \neq 0$ . Thus  $A$  is nonsingular, so let  $B := A^{-1}$ . To show  $B \in \mathcal{L}$ , we fix  $j$  and aim to show  $b_{ij} = 0$  for all  $i < j$ , hence  $B$  is lower triangular, and  $b_{jj} > 0$ , hence  $B$  has positive diagonal entries. From  $AB = I$  and  $A$  being lower triangular, hence  $a_{ik} = 0$  for  $k > i$ , we have

$$\sum_{k=1}^i a_{ik} b_{kj} = \delta_{ij}$$

Write out the corresponding equations for all  $i < j$ :

$$\begin{aligned} i = 1 : a_{11} b_{1j} &= 0 \\ i = 2 : a_{21} b_{1j} + a_{22} b_{2j} &= 0 \\ &\vdots \\ i = j : a_{j1} b_{1j} + a_{j2} b_{2j} + \cdots + a_{jj} b_{jj} &= 1 \end{aligned}$$

We forward solve, using the fact  $A$  has positive diagonal entries.

- The  $i = 1$  equation gives  $b_{1j} = 0$ .
- Substituting  $b_{1j} = 0$  into the  $i = 2$  equation gives  $a_{22} b_{2j} = 0$ , hence  $b_{2j} = 0$ .
- Substituting  $b_{1j} = b_{2j} = 0$  into the  $i = 3$  equation gives  $a_{33} b_{3j} = 0$ , hence  $b_{3j} = 0$ .
- Proceed in a similar fashion to get  $b_{ij} = 0$  for all  $i < j$ .
- Substituting  $b_{ij} = 0$  for all  $i < j$  into the  $i = j$  equation gives  $a_{jj} b_{jj} = 1$ , hence  $b_{jj} = \frac{1}{a_{jj}} > 0$ .

- (b) Assume that  $A$  has two Cholesky decompositions,

$$A = LL^T = MM^T, \quad L, M \in \mathcal{L}$$

Then

$$(M^{-1}L)(M^{-1}L)^T = M^{-1}LL^T M^{-T} = M^{-1}MM^T M^{-T} = I$$

hence  $U := M^{-1}L$  is orthogonal. Since  $\mathcal{L}$  is a group wrt matrix multiplication and  $M, L \in \mathcal{L}$ , we have  $U \in \mathcal{L}$ . The only orthogonal lower triangular matrix with positive diagonal entries is the identity, giving  $M^{-1}L = I$ . We conclude  $L = M$ .

3. Code: <https://github.com/RokettoJanpu/scientific-computing-1-redux/blob/main/hw5.ipynb>

```
import numpy as np
import numpy.linalg as la

def cholesky(A):
    n,m=A.shape
    if n!=m:
        print('The matrix is not square.')
        return
    L = np.zeros((n,n))
    for j in range(n):
        aux = A[j,j] - np.sum(L[j,:j]**2)
        if aux<=0:
            print('The matrix is not positive definite.')
            return
        L[j,j] = np.sqrt(aux)
        for i in range(j+1,n):
            L[i,j] = (A[i,j] - L[i,:j]@L[j,:j])/L[j,j]
    return L

A=np.random.rand(100,100)
B=A+A.T
C=A.T@A

for X in [B,C]:
    print('A+A^T:' if (X==B).all() else 'A^TA:')
    evals = la.eigvals(X)
    print('Minimum eigenvalue:',np.min(evals))
    L=cholesky(X)
    if L is not None:
        print('Matrix is SPD.')
        L_numpy = la.cholesky(X)
        print('Error between algorithms:',la.norm(L-L_numpy))

A+A^T:
Minimum eigenvalue: -7.788187643262669
The matrix is not positive definite.
A^TA:
Minimum eigenvalue: 9.176001668332046e-05
Matrix is SPD.
Error between algorithms: 2.2256399230782706e-12
```