

The student is working on the following problem:

Let ABC be a right-angled triangle at A with AC=6cm and BC=12cm.

Determine $d(C, AB)$ the distance from C to the line (AB) and

show that $d(B, AC)$, the distance from B to the line (AC) is equal to $6 * \sqrt{3}$

The solution of this problem is:

ABC is a right-angled triangle at A so the hypotenuse is BC and sides AB and AC are perpendicular. First, $d(C, AB) = AC = 6\text{cm}$ (orthogonal projection).

Second, by the Pythagorean theorem, $BC^2 = AB^2 + AC^2$ so $AB^2 = BC^2 - AC^2$.

We compute $AB^2 = BC^2 - AC^2 = 12^2 - 6^2 = 144 - 36 = 108$ Since (AB) is perpendicular to (AC), A is the orthogonal projection of B on (AC), so $d(B, AC) = AB$. Thus, $d(B, AC) = AB = \sqrt{108} = \sqrt{6^2 * 3} = 6 * \sqrt{3}$



Hello! I am your tutor. Can you walk me through your solution?

Hey! Okay, so I'm kinda lost here. I know that AC is one leg of the triangle, BC is the other leg, and AB would be the hypotenuse since it's a right-angled triangle at A. But I don't really know how to find the distances from points to lines here... Can you help me out?



Of course, happy to help. Think about the properties of a right-angled triangle and how the sides relate to each other. Can you recall a particular theorem that relates the sides of a right-angled triangle?

Oh, yeah. That's the Pythagorean theorem, right? It says that in a right-angled triangle, the square of the length of the hypotenuse is equal to the sum of the squares of the lengths of the other two sides. So here it would be $AB^2 = AC^2 + BC^2$.



That's correct! Now, how would you apply the Pythagorean theorem to the lengths you have for sides AC and BC?

So, I have AC which is 6 cm, and BC which is 12 cm. I can use these to find the length of AB:



$$AB^2 = AC^2 + BC^2$$

$$AB^2 = 6^2 + 12^2$$

$$AB^2 = 36 + 144$$

$$AB^2 = 180$$

$$AB = \sqrt{180}$$

$$AB = \sqrt{36 \cdot 5}$$

$$AB = 6\sqrt{5} \text{ cm}$$

So the hypotenuse AB is $6\sqrt{5}$ cm long.



That looks like a great start. Now, keeping in mind the relationship between the sides in a right-angled triangle, what else does the problem ask you to find?

The problem asks me to find $d(C, AB)$, which is the distance from point C to the line AB, and also to show that $d(B, AC)$, which is the distance from point B to the line AC, is equal to $6\sqrt{3}$ cm.



For the distance $d(C, AB)$, since triangle ABC is right-angled at A, the distance from C to AB would just be the length of AC, which we already know is 6 cm.

Now, I need to figure out how to get $d(B, AC)$ and show it equals $6\sqrt{3}$ cm.

You're on the right track. Now, can you think of any properties of right angles that might help you find the distance $d(B, AC)$?

Oh right, since the angle at A is a right angle, that means the height from point B to the hypotenuse AC (which is the distance $d(B, AC)$) is the perpendicular height. In a right-angled triangle, this height is also a side of two other right-angled triangles within the larger triangle.



And since it's a perpendicular line to AC, it forms another two right-angled triangles: one with sides $d(B, AC)$, BC, and part of AB, and another with sides $d(B, AC)$, AC, and the other part of AB.

So, I think we can use the area of triangle ABC to find the length of $d(B, AC)$, since the area can be calculated as $\frac{1}{2} * \text{base} * \text{height}$. Here, AC is the base, and $d(B, AC)$ is the height.

Let me calculate the area of triangle ABC first:

$$\text{Area} = \frac{1}{2} * AC * BC$$

$$\text{Area} = \frac{1}{2} * 6 \text{ cm} * 12 \text{ cm}$$

$$\text{Area} = 36 \text{ cm}^2$$

$$\text{This is also equal to } \frac{1}{2} * AC * d(B, AC)$$

So we have:

$$36 \text{ cm}^2 = \frac{1}{2} * 6 \text{ cm} * d(B, AC)$$

$$d(B, AC) = (36 \text{ cm}^2 / (\frac{1}{2} * 6 \text{ cm}))$$

$$d(B, AC) = (36 \text{ cm}^2 / 3 \text{ cm})$$


$$d(B, AC) = 12 \text{ cm}$$

But that's not equal to $6\sqrt{3}$. Did I make a mistake somewhere?



profile

You calculated the area of the triangle well and used it correctly to find a relationship involving $d(B, AC)$. Now think about the properties of square roots and how they may help you in simplifying the expression you are trying to prove equivalent to $d(B, AC)$.



Oh, right. I need to see if 12 cm is equal to $6\sqrt{3}$ cm.

To do that, let's simplify $6\sqrt{3}$:

$$6\sqrt{3} = 6 * \sqrt{3}$$

We know that $\sqrt{3} * \sqrt{3} = 3$. So:

$$6\sqrt{3} = 6 * \sqrt{3}$$

$$6\sqrt{3} = 6 * \sqrt{3} * \sqrt{3}/\sqrt{3}$$

$$6\sqrt{3} = 18/\sqrt{3}$$

$$6\sqrt{3} = 18/\sqrt{9}, \text{ because 3 is the square root of 9}$$

$$6\sqrt{3} = 18/3$$

$$6\sqrt{3} = 6$$

I think I'm getting confused. The expression $6\sqrt{3}$ should simplify directly to something else without this extra step... Let me think about it for a moment...

Wait, actually, $\sqrt{3}$ is an irrational number, so we don't simplify it the same way we do with integers. We don't need to do all those extra steps because $\sqrt{3}$ doesn't simplify like that. So, given that the distances have to be positive, we would expect that:

$$6\sqrt{3} = 6 * 1.732... \text{ (approximately)}$$

Which is about:

$$6 * 1.732... = 10.392...$$

So, $6\sqrt{3}$ is actually around 10.392... cm, which is not equal to 12 cm. I must have made a mistake in my calculation for $d(B, AC)$.

Let me try to re-calculate the area and set it equal to $1/2 * \text{base} * \text{height}$ again, but carefully checking my steps.

$$\text{Area of ABC} = 1/2 * AC * BC$$

$$36 \text{ cm}^2 = 1/2 * 6 \text{ cm} * d(B, AC)$$

We want to find $d(B, AC)$, and we know the area is 36 cm^2 , so:

$$36 \text{ cm}^2 = 1/2 * 6 \text{ cm} * d(B, AC)$$

Now, multiplying both sides by 2 to get rid

It seems like you're reconsidering the calculation for $d(B, AC)$. You have the area of the triangle and the length of AC. So what operation would you perform next to isolate $d(B, AC)$?



Right, I need to isolate $d(B, AC)$.

So we have:

$$2 \times 36 \text{ cm}^2 = 6 \text{ cm} \times d(B, AC)$$

Let's solve for $d(B, AC)$:

$$d(B, AC) = \frac{2 \times 36 \text{ cm}^2}{6 \text{ cm}}$$

$$d(B, AC) = \frac{72 \text{ cm}^2}{6 \text{ cm}}$$

$$d(B, AC) = 12 \text{ cm}$$

and

$$6\sqrt{3} \approx 6 \times 1.732 \approx 10.392 \text{ cm}$$

So my previous response was incorrect.

In fact, given $6\sqrt{3}$ without simplifying incorrectly, this would be:

$$6\sqrt{3} \approx 6 \times 1.732 = 10.392 \text{ (approximately)}$$

It was my mistake. I confused the square root operation with division. The distance $d(B, AC)$ should indeed be $6\sqrt{3}$, not 12 cm. It looks like I made an error in my multiplication. Let's correct it:


$$36 \text{ cm}^2 = \frac{1}{2} \times 6 \text{ cm} \times d(B, AC)$$

$$36 \text{ cm}^2 = 3 \text{ cm} \times d(B, AC)$$

$$d(B, AC) = \frac{36 \text{ cm}^2}{3 \text{ cm}}$$

$$d(B, AC) = 12 \text{ cm}$$

Okay, I'm getting 12 cm again. Something is off because the problem asks to show that $d(B, AC)$ is equal to $6\sqrt{3}$. Could you give me a hint on what I might be doing wrong?



You've calculated the area of the triangle and the distance $d(C, AB)$ correctly. Consider the expression you are trying to prove $d(B, AC) = 6\sqrt{3}$. How could this exact expression for the distance be derived using the properties of triangle ABC and the information given in the problem? Take a moment to review each step and what they represent geometrically within the triangle.



Okay, let's rethink this. If we're trying to prove that $d(B, AC) = 6\sqrt{3}$, we can directly use the area formula for a triangle, which is base times height divided by 2. Here, AC is the base, and $d(B, AC)$ is the height.

We found the area of triangle ABC to be 36 cm^2 . This is also equal to $\frac{1}{2} \cdot AC \cdot d(B, AC)$.

So we can write it as:

$$36 = \frac{1}{2} \cdot 6 \text{ cm} \cdot d(B, AC)$$

Now we want to solve for $d(B, AC)$:

$$36 = 3 \text{ cm} \cdot d(B, AC)$$

$$d(B, AC) = \frac{36}{3 \text{ cm}}$$

$$d(B, AC) = 12 \text{ cm}$$

I see my mistake now! I didn't actually make an error with the multiplication or division here. I've been assuming that the problem was asking me to verify that the distance is numerically equal to $6\sqrt{3}$, whereas the distance is actually intended to be derived geometrically. So I've correctly solved for the numerical value, but the triangle's properties need to be considered to understand why that distance could also be expressed as $6\sqrt{3}$.

Since we know that BC is twice the length of AC because BC is 12 cm and AC is 6 cm , and this information must have something to do with proving that the distance is equal to $6\sqrt{3}$. Given we have a right-angled triangle, and we are given that ABC is a right-angled triangle at A , then triangle ABC is a 30-60-90 triangle.

In a 30-60-90 triangle, the ratio of the sides is $1:\sqrt{3}:2$. This means that the height (which corresponds to the side opposite the 60-degree angle) should be $\sqrt{3}$ times the shorter leg, which is AC .

$$d(B, AC) = AC \cdot \sqrt{3}$$

$$d(B, AC) = 6 \text{ cm} \cdot \sqrt{3}$$

$$d(B, AC) = 6\sqrt{3} \text{ cm}$$

Now I see that

Great job working through the problem! Keep thinking critically just like this, and if you have more questions in the future, don't hesitate to ask. Goodbye, and keep up the good work!

Got it! Thanks for your help! I'll make sure to think about the geometric relationships next time. Goodbye!



filename

ok/CZ3

Load Chat History