

Confidence envelopes for the false discoveries with heterogeneous data

Romain Périer, joint work with Gilles Blanchard, Sebastian Döhler, Guillermo Durand and Étienne Roquain.



Multiple testing setting

Formal setting of multiple testing - notations

Let $X \in \mathcal{X}$ be a sample of distribution $\mu \in \mathcal{P}$ where $\mathcal{P}$ is a set of probability measures. Define :

- $H_1, \dots, H_m \subset \mathcal{P}$, m null hypotheses,
- For all i , a p -value $p_i(X)$ such that, if $\mu \in H_i$, $F_i(\alpha) := \mathbb{P}_{X \sim \mu}(p_i \leq \alpha) \leq \alpha$ for all $\alpha \in [0, 1]$,

Goal : find a "good" rejection set $R(X)$ under some error constraints.

Notations :

- $\mathcal{H}_0(\mu) = \{i : \mu \in H_i\}$,
- $m_0 = |\mathcal{H}_0(\mu)|$,
- $V(R) = |\mathcal{H}_0(\mu) \cap R|$.

Let R be a rejection set. Then we define the Family-Wise Error Rate (FWER) as

$$\text{FWER}(R) = \mathbb{P}(V(R) > 0).$$

Example 1 (Bonferonni procedure)

Let $\alpha \in [0, 1]$ and $R^{\text{Bonf}} = \{i \in \llbracket 1, m \rrbracket : p_i \leq \frac{\alpha}{m}\}$. Then $\text{FWER}(R^{\text{Bonf}}) \leq \alpha$.

We define the False Discovery Rate (FDR) as the mean of the False Discovery Proportion (FDP):

$$\text{FDP}(R) = \frac{V(R)}{|R| \vee 1}, \quad \text{FDR}(R) = \mathbb{E}[\text{FDP}(R)].$$

Example 2 (Benjamini-Hochberg procedure (Benjamini and Hochberg, 1995))

Let $\alpha \in [0, 1]$, $\hat{k}^{\text{BH}} = \max\{k : p_{(k)} \leq \frac{\alpha k}{m}\}$ and $R^{\text{BH}} = \{i \in \llbracket 1, m \rrbracket : p_i \leq \frac{\alpha \hat{k}}{m}\}$. Then, if $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values and is independent of $(p_i)_{i \in \mathcal{H}_1}$, $\text{FDR}(R^{\text{BH}}) \leq \alpha$.


Binomial tests : Assume $X = (X_1, \dots, X_m)$, where $(X_i)_{1 \leq i \leq m}$ are independent variables such that $X_i \sim \mathcal{B}(n_i, \pi_i)$ under H_i . Then :

- $p_i(X_i) := \mathbb{P}_{Y \sim \mathcal{B}(n_i, \pi_i)}(Y \leq X_i) = \sum_{j=0}^{X_i} \binom{n_i}{j} \pi_i^j (1 - \pi_i)^{n_i-j}$ is a p -value for i ,
- For all i , p_i is a discrete p -value and if the couples (n_i, π_i) are different, the respective cdf F_i are.

Other examples:

- Fisher's exact tests,
- Genome-wide association study.

CDFs of discrete and uniform p -values



Confidence envelopes

Definition 1 (Confidence envelope)

Let $\alpha \in (0, 1)$. A confidence envelope of level α is a statistic $\hat{V}_\alpha : \mathcal{P}(\mathbb{N}_m) \rightarrow \mathbb{N}$ such that, for all $\mu \in \mathcal{P}$,

$$\mathbb{P}_{X \sim \mu}(\forall S \subset \mathbb{N}_m, V(S) \leq \hat{V}_\alpha(S)) \geq 1 - \alpha.$$

- Allows user to choose freely their rejection set.

Definition 2 (Joint Error Rate)

For a reference family $\mathfrak{R} = ((R_k, \zeta_k))_{1 \leq k \leq K}$ and $\mu \in \mathcal{P}$, the joint error rate is :

$$JER(\mathfrak{R}, \mu) := \mathbb{P}_{X \sim \mu}(\exists k \in \mathbb{N}_K : V(R_k) > \zeta_k).$$

- Confidence envelope ideas restricted at our reference regions,
- Allows to compute a valid confidence envelope with interpolation (Blanchard et al., 2020, Durand et al., 2020).

In the following, we will be in a **top- k setting** :

$$R_k = \{\sigma(i), 1 \leq i \leq k\} \text{ (where } p_{\sigma(1)} \leq \dots \leq p_{\sigma(m)}).$$

Top-k confidence envelopes

Top-k confidence envelopes

Bretagnolle's inequality

Proposition 1 (DKWM inequality - Improved version from Reeve, 2024)

Let $X_1, \dots, X_m \stackrel{iid}{\sim} F$ and $\hat{F}_m(t) := \frac{1}{m} \sum_{1 \leq i \leq m} \mathbb{1}_{X_i \leq t}$ for all $t \in \mathbb{S}$.

Then, for all $\lambda \geq 0$,

$$\mathbb{P}\left(\sqrt{m} \sup_{t \in \mathbb{R}} (\hat{F}_m - F)(t) > \lambda\right) \leq \exp(-2\lambda^2).$$

When $(p_i)_{1 \leq i \leq m} \stackrel{iid}{\sim} \mathcal{U}([0, 1])$, $F(t) = t$.

Proposition 2 (Bretagnolle's inequality, Shorack and Wellner, 1986 with the improvement from Reeve, 2024)

Let $X_1, \dots, X_m$ be independent, denote by F_i the cdf of X_i and $\bar{F}$ the mean of F_i 's.

Then, for all $\lambda \geq 0$,

$$\mathbb{P}\left(\sqrt{m} \sup_{t \in \mathbb{R}} (\hat{F}_m - \bar{F})(t) > \lambda\right) \leq e \cdot \exp(-2\lambda^2).$$

Theorem 1 (Adaptive version of Bretagnolle's top- k calibration)

Let $\alpha \in (0, 1]$ and $\mathbb{G}(t) = (\mathbb{1}_{\{p_i \leq t\}} - F_i(t))_{1 \leq i \leq m}$. Suppose that $(p_i)_{i \in \mathcal{H}_0}$ is an independent family of p -values. Define $\lambda_\alpha = \sqrt{\frac{1 + \log(1/\alpha)}{2}}$, $\hat{m}_0$ such that

$$\left\{ \sup_{t \in [0, 1]} \sum_{i \in \mathcal{H}_0} \mathbb{G}_i(t) \leq \sqrt{m_0} \lambda_\alpha \right\} \subset \{m_0 \leq \hat{m}_0\}, \text{ and}$$

$$\zeta_k = k \wedge \left\lfloor \sum_{1 \leq i \leq \hat{m}_0} \left(\mathbf{F}(p_{(k)}) \right)_{[i]} + \sqrt{\hat{m}_0} \lambda_\alpha \right\rfloor^1.$$

Then, the reference family $\mathfrak{R} = ((R_k, \zeta_k))_{k \in \mathcal{K}}$ has its JER bounded by α .

Sketch of proof:

- $|R_k \cap \mathcal{H}_0| \leq \sum_{i \in \mathcal{H}_0} \mathbb{1}_{p_i \leq p_{(k)}}$,
- Evaluate Bretagnolle's inequality at point $p_{(k)}$,
- Bound $\sqrt{m_0}$ by $\sqrt{\hat{m}_0}$ and $\sum_{i \in \mathcal{H}_0} F_i(p_{(k)})$ by $\sum_{1 \leq i \leq \hat{m}_0} \left(\mathbb{F}(p_{(k)}) \right)_{[i]}$ (worst case).

¹ $x_{[i]}$ is the i -th largest coordinate of x .


Corollary 1 (Bretagnolle's top- k calibration)

Let $\alpha \in (0, 1]$. Suppose that $(p_i)_{i \in \mathcal{H}_0}$ is an independent family of p -values. Define


$$\zeta_k = k \wedge \left\lfloor m\bar{F}(p_{(k)}) + \sqrt{m}\lambda_\alpha \right\rfloor.$$

Then, the reference family $\mathfrak{R} = ((R_k, \zeta_k))_{k \in \mathcal{K}}$ has its JER bounded by α .


Comparison of bounds efficiency



Comparison of bounds efficiency



Comparison of confidence envelopes



Top-k confidence envelopes

Local test and estimation of m_0

Let $\alpha \in [0, 1]$ and, for $A \subset \{1, \dots, m\}$, $H_A := \bigcap_{i \in A} H_i$ the intersection hypothesis for A .

Suppose to have at hand, for all $A \subset \{1, \dots, m\}$, a local test φ_A (a test for H_A), and denote $\mathcal{N}(X) = \{A \subset \mathbb{N}_m : \varphi_A(X) = 0\}$.
 $\mathcal{N}(X)$ is the set of potential candidate for $\mathcal{H}_0(\mu)$.

Proposition 3 (Inversion procedure, Genovese and Wasserman, 2006)

Suppose that $X \sim \mu$ and $\varphi_{\mathcal{H}_0(\mu)}$ is a test of level α . Then, the statistic $\hat{V}_{\mathcal{N}}$ defined by $\hat{V}_{\mathcal{N}}(S) := \max_{A \in \mathcal{N}} |A \cap S|$ is a confidence envelope of level α .

- **Advantage** : Best bound on $V(S)$ from local tests,
- **Drawback** : Exponential computational cost.

Inversion procedure and concentration inequalities

Suppose that for all $A \subset \{1, \dots, m\}$, $\varphi_A = \sup_{t \in [0,1]} \varphi_{A,t}$.

Example 3

Let $\alpha \in (0, 1]$ and $A \subset \llbracket 1, m \rrbracket$. From Bretagnolle's inequality,

$$\varphi_{A,t} = \mathbb{1} \left\{ \sqrt{|A|} (\hat{F}_A - \bar{F}_A)(t) > \lambda_\alpha \right\} = \mathbb{1} \left\{ \sum_{i \in A} \mathbb{G}_i(t) > \sqrt{|A|} \lambda_\alpha \right\},$$

where $\mathbb{G}(t) = (\mathbb{1}_{\{p_i \leq t\}} - F_i(t))_{1 \leq i \leq m}$, is a local test of level α .

Lemma 1

For all $S \subset \mathbb{N}_m$,

$$\hat{V}_{\mathcal{N}}(S) = \max \left\{ \ell \in \{0, \dots, |S|\} : \min_{A: |A \cap S| = \ell} \sup_{t \in [0,1]} \varphi_{A,t} = 0 \right\}.$$

Shortcuts and application to Bretagnolle's inequality

Lemma 2

Denote by $\varphi_{S,\ell}^{SC1} := \max_{t \in [0,1]} \min_{A: |A \cap S| = \ell} \varphi_{A,t}$, then :

$$\begin{aligned} \hat{V}_{\mathcal{N}}(S) &\leq \max \left\{ \ell \in \{0, \dots, |S|\} : \varphi_{S,\ell}^{SC1} = 0 \right\} \\ &\leq \min_{t \in [0,1]} \left\{ \max \left\{ \ell \in \{0, \dots, |S|\} : \min_{A: |A \cap S| = \ell} \varphi_{A,t} = 0 \right\} \right\}. \end{aligned}$$

Example 4

With Bretagnolle's local test, for $t \in [0, 1]$,

$$\min_{A: |A \cap S| = \ell} \varphi_{A,t} = \min_{0 \leq j \leq m - |S|} \mathbb{1} \left\{ \sum_{1 \leq i \leq \ell} (\mathbb{G}(t))_{(S:i)} + \sum_{1 \leq i \leq j} (\mathbb{G}(t))_{(S^c:i)} > \lambda_{\alpha} \sqrt{\ell + j} \right\}.$$

- Reduces computation cost to $\mathcal{O}(m^2 \log(m))$.

Upper bound on m_0 via Bretagnolle's inequality

Theorem 2 (Bretagnolle's upper bound on m_0)

Let $\alpha \in (0, 1]$. Suppose that for all $k \in \mathcal{K}$, $(p_i)_{i \in \mathcal{H}_0}$ is an independent family of p -values.

Define $\lambda_\alpha = \sqrt{\frac{1 + \log(1/\alpha)}{2}}$, and

$$b(t) = \max \left\{ \ell : \ell^{-1/2} \left(\sum_{1 \leq j \leq \ell} (\mathbb{G}(t))_{(j)} \right) \leq \lambda_\alpha \right\}, \text{ for all } t \in [0, 1].$$

Then, within the event $\{\mathcal{H}_0 \in \mathcal{N}\}$, we have $m_0 \leq \min_{t \in [0, 1]} b(t) := \hat{m}_0$.

Sketch of proof :

- Use Example 4 with $S = \{1, \dots, m\}$,
- Apply lemma 2, second inequality.

Top-k confidence envelopes

Adaptive calibration

Theorem 3 (Adaptive version of Bretagnolle's top- k calibration)


Let $\alpha \in (0, 1]$. Suppose that $(p_i)_{i \in \mathcal{H}_0}$ is an independent family of p -values.

Define $\lambda_\alpha = \sqrt{\frac{1 + \log(1/\alpha)}{2}}$, $\mathbb{F}(t) := (F_i(t))_{1 \leq i \leq m}$, $\hat{m}_0$ as the estimator of m_0 from Theorem 2 and

$$\zeta_k = k \wedge \left\lfloor \sum_{1 \leq i \leq \hat{m}_0} \left(\mathbb{F}(p_{(k)}) \right)_{[i]} + \sqrt{\hat{m}_0} \lambda_\alpha \right\rfloor^2.$$


Then, the reference family $\mathfrak{R} = (R_k, \zeta_k)_{1 \leq k \leq m}$ has its JER bounded by α .

$^2x_{[i]}$ is the i -th largest coordinate of x .



Simulations (2/2)

Comparison of confidence envelopes



Heterogeneous simultaneous k -FWER

Heterogeneous simultaneous k -FWER

Come back to homogeneous setting

Derive a simultaneous k -FWER inequality from a FDR control procedure

Example 5 (Simes inequality)

Let $\alpha \in [0, 1]$. Assume that $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values. Then

$$\mathbb{P}\left(\exists i \in \{1, \dots, m_0\} : p_{(i:\mathcal{H}_0)} \leq \frac{\alpha i}{m}\right) \leq \alpha.$$

Derivation from BH procedure:

$$\begin{aligned}\mathbb{P}\left(\exists i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} \leq \frac{\alpha i}{m_0}\right) &= \mathbb{P}\left(\left|R^{\text{BH}}((p_i)_{i \in \mathcal{H}_0})\right| > 0\right) \\ &= \text{FWER}\left(R^{\text{BH}}((p_i)_{i \in \mathcal{H}_0})\right) \text{ since each } i \in \mathcal{H}_0 \text{ is true,} \\ &= \text{FDR}\left(R^{\text{BH}}((p_i)_{i \in \mathcal{H}_0})\right) \\ &\leq \alpha, \text{ by BH procedure.}\end{aligned}$$

Implies $JER(\mathfrak{R}) \leq \alpha$ for $R_k = \{i : p_i \leq \frac{\alpha k}{m}\}$ and $\zeta_k = k - 1$.

Definition 3

Let $\tau = (\tau_i)_{i \in \llbracket 1, m \rrbracket}$ be a family of thresholds.

Then the step up procedure associated to τ return

$$R^{\text{SU}(\tau)} = \{i \in \llbracket 1, m \rrbracket : p_i \leq \tau_{\hat{k}}\} \text{ and } \hat{k}^{\text{SU}(\tau)} = \max\{k : p_{(k)} \leq \tau_k\}.$$

The BH procedure is a **step-up** procedure.

Heterogeneous simultaneous k -FWER

Heterogeneous k -FWER method

A heterogeneous FDR control procedure

Theorem 4 (Döhler et al. (2018), Theorem 1)

Assume that $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values. Furthermore, assume that the two families $(p_i)_{i \in \mathcal{H}_0}$ and $(p_i)_{i \in \mathcal{H}_1}$ are independent. Let τ be a threshold family such that $\tau_m < 1$. Then

$$\text{FDR} \left(R^{\text{SU}(\tau)} \right) \leq \max_{1 \leq k \leq m} \left\{ \max_{\substack{A \subset \llbracket 1, m \rrbracket \\ |A| = m - k + 1}} \left\{ \frac{1}{k} \sum_{i \in A} \frac{F_i(\tau_k)}{1 - F_i(\tau_m)} \right\} \right\}^{(*)}.$$

Let, for all $(s, t) \in [0, 1) \times [0, 1]$,

$$\mathbf{H}(s, t) = \left(\frac{F_i(t)}{1 - F_i(s)} \right)_{1 \leq i \leq m},$$

for $i \in \llbracket 1, m \rrbracket$, $\mathcal{A}_i$ be the support of p_i and

$$\mathcal{A} = \{0\} \bigcup_{i \in \llbracket 1, m \rrbracket} \mathcal{A}_i.$$

Theorem 5 (Heterogeneous Simes-like inequality)

Let τ be a threshold family satisfying $(\star) \leq \alpha$, and assume that $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values. Then, the following inequality holds :

$$\mathbb{P}(\exists i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} \leq \tau_i) \leq \alpha.$$

Example 6 (Heterogeneous step-up procedure)

Let $\alpha \in (0, 1)$. Set


$$\tau_m \in \left\{ t \in \mathcal{A} : t < 1 \text{ and } (\mathbf{H}(t, t))_{[1]} \leq m\alpha \right\},$$

and, for all $k \in \llbracket 1, m-1 \rrbracket$, let


$$\tau_k = \max \left\{ t \in \mathcal{A}, t \leq \tau_m : (\mathbf{H}(\tau_m, t))_{[1]} + \cdots + (\mathbf{H}(\tau_m, t))_{[m-k+1]} \leq k\alpha \right\}.$$

Simulations (1/2)

Comparison of bounds efficiency



Comparison of bounds efficiency



Heterogeneous simultaneous k -FWER

**Adaptive (in m_0) heterogeneous k -FWER
method**

Proposition 4

From the proof of Theorem 4 in Döhler et al. (2018) the following slightly more refined result can be derived:

$$\text{FDR} \left(R^{\text{SU}(\tau)} \right) \leq \max_{1 \leq k \leq m} \left\{ \max_{\substack{A \subset \llbracket 1, m \rrbracket \\ |A| = m - k + 1}} \left\{ \frac{1}{k} \sum_{i \in A \cap \mathcal{H}_0} \frac{F_i(\tau_k)}{1 - F_i(\tau_m)} \right\} \right\}^{(*)}.$$

Example 7

Let $\alpha \in (0, 1)$ and τ_m under the same constraints than in Example 6 and, for all $k \in \llbracket 1, m - 1 \rrbracket$, let

$$\tilde{\tau}_k = \max \left\{ t \in \mathcal{A}, t \leq \tilde{\tau}_m : \left(\mathbf{H}(\tilde{\tau}_m, t) \right)_{[1]} + \cdots + \left(\mathbf{H}(\tilde{\tau}_m, t) \right)_{[(m-k+1) \wedge m_0 : \mathcal{H}_0]} \leq k\alpha \right\}.$$

Then $\tilde{\tau}$ satisfies $(\star) \leq \alpha$.

Theorem 6 ($\hat{m}_0$ -adaptivity of a threshold family)

Let $\tilde{\tau}$ be a threshold family such that $(\star) \leq \alpha$. Furthermore, assume that $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values. Then, for all (possibly data-dependant) threshold family τ such that $\tau_k \leq \tilde{\tau}_k$ for all $k \in \llbracket 1, m \rrbracket$ under the event $\left\{ \forall i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} > \tilde{\tau}_i \right\}$, the following inequality holds :

$$\mathbb{P}(\exists i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} \leq \tau_i) \leq \alpha.$$

Corollary 2 (Heterogeneous Simes like adaptive inequality)

Let $\tilde{\tau}$ be defined by Example 7 and $\hat{m}_0$ be a statistic such that

$$\{\forall i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} > \tilde{\tau}_i\} \subset \{m_0 \leq \hat{m}_0\}.$$

Furthermore, assume that $(p_i)_{i \in \mathcal{H}_0}$ is a family of independent p -values. Then the thresholds family defined by $\tau_m = \tilde{\tau}_m$ and $\forall k \in \llbracket 1, m-1 \rrbracket$,


$$\tau_k = \max \left\{ t \in \mathcal{A}, t \leq \tau_m : (\mathbf{H}(\tau_m, t))_{[1]} + \cdots + (\mathbf{H}(\tau_m, t))_{[(m-k+1) \wedge \hat{m}_0]} \leq k\alpha \right\}$$

verifies the inequality


$$\mathbb{P}(\exists i \in \mathcal{H}_0, p_{(i:\mathcal{H}_0)} \leq \tau_i) \leq \alpha.$$

Comparison with existing literature


Bound efficiency comparisons (1/2)



Bound efficiency comparisons (2/2)




Envelope comparisons (1/2)



Envelopes comparisons (2/2)

Comparison of confidence envelopes



Summary :

- Applied a heterogeneous version of DKWM inequality in the context of post hoc MT,
- Get an inexact shortcut for the inversion procedure practically applicable to heterogeneous inequalities,
- Get an m_0 adaptive version of the resulting confidence envelope,
- Get a heterogeneous Simes-like inequality and its adaptive version.

Perspectives :

- Numerical study of hybrid bounds,
- Get suitable heterogeneous versions of Wellner and K&R inequalities (work in progress).

Thanks a lot for your attention!

References

-  Benjamini, Yoav and Yosef Hochberg (1995). “Controlling the false discovery rate: a practical and powerful approach to multiple testing”. In: *J. Roy. Statist. Soc. Ser. B* 57.1, pp. 289–300. ISSN: 0035-9246. URL: [http://links.jstor.org/sici?sici=0035-9246\(1995\)57:1%3C289:CTFDRA%3E2.0.CO;2-E&origin=MSN](http://links.jstor.org/sici?sici=0035-9246(1995)57:1%3C289:CTFDRA%3E2.0.CO;2-E&origin=MSN).
-  Blanchard, Gilles, Pierre Neuvial, and Etienne Roquain (2020). “Post hoc confidence bounds on false positives using reference families”. In: *Ann. Statist.* 48.3, pp. 1281–1303. ISSN: 0090-5364, 2168-8966. DOI: 10.1214/19-AOS1847. URL: <https://doi.org/10.1214/19-AOS1847>.
-  Döhler, Sebastian, Guillermo Durand, and Etienne Roquain (2018). “New FDR bounds for discrete and heterogeneous tests”. In: *Electron. J. Stat.* 12.1, pp. 1867–1900. ISSN: 1935-7524. DOI: 10.1214/18-EJS1441. URL: <https://doi.org/10.1214/18-EJS1441>.

-  Durand, Guillermo, Gilles Blanchard, Pierre Neuvial, and Etienne Roquain (2020). “Post hoc false positive control for structured hypotheses”. In: *Scand. J. Stat.* 47.4, pp. 1114–1148. ISSN: 0303-6898,1467-9469. DOI: 10.1111/sjos.12453. URL: <https://doi.org/10.1111/sjos.12453>.
-  Durand, Guillermo, Florian Junge, Sebastian Döhler, and Etienne Roquain (2019). *DiscreteFDR: An R package for controlling the false discovery rate for discrete test statistics*. arXiv: 1904.02054 [stat.CO]. URL: <https://arxiv.org/abs/1904.02054>.
-  Genovese, Christopher R. and Larry Wasserman (2006). “Exceedance control of the false discovery proportion”. In: *J. Amer. Statist. Assoc.* 101.476, pp. 1408–1417. ISSN: 0162-1459,1537-274X. DOI: 10.1198/016214506000000339. URL: <https://doi.org/10.1198/016214506000000339>.

-  Katsevich, Eugene and Aaditya Ramdas (2020). “Simultaneous high-probability bounds on the false discovery proportion in structured, regression and online settings”. In: *Ann. Statist.* 48.6, pp. 3465–3487. ISSN: 0090-5364,2168-8966. DOI: 10.1214/19-AOS1938. URL: <https://doi.org/10.1214/19-AOS1938>.
-  Meah, Iqraa, Gilles Blanchard, and Etienne Roquain (2024). “False discovery proportion envelopes with m -consistency”. In: *J. Mach. Learn. Res.* 25, Paper No. 270, 52. ISSN: 1532-4435,1533-7928.
-  Reeve, Henry W J (2024). *A short proof of the Dvoretzky–Kiefer–Wolfowitz–Massart inequality*. arXiv: 2403.16651 [math.PR].
-  Shorack, Galen R. and Jon A. Wellner (1986). *Empirical processes with applications to statistics*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, pp. xxxviii+938. ISBN: 0-471-86725-X.

Precisions on interpolated bounds

Proposition 5 (Interpolated bound, Blanchard et al., 2020)

Let $\mathfrak{R} = ((R_k, \zeta_k))_{1 \leq k \leq K}$ a family with JER bounded by $\alpha \in [0, 1]$ and

$$\mathcal{A}(\mathcal{R}) := \left\{ A \subset \mathbb{N}_m : \forall k \in \mathbb{N}_K, |R_k \cap A| \leq \zeta_k \right\}.$$

Then $V_{\mathfrak{R}}^*(S) := \max_{A \in \mathcal{A}(\mathcal{R})} |R \cap A|$ is a level α post hoc bound.

Proposition 6 (Interpolated bound nested setting)

Assume $(R_k)_k$ is a family of regions such that

$$\forall k \in \{1, \dots, K\}, R_k \subset R_{k+1}.$$

Then:

$$V_{\mathfrak{R}}^*(S) = \min_{1 \leq k \leq K} (\zeta_k + |S \setminus R_k|) \wedge |S|.$$

Complements on Uniform Wellner and K& R inequalities

Proposition 7

Let $\alpha \leq 0.31$. Then, in the top- k setting with p -values under the null independent,

$$\zeta_k := \left\lfloor \frac{\log(1/\alpha)}{\log(1 + \log(1/\alpha))} (1 + mp_{(k)}) \right\rfloor$$

gives a reference family with JER bounded by α .

Proposition 8 (Uniform version of Wellner inequality, Meah et al., 2024)

Let $U_1, \dots, U_m \stackrel{iid}{\sim} \mathcal{U}([0, 1])$ and $\kappa = \frac{\pi^2}{6}$. For all $\delta \in (0, 1]$,

$$\mathbb{P} \left(\forall t \in [0, 1], m^{-1} \sum_{1 \leq i \leq m} \mathbb{1}_{U_i \leq t} \leq th^{-1} \left(\frac{2 \log(\kappa/\delta) + 4 \log(1 + \log_2(1/t))}{mt} \right) \right) \geq 1 - \delta,$$


with $h(\lambda) = \lambda(\log \lambda - 1) + 1$.

Simulations settings

Same simulation settings than Döhler et al., 2018, with $N = 50$, $m = 2000$, $\pi_0 = 0.8$ and the response under H_1 simulated by binomials of parameter 0,25 against 0.1 under H_0 .

Deterministic setting simulations

Deterministic setting : the R_k are deterministic and have a forest structure, meaning that for all $(k, k') \in \{1, \dots, K\}^2$, $R_k \cap R_{k'} \in \{\emptyset, R_k, R_{k'}\}$.



Old simulations

- **top- k** : $m = 3200$ and $m_0 = 2400$. Continuous p -values sampled from a Gaussian model, with mean equal to 0 under H_0 and 3 under H_1 . Each p -value is discretized using a support of length 50 sampled from a uniform law.
- **k-FWER** : Same simulation setting with the support equal to 300.
- **Deterministic setting** : $m = 3200$ and $m_0 = 2800$, 8 active leaves of length 100 and each active leaf contains 50% of p -values under H_1 .

Comparison with bounds found in the literature

