

DTU



47201 Engineering thermodynamics

# Module 10

# Recap: Enthalpy, $C_v$ , and $C_p$

Variation of internal energy:

$$dU = \delta Q + \delta W$$

Boundary work

$$\delta W = -PdV$$

Enthalpy:  $H = U + PV$

$$dH = dU + d(PV) = dU + PdV + VdP$$

At constant volume

$$dU = \delta Q \longrightarrow C_V = \left( \frac{\partial U}{\partial T} \right)_V$$


At constant pressure:

$$dH = \delta Q \longrightarrow C_P = \left( \frac{\partial H}{\partial T} \right)_P$$

Heat capacity

$$C = \frac{\delta Q}{\delta T}$$

At a given temperature, which one  
has more internal energy?




or



# Reaction heat (const. volume)

$$\Delta u = q_V + \cancel{w}$$




The heat of reaction,  $q_V$ , is the quantity of heat associated with a chemical reaction occurring within the system.

If the reaction occurs at *constant temperature*, the heat of reaction is exchanged between the system and its surroundings.

# Reaction heat (const. volume)

$$\Delta u = q_V + \cancel{w}$$




The heat of reaction,  $q_V$ , is the quantity of heat associated with a chemical reaction occurring within the system.

If the reaction occurs at *constant temperature*, the heat of reaction is exchanged between the system and its surroundings.

exothermic reaction

$$q_V < 0$$




releases heat  
to the  
surroundings

produces a  
temperature increase

endothermic reaction

$$q_V > 0$$




absorbs heat  
from the  
surroundings

produces a  
temperature decrease

# Reaction heat (const. volume)

$$\Delta u = q_V + \cancel{w}$$




The heat of reaction,  $q_V$ , is the quantity of heat associated with a chemical reaction occurring within the system.

If the reaction occurs in an *isolated system*, it produces a change in the temperature of the system.

exothermic reaction

$$q_V < 0$$




reaction  
releases heat

produces a  
temperature increase

endothermic reaction

$$q_V > 0$$




reaction  
absorbs heat

produces a  
temperature decrease

# Hess's law



*The enthalpy of a chemical process is independent of the path taken from the initial to the final state*





# Hess's law




*The enthalpy of a chemical process is independent of the path taken from the initial to the final state*



$$\Delta H^\circ = ?$$





# Hess's law

*The enthalpy of a chemical process is independent of the path taken from the initial to the final state*






# Hess's law

*The enthalpy of a chemical process is independent of the path taken from the initial to the final state*



# Hess's law

*The enthalpy of a chemical process is independent of the path taken from the initial to the final state*



# Enthalpy of formation

*Change of enthalpy during the formation of 1 mole of the substance from its constituent elements in their reference form.*

The standard enthalpy of formation of a pure element in its reference form is 0.

*Reference form* means the molecules spontaneously formed by elements, e.g.  $\text{H}_2$

# Enthalpy of formation

*Change of enthalpy during the formation of 1 mole of the substance from its constituent elements in their reference form.*


The standard enthalpy of formation of a pure element in its reference form is 0.

*Reference form* means the molecules spontaneously formed by elements, e.g.  $\text{H}_2$

Why is this useful?

# Enthalpy of formation


*Change of enthalpy during the formation of 1 mole of the substance from its constituent elements in their reference form.*







$$\Delta H^\circ = ?$$

# Enthalpy of formation

## Decomposition



## Recombination





# Enthalpy of formation

[See wikipedia page](#)


## Inorganic substances [\[ edit \]](#)

Species	Phase	Chemical formula	$\Delta_f H^\ominus / (\text{kJ/mol})$
Aluminium	Solid	Al	0
Aluminium chloride	Solid	AlCl <sub>3</sub>	-705.63
Aluminium oxide	Solid	Al <sub>2</sub> O <sub>3</sub>	-1675.5
Aluminium hydroxide	Solid	Al(OH) <sub>3</sub>	-1277
Aluminium sulphate	Solid	Al <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub>	-3440
Barium chloride	Solid	BaCl <sub>2</sub>	-858.6
Barium carbonate	Solid	BaCO <sub>3</sub>	-1216
Barium hydroxide	Solid	Ba(OH) <sub>2</sub>	-944.7
Barium oxide	Solid	BaO	-548.1
Barium sulfate	Solid	BaSO <sub>4</sub>	-1473.3
Beryllium	Solid	Be	0
Beryllium hydroxide	Solid	Be(OH) <sub>2</sub>	-903
Beryllium oxide	Solid	BeO	-609.4
Boron trichloride	Solid	BCl <sub>3</sub>	-402.96
Bromine	Liquid	Br <sub>2</sub>	0
Bromide ion	Aqueous	Br <sup>-</sup>	-121

# Temperature dependence

*Standard* enthalpies of formation refer to standard pressure ( $10^5$  Pa) and specified temperature  $T_1$  (which may vary, but often is  $25\text{ }^\circ\text{C} = 298.15\text{ K}$ ).

How do we calculate the reaction enthalpy if the reaction happens at a different temperature  $T_2$ ?



# Temperature dependence


*Standard* enthalpies of formation refer to standard pressure ( $10^5$  Pa) and specified temperature  $T_1$  (which may vary, but often is  $25\text{ °C} = 298.15\text{ K}$ ).

How do we calculate the reaction enthalpy if the reaction happens at a different temperature  $T_2$ ?

We split it in steps and apply conservation of energy:

1. We bring the reactants from  $T_2$  to  $T_1$ :  $\Delta H = \sum_{\text{Reactants}} \int_{T_2}^{T_1} C_p(T) dT$
2. We perform the reaction at standard temperature  $T_1$
3. We bring the products back to  $T_2$ :  $\Delta H = \sum_{\text{Products}} \int_{T_1}^{T_2} C_p(T) dT$

We add the enthalpy variations for the three steps.



# Temperature dependence

*Standard* enthalpies of formation refer to standard pressure ( $10^5$  Pa) and specified temperature  $T_1$  (which may vary, but often is  $25\text{ }^\circ\text{C} = 298.15\text{ K}$ ).


How do we calculate the reaction enthalpy if the reaction happens at a different temperature  $T_2$ ?

We split it in steps and apply conservation of energy:

1. We bring the reactants from  $T_2$  to  $T_1$ :  $\Delta H = \sum_{\text{Reactants}} \int_{T_2}^{T_1} C_p(T) dT$
2. We perform the reaction at standard temperature  $T_1$
3. We bring the products back to  $T_2$ :  $\Delta H = \sum_{\text{Products}} \int_{T_1}^{T_2} C_p(T) dT$

We add the enthalpy variations for the three steps.

The result might be different due to the different heat capacities between reactants and products.



# Constant volume reactions

How do we calculate the heat of reaction at constant volume from the reaction enthalpy?

# Constant volume reactions

How do we calculate the heat of reaction at constant volume from the reaction enthalpy?

For a constant pressure reaction, the enthalpy is equal to the heat of reaction.

For obtaining the heat of reaction at constant pressure, we need to consider the work done against the external pressure.

$$q_V = q_P + w \quad \text{Is it clear why do we have a plus sign?}$$

# Constant volume reactions

How do we calculate the heat of reaction at constant volume from the reaction enthalpy?

For a constant pressure reaction, the enthalpy is equal to the heat of reaction.

For obtaining the heat of reaction at constant pressure, we need to consider the work done against the external pressure.

$$q_V = q_P + w \quad \text{Is it clear why do we have a plus sign?}$$

While at constant pressure some energy is used for boundary work, at constant volume all the energy is entirely converted into heat.

# Constant volume reactions

How do we calculate the heat of reaction at constant volume from the reaction enthalpy?

For a constant pressure reaction, the enthalpy is equal to the heat of reaction.

For obtaining the heat of reaction at constant pressure, we need to consider the work done against the external pressure.

$$q_V = q_P + w$$

To calculate the work, we need to know the pressure and specific volumes of all the reactants and products.

$$w = -P\Delta v \quad \text{work that would be done in const. P}$$



# Constant volume reactions

How do we calculate the heat of reaction at constant volume from the reaction enthalpy?

For a constant pressure reaction, the enthalpy is equal to the heat of reaction.

For obtaining the heat of reaction at constant pressure, we need to consider the work done against the external pressure.

$$q_V = q_P + w$$

To calculate the work, we need to know the pressure and specific volumes of all the reactants and products.

$$w = -P\Delta v \quad \text{work that would be done in const. P}$$

Sometimes, we just need to consider the gases (using the ideal gas law) since the volumes of solids and liquids are very small in comparison.

# Reactions happening at a different pressure

*Standard* enthalpies of formation refer to standard pressure ( $P_1 = 10^5$  Pa) and specified temperature  $T$ .

How do we calculate the reaction enthalpy (heat of reaction) if the reaction happens at a different pressure  $P_2$ ?

# Reactions happening at a different pressure

*Standard* enthalpies of formation refer to standard pressure ( $P_1 = 10^5$  Pa) and specified temperature  $T$ .

How do we calculate the reaction enthalpy (heat of reaction) if the reaction happens at a different pressure  $P_2$ ?

We need to consider again the work done against the external pressure.

$$q_V = q_{P_1} + w_1 = q_{P_2} + w_2$$

# Reactions happening at a different pressure

*Standard* enthalpies of formation refer to standard pressure ( $P_1 = 10^5$  Pa) and specified temperature  $T$ .

How do we calculate the reaction enthalpy (heat of reaction) if the reaction happens at a different pressure  $P_2$ ?

We need to consider again the work done against the external pressure.

$$q_V = q_{P_1} + w_1 = q_{P_2} + w_2 \rightarrow q_{P_2} = q_{P_1} + w_1 - w_2$$

# Reactions happening at a different pressure

*Standard* enthalpies of formation refer to standard pressure ( $P_1 = 10^5$  Pa) and specified temperature  $T$ .

How do we calculate the reaction enthalpy (heat of reaction) if the reaction happens at a different pressure  $P_2$ ?

We need to consider again the work done against the external pressure.

$$q_V = q_{P_1} + w_1 = q_{P_2} + w_2 \rightarrow q_{P_2} = q_{P_1} + w_1 - w_2$$

To calculate the work, we need to know the pressure and specific volumes of all the reactants and products (at the right pressure!).

$$w_1 = -P_1 \Delta v_1$$

$$w_2 = -P_2 \Delta v_2$$

# Conclusions

There is some book-keeping to be done, but ultimately it always boils down to conservation of energy.

One always has to remember that specific enthalpy and specific internal energy are state functions: for each substance they only depend on the temperature.

Since they are extensive properties, they are proportional to the amount of each substance, and we have to sum the contributions for each chemical compound.