

## Lecture 21. Solving Recurrence Relations

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

**Example 20.1.** Ahmed lends money at outrageous rates of interest. He demands to be paid 10% interest *per week* on a loan, compounded weekly. Suppose you borrow 500 Dhs from him. Let  $M(n)$  = the money you owed at  $n$ -th week.

- (a) Find the recurrence relation for  $M(n)$ .
- (b) If you wait four weeks to pay him back, how much will you owe?

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

**Example 20.1.** Ahmed lends money at outrageous rates of interest. He demands to be paid 10% interest *per week* on a loan, compounded weekly. Suppose you borrow 500 Dhs from him. Let  $M(n)$  = the money you owed at  $n$ -th week.

- (a) Find the recurrence relation for  $M(n)$ .
- (b) If you wait four weeks to pay him back, how much will you owe?

**Answer:**

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.1 M(n-1) & \text{if } n > 0 \end{cases} \quad M(4) = \$732.05$$

**Practice 20.2.** Find a closed-form solution for the recurrence relation from

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.1 M(n-1) & \text{if } n > 0 \end{cases}$$

**Practice 20.2.** Find a closed-form solution for the recurrence relation from

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.1 M(n-1) & \text{if } n > 0 \end{cases}$$

**Answer:**  $M(n) = 500(1.1)^n$

## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

**One Idea.** If we knew that

$$P(n) = \text{a polynomial function of } n$$

then we can detect such a sequence by looking at the differences between terms.



## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

**One Idea.** If we knew that

$$P(n) = \text{a polynomial function of } n$$

then we can detect such a sequence by looking at the differences between terms.  
Given any sequence,

$$a_0, a_1, a_2, \dots, a_{n-1}, a_n$$

the differences

$$a_1 - a_0, a_2 - a_1, a_3 - a_2, \dots, a_n - a_{n-1}$$

form another sequence, called the [sequence of differences](#).

Given any sequence,

$$a_0, a_1, a_2, \dots, a_{n-1}, a_n$$

the differences

$$a_1 - a_0, a_2 - a_1, a_3 - a_2, \dots, a_n - a_{n-1}$$

form another sequence, called the [sequence of differences](#).

- A linear sequence  $a_n = An + B$  will have a constant sequence of differences (because a line has constant slope).
- A quadratic sequence  $a_n = An^2 + Bn + C$  will have a linear sequence of differences.
- a cubic sequence  $a_n = An^3 + Bn^2 + Cn + D$  will have a quadratic sequence of differences, etc.
- If we eventually end up with a constant sequence, then the original sequence is given by a polynomial function.
- The degree of the conjectured polynomial is the number of times we had to calculate the sequence of differences.

**Example 20.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Example 20.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Example 20.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Practice 20.4.** Solve the recurrence relation  $a_n = a_{n-1} + n$  with initial term  $a_0 = 4$ .

**Example 20.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Practice 20.4.** Solve the recurrence relation  $a_n = a_{n-1} + n$  with initial term  $a_0 = 4$ .

**Answer.**  $a_n = \frac{n(n+1)}{2} + 4$  for  $n \geq 0$ .