

Lecture 22. Linear Homogeneous Recurrence Relations

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

- **Linear:** Each term a_j in the relation appears with an exponent of 1.

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

- **Linear:** Each term a_j in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on n (i.e., no external or independent functions of n).

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

- **Linear:** Each term a_j in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on n (i.e., no external or independent functions of n).
- **Degree k :** Each term a_n is defined by the previous k terms in the sequence.

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

- **Linear:** Each term a_j in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on n (i.e., no external or independent functions of n).
- **Degree k :** Each term a_n is defined by the previous k terms in the sequence.

Example 22.2. Linear and/or Homogeneous?

$$(a) M_n = (1.1)M_{n-1} \quad (b) F_n = F_{n-1} + F_{n-2} \quad (c) a_n = a_{n-1} + a_{n-2}^2 \quad (d) h_n = 2h_{n-1} + 1$$

Definition 22.1. A linear homogeneous recurrence relation of degree k with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where r_1, r_2, \dots, r_k are real numbers and $r_k \neq 0$ with $k < n$. This recurrence includes k initial conditions, $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$.

- **Linear:** Each term a_j in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on n (i.e., no external or independent functions of n).
- **Degree k :** Each term a_n is defined by the previous k terms in the sequence.

Example 22.2. Linear and/or Homogeneous?

$$(a) M_n = (1.1)M_{n-1} \quad (b) F_n = F_{n-1} + F_{n-2} \quad (c) a_n = a_{n-1} + a_{n-2}^2 \quad (d) h_n = 2h_{n-1} + 1$$

(a) linear homogeneous recurrence relation of degree 1, (b) linear homogeneous recurrence relation of degree 2, (c) not linear, (d) not homogeneous,

- The basic approach is to look for solutions of the form $a_n = x^n$, where x is a constant.

- The basic approach is to look for solutions of the form $a_n = x^n$, where x is a constant.
- $a_n = x^n$ is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

- The basic approach is to look for solutions of the form $a_n = x^n$, where x is a constant.
- $a_n = x^n$ is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

$$x^n = r_1 x^{n-1} + r_2 x^{n-2} + \cdots + r_k x^{n-k}$$

- The basic approach is to look for solutions of the form $a_n = x^n$, where x is a constant.
- $a_n = x^n$ is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

$$x^n = r_1 x^{n-1} + r_2 x^{n-2} + \cdots + r_k x^{n-k}$$

- Rearranging terms leads to the **characteristic equation**:

$$x^n - r_1 x^{n-1} - r_2 x^{n-2} - \cdots - r_k x^{n-k} = 0.$$

The characteristic equation:

$$x^n - r_1x^{n-1} - r_2x^{n-2} - \dots - r_kx^{n-k} = 0.$$

Theorem 22.3. The characteristic equation of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ is

$$x^2 - r_1x - r_2 = 0.$$

If the characteristic equation has two distinct roots, x_1 and x_2 , then

$$a_n = px_1^n + qx_2^n \text{ for some } p, q$$

is the explicit formula for the sequence. Here, p and q depend on the initial conditions.

Example 22.4. Find an explicit formula for the sequence defined by $a_n = 7a_{n-1} - 10a_{n-2}$ with $a_0 = 2$ and $a_1 = 3$.

The characteristic equation:

$$x^n - r_1x^{n-1} - r_2x^{n-2} - \dots - r_kx^{n-k} = 0.$$

Theorem 22.3. The characteristic equation of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ is

$$x^2 - r_1x - r_2 = 0.$$

If the characteristic equation has two distinct roots, x_1 and x_2 , then

$$a_n = px_1^n + qx_2^n \text{ for some } p, q$$

is the explicit formula for the sequence. Here, p and q depend on the initial conditions.

Example 22.4. Find an explicit formula for the sequence defined by $a_n = 7a_{n-1} - 10a_{n-2}$ with $a_0 = 2$ and $a_1 = 3$.

Answer: $a_n = \frac{7}{3}2^n - \frac{1}{3}5^n$.

Theorem 22.5. If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ has two distinct roots, x_1 and x_2 , then

$$a_n = px_1^n + qx_2^n$$

where p and q depend on the initial conditions, is the explicit formula for the sequence.

Practice 22.6. (Fibonacci Sequence) Solve the recurrence relation $f_n = f_{n-1} + f_{n-2}$ with the initial conditions $f_0 = 0$ and $f_1 = 1$.

Theorem 22.5. If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ has two distinct roots, x_1 and x_2 , then

$$a_n = px_1^n + qx_2^n$$

where p and q depend on the initial conditions, is the explicit formula for the sequence.

Practice 22.6. (Fibonacci Sequence) Solve the recurrence relation $f_n = f_{n-1} + f_{n-2}$ with the initial conditions $f_0 = 0$ and $f_1 = 1$.

Answer: $f_n = \frac{\phi^n - (1-\phi)^n}{\sqrt{5}}$, where $\phi = \frac{1+\sqrt{5}}{2}$. (Note that $\phi \approx 1.618\ 033\ 988\ 749\dots$ is a golden ratio.)

Theorem 22.7. If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ has a single root, x , then

$$a_n = px_1^n + qnx_2^n$$

where p and q depend on the initial conditions, is the explicit formula for the sequence.

Practice 22.8. Solve the recurrence relation $b_n = 6b_{n-1} - 9b_{n-2}$ with the initial conditions $b_0 = 1$ and $b_1 = 4$.

Theorem 22.7. If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation $a_n = r_1a_{n-1} + r_2a_{n-2}$ has a single root, x , then

$$a_n = px_1^n + qnx_2^n$$

where p and q depend on the initial conditions, is the explicit formula for the sequence.

Practice 22.8. Solve the recurrence relation $b_n = 6b_{n-1} - 9b_{n-2}$ with the initial conditions $b_0 = 1$ and $b_1 = 4$.

Answer: $b_n = 3^n + \frac{1}{3}n3^n$

Although we will not consider examples more complicated than these, this characteristic root technique can be applied to much more complicated recurrence relations.