

## Lecture 19. Solving Recurrence Relations

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

**Example 19.1.** Ahmed lends money at outrageous rates of interest. He demands to be paid 10% interest *per week* on a loan, compounded weekly. Suppose you borrow 500 Dhs from him. Let  $M(n)$  = the money you owed at  $n$ -th week.

- (a) Find the recurrence relation for  $M(n)$ .
- (b) If you wait four weeks to pay him back, how much will you owe?

## Recursive Thinking

In the previous lecture note, we have seen some examples how to think recursively about a problem by describing it with a recurrence relation. Remember that any recurrence relation has two parts: a **base case** that describes some initial conditions, and a **recursive case** that describes a **future value in terms of previous values**. Armed with this way of thinking, we can model other problems using recurrence relations.

**Example 19.1.** Ahmed lends money at outrageous rates of interest. He demands to be paid 10% interest *per week* on a loan, compounded weekly. Suppose you borrow 500 Dhs from him. Let  $M(n)$  = the money you owed at  $n$ -th week.

- (a) Find the recurrence relation for  $M(n)$ .
- (b) If you wait four weeks to pay him back, how much will you owe?

**Answer:**

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.10 M(n-1) & \text{if } n > 0 \end{cases} \quad M(4) = \$732.05$$

**Practice 19.2.** Find a closed-form solution for the recurrence relation from

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.10 M(n-1) & \text{if } n > 0 \end{cases}$$

**Practice 19.2.** Find a closed-form solution for the recurrence relation from

$$M(n) = \begin{cases} 500 & \text{if } n = 0 \\ 1.10 M(n-1) & \text{if } n > 0 \end{cases}$$

**Answer:**  $M(n) = 500(1.10)^n$

## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

**One Idea.** If we knew that

$$P(n) = \text{a polynomial function of } n$$

then we can detect such a sequence by looking at the differences between terms.



## Polynomial sequences: Using differences

Finding a closed-form solution for  $P(n)$  can be thought of as guessing a formula for the sequence  $P(0), P(1), P(2), \dots$

**One Idea.** If we knew that

$$P(n) = \text{a polynomial function of } n$$

then we can detect such a sequence by looking at the differences between terms.  
Given any sequence,

$$a_0, a_1, a_2, \dots, a_{n-1}, a_n$$

the differences

$$a_1 - a_0, a_2 - a_1, a_3 - a_2, \dots, a_n - a_{n-1}$$

form another sequence, called the [sequence of differences](#).

Given any sequence,

$$a_0, a_1, a_2, \dots, a_{n-1}, a_n$$

the differences

$$a_1 - a_0, a_2 - a_1, a_3 - a_2, \dots, a_n - a_{n-1}$$

form another sequence, called the [sequence of differences](#).

- A linear sequence  $a_n = An + B$  will have a constant sequence of differences (because a line has constant slope).
- A quadratic sequence  $a_n = An^2 + Bn + C$  will have a linear sequence of differences.
- a cubic sequence  $a_n = An^3 + Bn^2 + Cn + D$  will have a quadratic sequence of differences, etc.
- If we eventually end up with a constant sequence, then the original sequence is given by a polynomial function.
- The degree of the conjectured polynomial is the number of times we had to calculate the sequence of differences.

**Example 19.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Example 19.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Example 19.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Practice 19.4.** Solve the recurrence relation  $a_n = a_{n-1} + n$  with initial term  $a_0 = 4$ .

**Example 19.3.** Find a closed-form solution  $f(n)$  for the recurrence relation

$$H(n) = \begin{cases} 1 & \text{if } n = 1 \\ H(n-1) + 6n - 6 & \text{if } n > 1 \end{cases}$$

**Answer.**  $H(n) = 3n^2 - 3n + 1$  is a **good candidate** for a closed-form solution.

**Remark.** The result of these procedures is still only a guess. To be sure that our guess is right, we need to prove that the formula matches the recurrence relation **for all  $n$** .

**Practice 19.4.** Solve the recurrence relation  $a_n = a_{n-1} + n$  with initial term  $a_0 = 4$ .

**Answer.**  $a_n = \frac{n(n+1)}{2} + 4$  for  $n \geq 0$ .

## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

- **Linear:** Each term  $a_j$  in the relation appears with an exponent of 1.



## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

- **Linear:** Each term  $a_j$  in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on  $n$  (i.e., no external or independent functions of  $n$ ).

## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

- **Linear:** Each term  $a_j$  in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on  $n$  (i.e., no external or independent functions of  $n$ ).
- **Degree  $k$ :** Each term  $a_n$  is defined by the previous  $k$  terms in the sequence.

## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

- **Linear:** Each term  $a_j$  in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on  $n$  (i.e., no external or independent functions of  $n$ ).
- **Degree  $k$ :** Each term  $a_n$  is defined by the previous  $k$  terms in the sequence.

**Example 19.6.** Linear and/or Homogeneous?

$$(a) M_n = (1.1)M_{n-1} \quad (b) F_n = F_{n-1} + F_{n-2} \quad (c) a_n = a_{n-1} + a_{n-2}^2 \quad (d) h_n = 2h_{n-1} + 1$$

## Linear Homogeneous Recurrence Relations

**Definition 19.5.** A linear homogeneous recurrence relation of degree  $k$  with constant coefficients is a recurrence relation of the form

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

where  $r_1, r_2, \dots, r_k$  are real numbers and  $r_k \neq 0$  with  $k < n$ . This recurrence includes  $k$  initial conditions,  $a_0 = \alpha_0, a_1 = \alpha_1, \dots, a_k = \alpha_k$ .

- **Linear:** Each term  $a_j$  in the relation appears with an exponent of 1.
- **Homogeneous:** The relation contains no additional terms that directly depend on  $n$  (i.e., no external or independent functions of  $n$ ).
- **Degree  $k$ :** Each term  $a_n$  is defined by the previous  $k$  terms in the sequence.

**Example 19.6.** Linear and/or Homogeneous?

$$(a) M_n = (1.1)M_{n-1} \quad (b) F_n = F_{n-1} + F_{n-2} \quad (c) a_n = a_{n-1} + a_{n-2}^2 \quad (d) h_n = 2h_{n-1} + 1$$

(a) linear homogeneous recurrence relation of degree 1, (b) linear homogeneous recurrence relation of degree 2, (c) not linear, (d) not homogeneous,

- The basic approach is to look for solutions of the form  $a_n = x^n$ , where  $x$  is a constant.

- The basic approach is to look for solutions of the form  $a_n = x^n$ , where  $x$  is a constant.
- $a_n = x^n$  is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

- The basic approach is to look for solutions of the form  $a_n = x^n$ , where  $x$  is a constant.
- $a_n = x^n$  is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

$$x^n = r_1 x^{n-1} + r_2 x^{n-2} + \cdots + r_k x^{n-k}$$

- The basic approach is to look for solutions of the form  $a_n = x^n$ , where  $x$  is a constant.
- $a_n = x^n$  is a solution to the recurrence relation

$$a_n = r_1 a_{n-1} + r_2 a_{n-2} + \cdots + r_k a_{n-k}$$

if and only if

$$x^n = r_1 x^{n-1} + r_2 x^{n-2} + \cdots + r_k x^{n-k}$$

- Rearranging terms leads to the **characteristic equation**:

$$x^n - r_1 x^{n-1} - r_2 x^{n-2} - \cdots - r_k x^{n-k} = 0.$$



The characteristic equation:

$$x^n - r_1x^{n-1} - r_2x^{n-2} - \dots - r_kx^{n-k} = 0.$$

**Theorem 19.7.** The characteristic equation of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  is

$$x^2 - r_1x - r_2 = 0.$$

If the characteristic equation has two distinct roots,  $x_1$  and  $x_2$ , then

$$a_n = px_1^n + qx_2^n \text{ for some } p, q$$

is the explicit formula for the sequence. Here,  $p$  and  $q$  depend on the initial conditions.

**Example 19.8.** Find an explicit formula for the sequence defined by  $a_n = 7a_{n-1} - 10a_{n-2}$  with  $a_0 = 2$  and  $a_1 = 3$ .

The characteristic equation:

$$x^n - r_1x^{n-1} - r_2x^{n-2} - \dots - r_kx^{n-k} = 0.$$

**Theorem 19.7.** The characteristic equation of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  is

$$x^2 - r_1x - r_2 = 0.$$

If the characteristic equation has two distinct roots,  $x_1$  and  $x_2$ , then

$$a_n = px_1^n + qx_2^n \text{ for some } p, q$$

is the explicit formula for the sequence. Here,  $p$  and  $q$  depend on the initial conditions.

**Example 19.8.** Find an explicit formula for the sequence defined by  $a_n = 7a_{n-1} - 10a_{n-2}$  with  $a_0 = 2$  and  $a_1 = 3$ .

**Answer:**  $a_n = \frac{7}{3}2^n - \frac{1}{3}5^n$ .

**Theorem 19.7.** If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  has two distinct roots,  $x_1$  and  $x_2$ , then

$$a_n = px_1^n + qx_2^n$$

where  $p$  and  $q$  depend on the initial conditions, is the explicit formula for the sequence.

**Practice 19.9.** (Fibonacci Sequence) Solve the recurrence relation  $f_n = f_{n-1} + f_{n-2}$  with the initial conditions  $f_0 = 0$  and  $f_1 = 1$ .

**Theorem 19.7.** If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  has two distinct roots,  $x_1$  and  $x_2$ , then

$$a_n = px_1^n + qx_2^n$$

where  $p$  and  $q$  depend on the initial conditions, is the explicit formula for the sequence.

**Practice 19.9.** (Fibonacci Sequence) Solve the recurrence relation  $f_n = f_{n-1} + f_{n-2}$  with the initial conditions  $f_0 = 0$  and  $f_1 = 1$ .

**Answer:**  $f_n = \frac{\phi^n - (1-\phi)^n}{\sqrt{5}}$ , where  $\phi = \frac{1+\sqrt{5}}{2}$ . (Note that  $\phi \approx 1.618\ 033\ 988\ 749\dots$  is a golden ratio.)

**Theorem 19.10.** If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  has a single root,  $x$ , then

$$a_n = px_1^n + qnx_2^n$$

where  $p$  and  $q$  depend on the initial conditions, is the explicit formula for the sequence.

**Practice 19.11.** Solve the recurrence relation  $b_n = 6b_{n-1} - 9b_{n-2}$  with the initial conditions  $b_0 = 1$  and  $b_1 = 4$ .

**Theorem 19.10.** If the characteristic equation

$$x^2 - r_1x - r_2 = 0$$

of the recurrence relation  $a_n = r_1a_{n-1} + r_2a_{n-2}$  has a single root,  $x$ , then

$$a_n = px_1^n + qnx_2^n$$

where  $p$  and  $q$  depend on the initial conditions, is the explicit formula for the sequence.

**Practice 19.11.** Solve the recurrence relation  $b_n = 6b_{n-1} - 9b_{n-2}$  with the initial conditions  $b_0 = 1$  and  $b_1 = 4$ .

**Answer:**  $b_n = 3^n + \frac{1}{3}n3^n$

Although we will not consider examples more complicated than these, this characteristic root technique can be applied to much more complicated recurrence relations.