135

05

23738
44157.2
744235
10997700

SMP

10997700

MATH1061

work submission coversheet

Submit via Blackboard Course Link

School of Mathematics and Physics
Faculty of Science

This cover sheet should be attached to the front of your submission (not behind plastic). It is important that each barcode on this page is read. Make sure that the barcode sections are devoid of ink or stains etc.

Last Name	**Lin**	Student Number	**45791439**
First Name	Xianglong		
Submission Topic	Assignment 2		

Due date of submission	10/09/2020		
Due time of submission	1:00 PM	Name of tutor (if appropriate)	
Group	05	Session attended (if appropriate)	
Actual submission date		day	time

Marked

Yes / No

IMPORTANT POINTS:

- **Place the submission through the slot with the barcode FACE UP.**
- **It is your responsibility to ensure that you keep a copy of any work submitted.**
- **Please read the Statement of Original Authorship at the bottom of the page.**
- **Sign this coversheet in the space provided below.**

Statement of Original Authorship

I, **Xianglong Lin** hereby state that the work contained in this assignment has not previously been submitted for assessment, either in whole or in part, by either myself or any other student at either The University of Queensland or at any other tertiary institution except where explicitly acknowledged. To the best of my knowledge and belief, the assignment contains no material that has been previously published or written by another person except where due reference is made. I make this Statement in full knowledge of an understanding that, should it be found to be false, I will be subject to disciplinary action under Student Integrity and Misconduct Policy 3.60.04 of the University of Queensland.

The University of Queensland 's policy on plagiarism can be found at
http://ppl.app.uq.edu.au/content/3.60.04-student-integrity-and-misconduct (Reference 3.60.04)

Signed .....Xianglong LIN..... . Date .....10.09.2020.....

Assignment 2

Q1. (a) $7260 = 2^2 \times 3 \times 5 \times 11^2$

$5720 = 2^3 \times 5 \times 11 \times 13$

$\gcd(7260, 5720) = 2^2 \times 5 \times 11$

$\text{lcm}(7260, 5720) = 2^3 \times 3 \times 5 \times 11^2 \times 13 = 188760$

(b) $434346 = 43095 \times 2 + 18156$ $\gcd(434346, 43095) = \gcd(43095, 18156)$

$43095 = 18156 \times 2 + 6783$ $\gcd(43095, 18156) = \gcd(18156, 6783)$

$18156 = 6783 \times 2 + 4590$ $\gcd(18156, 6783) = \gcd(6783, 4590)$

$6783 = 4590 \times 1 + 2193$ $\gcd(6783, 4590) = \gcd(4590, 2193)$

$4590 = 2193 \times 2 + 204$ $\gcd(4590, 2193) = \gcd(2193, 204)$

$2193 = 204 \times 10 + 153$ $\gcd(2193, 204) = \gcd(204, 153)$

$204 = 153 \times 1 + 51$ $\gcd(204, 153) = \gcd(153, 51)$

$153 = 51 \times 3 + 0$ $\gcd(153, 51) = \gcd(51, 0) = 51$.

True.

Q2. (a) Proof: $\because d \mid a$

$\therefore a = dk$ for $k \in \mathbb{Z}$.

$\because d \mid b$

$\therefore b = dx$ for $x \in \mathbb{Z}$.

$$\frac{ab}{d} = \frac{dk \cdot dx}{d}$$

$$= \frac{d^2 kx}{d}$$

$= dkx$ since $d \in \mathbb{N}$

$= ax$ bk

$\therefore \frac{ab}{d}$ is an integer that is a common multiple of both a and b. □

(b) False.

Take $a = 4, b = 2, d = 6 \in \mathbb{N}$.

(a) Then $a^2 = 4^2 = 16$, $b^2 = 2^2 = 4$

$\because 16 \bmod 6 = 4$, $4 \bmod 6 = 4 \in \mathbb{N}$

$\therefore 16 \equiv 4 \pmod 6$

$\therefore a^2 \equiv b^2 \pmod 6$

Also $\because 4 \bmod 6 = 4$, $2 \bmod 6 = 2$

$\therefore 4 \not\equiv 2 \pmod 6$.

$\therefore a \not\equiv b \pmod 6$. $\square$

(c) True.

Proof (by contradiction):

Suppose this statement is false, there is, $\exists a, b \in \mathbb{N}$, $a^3 \equiv b^3 \pmod 3 \rightarrow a \not\equiv b \pmod 3$.

Then, $a^3 - b^3 = 3k$ for some $k \in \mathbb{Z}$

So $(a-b)(a^2+b^2+ab) = 3k$

Also $\because a \not\equiv b \pmod 3$

$\therefore a - b \neq 3x$ for some $x \in \mathbb{Z}$

$\therefore a^2 + b^2 + ab = 3y$ for some $y \in \mathbb{Z}$.

There are 2 possibilities for a, b such that $a - b \neq 3x$.

① $(a-b) \equiv 1 \pmod 3$

$a \equiv 1 \pmod 3$, $b \equiv 0 \pmod 3$ or vice versa

$\therefore a^2 + b^2 + ab \equiv 1 \pmod 3$

$\therefore a^2 + b^2 + ab \neq 3y$

So we have a contradiction.

② $(a-b) \equiv 2 \pmod 3$

$\therefore a \equiv 0 \pmod 3$, $b \equiv 2 \pmod 3$ or vice versa

$\therefore a^2 + b^2 + ab \equiv 4 \equiv 1 \pmod 3$

$\therefore a^2 + b^2 + ab \neq 3y$

So we have a contradiction.

Therefore, for all $a, b \in \mathbb{N}$, if $a^3 \equiv b^3 \pmod 3$, then $a \equiv b \pmod 3$ $\square$

(d) True.

Proof (by contraposition).

Suppose that $\sqrt{x}$ is not irrational, namely rational.

$\therefore \sqrt{x} = \frac{a}{b}$ where $a, b \in \mathbb{N}$, $\gcd(a, b) = 1$

$\therefore (\sqrt{x})^2 = \frac{a^2}{b^2}$

$x = \frac{a^2}{b^2}$

$\therefore x$ is rational number

Therefore, by contraposition, we conclude that For all $x \in \mathbb{R}$, if x is irrational and $x \geq 0$, then $\sqrt{x}$ is also irrational. $\square$

(e). False.

Take $x = \frac{16}{25}$

Then $x \geq 0$ and $x \notin \mathbb{Z}$

But $\sqrt{x} = \sqrt{\frac{16}{25}} = \frac{4}{5}$ is rational. $\square$

(f) True.

Proof: $\because x \leq y$, $\lfloor x \rfloor = \lfloor y \rfloor$

$\therefore x = a + b$, $y = a + c$ where $a \in \mathbb{N}$, $b, c \in [0, 1)$, $b \leq c$

$\therefore 2x = 2(a+b) = 2a + 2b$

$2y = 2(a+c) = 2a + 2c$

There are 3 possibilities for $\lfloor x \rfloor = \lfloor y \rfloor$.

① $b \in [0, \frac{1}{2})$, $c \in [\frac{1}{2}, 1)$

Then $\lfloor 2x \rfloor = 2a$, $\lfloor 2y \rfloor = 2a$

$\therefore \lfloor 2x \rfloor = \lfloor 2y \rfloor$

② $b \in [0, \frac{1}{2})$, $c \in [\frac{1}{2}, 1)$

Then $\lfloor 2x \rfloor = 2a$, $\lfloor 2y \rfloor = 2a + 1$

$\therefore \lfloor 2x \rfloor = \lfloor 2y \rfloor - 1$

③ $b \in [\frac{1}{2}, 1)$, $c \in [\frac{1}{2}, 1)$

Then $\lfloor 2x \rfloor = 2a + 1$, $\lfloor 2y \rfloor = 2a + 1$

$\therefore \lfloor 2x \rfloor = \lfloor 2y \rfloor$

Therefore, for all $x, y \in \mathbb{R}$, if $x \leq y$ and $\lfloor x \rfloor = \lfloor y \rfloor$, the either $\lfloor 2x \rfloor = \lfloor 2y \rfloor$ or $\lfloor 2x \rfloor = \lfloor 2y \rfloor - 1$. $\square$

Q3. Proof (by contradiction): Suppose $\sqrt{10}$ is rational.

Then $\sqrt{10} = \frac{a}{b}$ where $a \in \mathbb{W}$, $b \in \mathbb{N}$, $\gcd(a, b) = 1$

$\therefore (\sqrt{10})^2 = \frac{a^2}{b^2}$

$10 = \frac{a^2}{b^2}$

$a^2 = 10b^2$

$a^2 = 2 \times 5b^2$

$\therefore a^2$ is an even number

Also $\because$ odd number times odd number is always odd

$\therefore a$ is also an even number

$\therefore a = 2x$ for some $x \in \mathbb{N}$

$\therefore (2x)^2 = 2 \cdot 5b^2$

$4x^2 = 10b^2$

$2x^2 = 5b^2$

$\because 2x^2$ is even, 5 is odd

$\therefore b^2$ must be even.

$\therefore b$ must be even

Hence, both a, b are even

$\therefore \gcd(a, b) = 2 \neq 1$

So we have a contradiction

Therefore $\sqrt{10}$ cannot be rational, that is $\sqrt{10}$ is irrational. $\square$