

Quantum walks for 1D anyons in a cavity

Ronaldo Navarro Ambriz* and Santiago Francisco Caballero Benítez†
Instituto de Física, Universidad Nacional Autónoma de México, Ciudad de México, México
(Dated: August 17, 2026)

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Keywords: anyons, quantum matter, ultracold atoms

I. INTRODUCTION

Los anyones son ...

Añadiendo el término de cavidad:

$$\mathcal{H}_{cav} = g_{cav} \sum_{i,j=1}^M (-1)^{i-j} \hat{n}_i \hat{n}_j = g_{cav} \left(\sum_{i=1}^M (-1)^i \hat{n}_i \right)^2, \quad (2)$$

Obteniendo el Hamiltoniano extendido:

$$\mathcal{H}_{ext}^b = \mathcal{H}^b + \mathcal{H}_{cav} \quad (3)$$

The original Hamiltonian (1) breaks the inversion symmetry $\mathcal{I}$, as the particles in the chain acquire a phase due a site occupation. Result, only for a change of sign in both term we will observe a total reflection in the quantum walk

II. MAIN IDEA

Del Hamiltoniano de bosones condicionados:

$$\mathcal{H}^b = -J \sum_{i=1}^M \left(\hat{b}_i^\dagger \hat{b}_{i+1} e^{i\theta \hat{n}_i} + h.c. \right) + \frac{U}{2} \sum_{i=1}^M \hat{n}_i (\hat{n}_i - 1), \quad (1)$$

$$\langle \hat{n}_j(t) \rangle_{+U,+g} = \langle \hat{n}'_j(t) \rangle_{-U,-g} \quad (4)$$

$$\langle \hat{n}_j(t) \rangle_{+U,-g} = \langle \hat{n}'_j(t) \rangle_{-U,+g} \quad (5)$$

III. CONCLUSION

Appendix: Appendix

Del operador de paridad:

$$\mathcal{P} = e^{i\pi \sum_r \hat{n}_{2r+1}} \quad (A.1)$$

* Correspondence email address: ron@estudiantes.unam.fisi

† Correspondence email address: scaballero@fisica.unam.mx