

Wave guides as an analogous system of 1D optical lattice for anyons

Ronaldo Navarro Ambriz* and Santiago Francisco Caballero Benítez†
Instituto de Física, Universidad Nacional Autónoma de México, Ciudad de México, México
 (Dated: September 30, 2026)


Lorem ipsum dolor sit amet, consectetur adipiscing elit. Ut purus elit, vestibulum ut, placerat ac, adipiscing vitae, felis. Curabitur dictum gravida mauris. Nam arcu libero, nonummy eget, consectetur id, vulputate a, magna. Donec vehicula augue eu neque. Pellentesque habitant morbi tristique senectus et netus et malesuada fames ac turpis egestas. Mauris ut leo. Cras viverra metus rhoncus sem. Nulla et lectus vestibulum urna fringilla ultrices. Phasellus eu tellus sit amet tortor gravida placerat. Integer sapien est, iaculis in, pretium quis, viverra ac, nunc. Praesent eget sem vel leo ultrices bibendum. Aenean faucibus. Morbi dolor nulla, malesuada eu, pulvinar at, mollis ac, nulla. Curabitur auctor semper nulla. Donec varius orci eget risus. Duis nibh mi, congue eu, accumsan eleifend, sagittis quis, diam. Duis eget orci sit amet orci dignissim rutrum.

Keywords: anyons, quantum matter, ultracold atoms

I. INTRODUCTION

Los anyones son ...

II. MAIN IDEA



Para las guías de onda:

- Una guía de onda posee un índice de refracción en el otro núcleo, rodeado por otro de menor índice $n_1 < n_2$
- Para una única guía de onda, los modos admitidos están dados por sus características físicas.
- Un arreglo de guías de onda, el acoplamiento permitirá la transferencia entre los elementos del arreglo.

From an infinite waveguides array with homogeneous optical single mode waveguides and equally spaced along z-axis. This is described by the following differential equation [1]

$$i \frac{dE_j(z)}{dz} = g_1(E_{j-1} + E_{j+1}), \quad j = -\infty, \dots, \infty, \quad (1)$$

From this

$$i \frac{d|\psi(z)\rangle}{dz} = \hat{H}|\psi(z)\rangle, \quad \hat{H} = g_1 (\hat{V}_\infty + \hat{V}_\infty^\dagger), \quad (2)$$

where the operators $\hat{V}_\infty$ and $\hat{V}_\infty^\dagger$ are the infinite counterparts of the London-Susskind-Glogower operators [2, 3], defined as

$$\hat{V}_\infty = \sum_{n=-\infty}^{\infty} |n\rangle\langle n+1|, \quad \hat{V}_\infty^\dagger = \sum_{n=-\infty}^{\infty} |n+1\rangle\langle n|, \quad (3)$$

For an analytical solution we need to satisfy in our Hilbert space the following conditions for the LSG operator

$$[\hat{V}_\infty, \hat{V}_\infty^\dagger] = 0, \quad \hat{V}_\infty \hat{V}_\infty^\dagger = \hat{I},$$

In this framework the initial state is a photon in the n site of an infinite array of waveguides $|\psi(0)\rangle = |n_0\rangle$;

$$|\psi_{n_0}(z)\rangle = \exp \left[-g_1 z \left(i\hat{V}_\infty^\dagger - \frac{1}{i\hat{V}_\infty} \right) \right] |n_0\rangle.$$

Donde podemos identificar la función generadora de las funciones Bessel, para obtener

$$|\psi_{n_0}(z)\rangle = \sum_{n=-\infty}^{\infty} J_n(-2g_1 z) i^n \hat{V}_\infty^{\dagger n} |n_0\rangle.$$

para este modelo podemos hacer una analogía con un modelo de bosones en una red óptica partiendo del operador $\hat{V}_\infty$ y la evolución será escrita en función del tiempo:

$$\hat{V}_\infty = \sum_{n=-\infty}^{\infty} \hat{b}_n^\dagger \hat{b}_{n+1}, \quad \hat{V}_\infty^\dagger = \sum_{n=-\infty}^{\infty} \hat{b}_{n+1}^\dagger \hat{b}_n$$

The operator $\hat{V}_\infty^{\dagger n}$ is equivalent of create a particle in n site in the framework of optical lattices for a subspace of one particle per site:

$$\hat{V}_\infty^{\dagger n} |1_0\rangle = \hat{b}_n^\dagger |0\rangle, \quad (4)$$

* Correspondence email address: ron@estudiantes.unam.fisi

† Correspondence email address: scaballero@fisica.unam.mx

Obtaining the dynamics for a particle in 0-site

$$|\psi_0(t)\rangle = \sum_{n=-\infty}^{\infty} J_n(-2Jt) i^n \hat{b}_n^\dagger |0\rangle. \quad (5)$$

We could obtain a rich dynamics for n-particles system working with a Heisenberg representation for one body operators $\hat{b}^\dagger$ applied in the sites, obtaining in

general the expression:

$$\langle \hat{n}_j \rangle = \sum_{m=-\infty}^{\infty} |J_{m-j}(-2Jt)|^2 \langle \hat{n}_m(0) \rangle \quad (6)$$

$$S(q) = \sum_{l=0}^M |J_l(-2Jt)|^2 - \sum_{l,m=0}^M e^{iq(l-m)} |J_l(-2Jt)|^2 |J_m(-2Jt)|^2$$

III. CONCLUSION

-
- [1] Alan L. Jones. Coupling of optical fibers and scattering in fibers. *J. Opt. Soc. Am.*, 55:261–271, Mar 1965.
 - [2] Fritz London. Über die jacobischen transformationen der quantenmechanik. *Z. Phys.*, 37:915–925, Dec 1926.
 - [3] Leonard Susskind and Jonathan Glogower. Quantum mechanical phase and time operator. *Physics Physique Fizika*, 1:49–61, Jul 1964.

so the expected value of $\hat{n}_j$ is:

$$\begin{aligned} \langle \hat{n}_j(t) \rangle &= \langle \psi(0) | \hat{b}_j^\dagger(t) \hat{b}_j(t) | \psi(0) \rangle, \\ &= \langle \psi(0) | \sum_{m,m'=-\infty}^{\infty} i^{m-m'} J_{m-j}(-2Jt) J_{m'-j}(-2Jt) \\ &\quad \hat{b}_m^\dagger \hat{b}_m | \psi(0) \rangle, \end{aligned} \quad (A.2)$$

In general we get a 1-d chain, if we annihilate at m' site and create at m we will obtain an orthogonal state, so for the different values of $m \neq m'$ we will obtain zero, so for an arbitrary initial state we get:

$$\langle \hat{n}(t) \rangle = \sum_{m=-\infty}^{\infty} |J_{m-j}(-2Jt)|^2 \langle \hat{n}_m(0) \rangle \quad (A.3)$$

2. Passive representation of $\hat{n}_j$ operator

For this framework we need to specify the initial state, first, for 2 particles in the sites j, j' respectively, with $j \neq j'$, the state as function of time is

$$|\psi(t)\rangle = e^{-i\mathcal{H}t} |\psi(0)\rangle = e^{-i\mathcal{H}t} \hat{b}_j^\dagger \hat{b}_{j'}^\dagger |0\rangle, \quad (A.4)$$

as the temporal propagator is unitary we use the identity to get the following expressions

$$\begin{aligned} |\psi(t)\rangle &= e^{-i\mathcal{H}t} \hat{b}_j^\dagger \mathcal{I} \hat{b}_{j'}^\dagger \mathcal{I} |0\rangle \\ &= e^{-i\mathcal{H}t} \hat{b}_j^\dagger e^{i\mathcal{H}t} e^{-i\mathcal{H}t} \hat{b}_{j'}^\dagger e^{i\mathcal{H}t} e^{-i\mathcal{H}t} |0\rangle \\ &= e^{-i\mathcal{H}t} \hat{b}_j^\dagger e^{i\mathcal{H}t} e^{-i\mathcal{H}t} \hat{b}_{j'}^\dagger e^{i\mathcal{H}t} |0\rangle, \end{aligned} \quad (A.5)$$

Appendix: Appendix

In this paper we suggest to calculate the operator from active representations due the feasibility of operations

1. Active representation of $\hat{n}_j$ operator

We obtain for a $|\psi\rangle$ as steady state and $\hat{b}^\dagger$ as dynamic operator, write as:

$$\hat{b}_j^\dagger(t) = \sum_{m=-\infty}^{\infty} i^{m-j} J_{m-j}(-2Jt) \hat{b}_m^\dagger(0), \quad (A.1)$$