

Quantum Mechanics

RongYu

September 23, 2025

This note is compiled based on the content of the Quantum Mechanics course taught by Professor Hong Guo from School of Electronics, Peking University.

Contents

1	Review of Preparatory Knowledge	5
1.1	Classical Mechanics	5
1.2	Newtonian Mechanics	5
1.3	Lagrangian Mechanics - Analytical M.	5
1.3.1	Constraint	5
1.3.2	Coordinates and Space	5
1.3.3	Functional	6
1.3.4	Difference & Differential & Variation	6
1.3.5	Principle of Least Action (Hamilton's Principle)	6
1.3.6	Euler-Lagrange Equations	6
1.4	Hamiltonian Mechanics - Analytical M.	7
1.4.1	Legendre Transformations	7
1.4.2	Hamilton's Canonical Equations of Motion	7
1.4.3	Poisson Bracket (Classical Canonical Commutator)	8
1.4.4	Fundamental Commutation Relations in CM	8
1.4.5	Properties of Poisson Bracket	8
1.4.6	Hamilton-Jacobi Equation	9
1.4.7	Phase Space (Dynamics)	9
1.4.8	Regular Lagrangian	9
2	Classical Electrodynamics	10
2.1	Vector Analysis	10
2.1.1	Generalized Stokes' Theorem	10
2.1.2	Helmholtz's theorem	10
2.2	Experimental Laws and Maxwell's Equations	11
2.2.1	Coulomb's Law	11
2.2.2	Biot-Savart's Law	11
3	Fundamental Postulates (Axioms)	11

4	SOME IS	11
4.1	State	11
4.2	Observable	12
4.3	Measurement	12
4.3.1	Measurement + Outcomes	12
4.3.2	Probability of Measurement Outcomes	12
4.3.3	State Collapse	12
4.4	Evolution (Dynamics)	12
4.4.1	Schrödinger Picture	12
4.4.2	Heisenberg Picture	13
4.5	Identical Particles and Symmetrization	13
4.5.1	Identical Particles	13
4.5.2	Symmetrization	13
5	State Vector	13
5.1	Euclidean Space \mathbb{R}^3	13
5.2	State Vector Space \mathbb{H} (Hilbert Space)	14
6	Operator	14
6.1	Linear Operator	15
6.1.1	Two Kinds of Linear Operators	15
6.1.2	Rules	15
6.1.3	Properties	15
7	Overview of QM	15
8	Roles: State Vector, Operator	16
8.0.1	Operator	16
8.0.2	State	16
8.0.3	Eigenspectrum	16
8.1	Properties	16
8.2	Problems	16
8.2.1	Eigenvalue Problem	16
8.2.2	Evolution Problem	16
8.3	Picture	16
8.3.1	Schrödinger Picture	16
8.3.2	Heisenberg Picture	17
8.4	Particles	17
8.5	Perturbation Theories	17
8.6	Angular Momentum and Magnetic Moment	17
8.7	Mechanics	17
8.8	Theorem of Hermitian Operator	17
9	Wave Mechanics & Matrix Mechanics	18
9.1	Wave Mechanics: wavefunction	18
9.2	Matrix Mechanics: State Vector	19
9.3	Matrix Mechanics: Hermitian Conjugate	19

10 Commutation	20
10.0.1 Definition	20
10.0.2 Compare	20
10.0.3 Relation	20
11 Properties of Commutations	20
12 Fundamental Commutation Relations	21
13 Commutation Relations of Orbital Angular Momentum	22
14 Eigenvalue Problem	23
15 Observables	23
16 Eigenvalue & Eigenstates	24
17 Spin and Pauli Matrices	25
17.1 Stern-Gerlach Experiment	25
17.1.1 Meanings	25
17.1.2 Analysis	25
17.1.3 History	25
17.1.4 Cascaded S-G Experiment	25
18 Spin Angular Momentum	26
18.0.1 Significance	26
19 Algebraic Method to Eigenvalue Problem of Angular Momentum	27
20 Review	27
20.1 Orbital Angular Momentum	27
20.1.1 Definition of Orbital AM	27
20.1.2 Commutation Relations of Orbital AM	27
20.1.3 Eigenvalue Equation	27
20.2 Intrinsic Angular Momentum (Spin)	27
20.2.1 Eigenvalue Equation	27
20.3 Pauli Matrix	28
20.3.1 Ladder / Raising & Lowering / Transition Operator	28
20.3.2 Pauli Matrices ($S_z(\sigma_z)$ -representation)	28
21 Arbitrary Angular Momentum	28
21.1 Ladder Operator	29
21.1.1 Definition of Ladder Operators	29
21.1.2 Effect of Ladder Operators on Common Eigenstates	29
21.1.3 Restriction	29
21.1.4 Commutations	29
21.2 Eigenvalue Equation of Angular Momentum Operator	30

22	Commutation Relations between Angular Momentum Operators and Hamiltonian	31
23	Addition of Angular Momenta	31
24	Uncertainty Principle	32
24.1	Heisenberg Uncertainty Principle	32
24.2	Generalized Uncertainty Principle	32
24.2.1	Essential Mathematical Tools	32
24.2.2	Define Deviation Operator	33
24.2.3	Properties of $\hat{\sigma}_A, \hat{\sigma}_B$	33
24.2.4	Proof of GUP	33
24.3	Energy-Time Uncertainty Relation	33
24.3.1	Ehrenfest Theorem	33
24.3.2	Proof	34
24.4	Minimum Uncertainty State	34
24.4.1	MUS of \hat{x} & \hat{p}	34
25	Quantum Dynamics	35
25.1	Schrödinger Equation	35
25.1.1	Remark	35
25.1.2	Representation of Position	35
25.2	Time Evolution Operator	35
25.2.1	TEO for Conservative System	36
25.2.2	Properties of $\hat{U}(t, t_0)$	36
26	title	37

1 Review of Preparatory Knowledge

1.1 Classical Mechanics

1. $\delta S \equiv 0$ Principle of Least Action
2. $\frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha} = \frac{\partial \mathcal{L}}{\partial q_\alpha} \quad (\alpha = 1, 2, \dots, D)$ Euler-Lagrange eqs.
3. $\begin{cases} \dot{q}_\alpha = \frac{\partial H}{\partial p_\alpha} \\ \dot{p}_\alpha = -\frac{\partial H}{\partial q_\alpha} \end{cases} \quad (\alpha = 1, 2, \dots, D)$ Hamilton's Canonical eqs.
4. $\frac{df}{dt} = \frac{\partial f}{\partial t} + [f, H]_{\text{PB}}$ Poisson Bracket
5. $\frac{\partial S}{\partial t} + H(q, \frac{\partial S}{\partial q}, t) = 0$ Hamilton-Jacobi eq.

1.2 Newtonian Mechanics

Euclidean Space (Physical Space) \longrightarrow Cartesian (Descartes) Coordinates

$$\frac{d\mathbf{p}}{dt} = \mathbf{F}, \quad \mathbf{p} = m\mathbf{v} \quad (\text{Momentum \& Force})$$

$$\frac{d\mathbf{L}}{dt} = \mathbf{M}, \quad \mathbf{L} = \mathbf{x} \times \mathbf{p}, \quad \mathbf{M} = \mathbf{x} \times \mathbf{F} \quad (\text{Angular Momentum \& Torque})$$

Mass point \longrightarrow Mass point system \longrightarrow Rigid Body \longrightarrow Inertia $\begin{cases} \text{Mass: } m \\ \text{Inertia of Rotation: } mr^2 \end{cases}$

1.3 Lagrangian Mechanics - Analytical M.

1.3.1 Constraint

x_1, x_2, \dots, x_{3N} are dependent $\implies q_1, q_2, \dots, q_D$ are independent $\left(\frac{\partial q_i}{\partial q_j} = 0, \quad i \neq j \right)$

constraint $\begin{cases} \text{holonomic: } f(x; t) \equiv 0 \\ \text{non-holonomic: } f(x, \dot{x}; t) \equiv 0 \end{cases}$

1.3.2 Coordinates and Space

Generalized Coordinates $q_\alpha(t)$, $\alpha = 1, 2, \dots, D \leq 3N$

Configuration Space (Abstract Space): D -dim

1.3.3 Functional

Example of functionals (function of function):

1. Lagrangian:

$$\mathcal{L}(q(t), \dot{q}(t); t)$$

generalized velocity: $\dot{q} = \frac{dq}{dt}$

2. Hamiltonian:

$$H(q(t), p(t); t)$$

canonical position: $q(t)$; canonical momentum: $p_\alpha = \frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha}$ (in Phase Space)

3. Action:

$$S[q(t)] = \int_{t_1}^{t_2} \mathcal{L}(q(t), \dot{q}(t); t) dt$$

Dimensions of these functionals:

$$[\mathcal{L}] = [H] = [\text{Energy}], [S] = [\text{Energy}] \cdot [\text{Time}] = [\text{Length}] \cdot [\text{Momentum}] = [\hbar]$$

Reduced Planck's Constant: $\hbar = \frac{h}{2\pi}$

1.3.4 Difference & Differential & Variation

$$f(x_2) - f(x_1) = \underbrace{\Delta f}_{\text{difference}} \xrightarrow[\Delta x = x_2 - x_1]{\Delta x \rightarrow 0} \underbrace{df}_{\text{differential}}, \text{ variation: } \delta q(t) = q_2(t) - q_1(t)$$

$$f(q, \dot{q}; t) : \begin{cases} \Delta f = \frac{\partial f}{\partial q} \Delta q + \frac{\partial f}{\partial \dot{q}} \Delta \dot{q} \\ df = \frac{\partial f}{\partial q} dq + \frac{\partial f}{\partial \dot{q}} d\dot{q} \\ \delta f = \frac{\partial f}{\partial q} \delta q + \frac{\partial f}{\partial \dot{q}} \delta \dot{q} \quad (\delta q(t_1) = \delta q(t_2) = 0; \delta t = 0) \end{cases}$$

1.3.5 Principle of Least Action (Hamilton's Principle)

$$\boxed{\delta S \equiv 0} \implies \text{dynamical equations}$$

1.3.6 Euler-Lagrange Equations

$$\boxed{\frac{\partial \mathcal{L}}{\partial q_\alpha} = \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha} \right), \quad \alpha = 1, 2, \dots, D}$$

D 2nd-order ODE

$$\begin{aligned}
0 \equiv \delta S &= \delta \int_{t_1}^{t_2} \mathcal{L}(q, \dot{q}; t) dt = \int_{t_1}^{t_2} \left(\frac{\partial \mathcal{L}}{\partial q} \delta q + \frac{\partial \mathcal{L}}{\partial \dot{q}} \delta \dot{q} \right) dt \\
&= \int_{t_1}^{t_2} \left[\frac{\partial \mathcal{L}}{\partial q} \delta q + \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q \right) - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \delta q \right] dt \\
&= \int_{t_1}^{t_2} \left[\frac{\partial \mathcal{L}}{\partial q} - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \right] \delta q dt + \underbrace{\frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q \Big|_{t_1}^{t_2}}_{=0} \implies \frac{\partial \mathcal{L}}{\partial q} - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \equiv 0
\end{aligned}$$

1.4 Hamiltonian Mechanics - Analytical M.

1.4.1 Legendre Transformations

$$\begin{cases} p_\alpha = \frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha}, \alpha = 1, 2, \dots, D \\ H(q, p; t) = \sum_\alpha p_\alpha \dot{q}_\alpha - \mathcal{L}(q, \dot{q}; t) \end{cases}$$

$$\implies \frac{\partial H}{\partial \dot{q}_\alpha} = p_\alpha - \frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha} \equiv 0 \implies H \text{ is } \dot{q} \text{ - independent, } \mathcal{L} \text{ is } p \text{ - independent}$$

1.4.2 Hamilton's Canonical Equations of Motion

$$\boxed{\begin{cases} \dot{q}_\alpha = \frac{\partial H}{\partial p_\alpha} \\ \dot{p}_\alpha = -\frac{\partial H}{\partial q_\alpha} \end{cases} \quad \alpha = 1, 2, \dots, D}$$

2D 1st-order ODE

$$\begin{aligned}
0 \equiv \frac{\partial \mathcal{L}}{\partial p_\alpha} &= \dot{q}_\alpha - \frac{\partial H}{\partial p_\alpha} \implies \dot{q}_\alpha = \frac{\partial H}{\partial p_\alpha}; \dot{p}_\alpha = \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}_\alpha} \right) = \frac{\partial \mathcal{L}}{\partial q_\alpha}, \frac{\partial H}{\partial q_\alpha} = -\frac{\partial \mathcal{L}}{\partial q_\alpha} \implies \dot{p}_\alpha = -\frac{\partial H}{\partial q_\alpha} \\
dH &= \frac{\partial H}{\partial t} dt + \frac{\partial H}{\partial q} dq + \frac{\partial H}{\partial p} dp = d(p\dot{q} - \mathcal{L}(q, \dot{q}; t)) \\
&= (dp)\dot{q} + p d\dot{q} - \frac{\partial \mathcal{L}}{\partial t} dt - \frac{\partial \mathcal{L}}{\partial q} dq - \frac{\partial \mathcal{L}}{\partial \dot{q}} d\dot{q} = \dot{q} dp - \dot{p} dq - \frac{\partial \mathcal{L}}{\partial t} dt \\
&\implies \dot{q} = \frac{\partial H}{\partial p}, \dot{p} = -\frac{\partial H}{\partial q} \\
0 \equiv \delta S &= \delta \int (p\dot{q} - H(q, p; t)) dt \implies \dot{q} = \frac{\partial H}{\partial p}, \dot{p} = -\frac{\partial H}{\partial q}
\end{aligned}$$

1.4.3 Poisson Bracket (Classical Canonical Commutator)

$$[A, B]_{\text{PB}} = \sum_{\alpha} \left(\frac{\partial A}{\partial q_{\alpha}} \frac{\partial B}{\partial p_{\alpha}} - \frac{\partial A}{\partial p_{\alpha}} \frac{\partial B}{\partial q_{\alpha}} \right)$$

$$\Rightarrow \text{Quantum Canonical Commutator } [\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A}$$

$$\forall f(q, p; t), \quad \frac{df}{dt} = \frac{\partial f}{\partial t} + \sum_{\alpha} \left(\frac{\partial f}{\partial q_{\alpha}} \dot{q}_{\alpha} + \frac{\partial f}{\partial p_{\alpha}} \dot{p}_{\alpha} \right) = \frac{\partial f}{\partial t} + \sum_{\alpha} \left(\frac{\partial f}{\partial q_{\alpha}} \frac{\partial H}{\partial p_{\alpha}} - \frac{\partial f}{\partial p_{\alpha}} \frac{\partial H}{\partial q_{\alpha}} \right) = \frac{\partial f}{\partial t} + [f, H]_{\text{PB}}$$

$$\text{if } f(q, p; t) = q_{\alpha} \Rightarrow \frac{dq_{\alpha}}{dt} = \frac{\partial H}{\partial p_{\alpha}}, \text{ if } f(q, p; t) = p_{\alpha} \Rightarrow \frac{dp_{\alpha}}{dt} = -\frac{\partial H}{\partial q_{\alpha}}$$

1.4.4 Fundamental Commutation Relations in CM

$$[q_{\alpha}, p_{\beta}]_{\text{PB}} = \delta_{\alpha\beta} \longleftrightarrow [\hat{q}_{\alpha}, \hat{p}_{\beta}] = i\hbar\delta_{\alpha\beta}$$

$$[q_{\alpha}, q_{\beta}]_{\text{PB}} = [p_{\alpha}, p_{\beta}]_{\text{PB}} \equiv 0 \longleftrightarrow [\hat{q}_{\alpha}, \hat{q}_{\beta}] = [\hat{p}_{\alpha}, \hat{p}_{\beta}] \equiv 0$$

$$[q_{\alpha}, p_{\beta}] = \sum_{\gamma} \left(\frac{\partial q_{\alpha}}{\partial q_{\gamma}} \frac{\partial p_{\beta}}{\partial p_{\gamma}} - \frac{\partial q_{\alpha}}{\partial p_{\gamma}} \frac{\partial p_{\beta}}{\partial q_{\gamma}} \right) = \delta_{\alpha\gamma} \delta_{\beta\gamma} - 0 = \delta_{\alpha\beta}$$

$$[q_{\alpha}, q_{\beta}] = \sum_{\gamma} \left(\frac{\partial q_{\alpha}}{\partial q_{\gamma}} \frac{\partial q_{\beta}}{\partial p_{\gamma}} - \frac{\partial q_{\alpha}}{\partial p_{\gamma}} \frac{\partial q_{\beta}}{\partial q_{\gamma}} \right) = \delta_{\alpha\gamma} \cdot 0 - 0 \cdot \delta_{\beta\gamma} = 0$$

1.4.5 Properties of Poisson Bracket

1. $[A, B]_{\text{PB}} = -[B, A]_{\text{PB}}$
2. $[A + B, C]_{\text{PB}} = [A, C]_{\text{PB}} + [B, C]_{\text{PB}}, [A, B + C]_{\text{PB}} = [A, B]_{\text{PB}} + [A, C]_{\text{PB}}$
3. $[A, BC]_{\text{PB}} = [A, B]_{\text{PB}}C + B[A, C]_{\text{PB}}, [AB, C]_{\text{PB}} = [A, C]_{\text{PB}}B + A[B, C]_{\text{PB}}$
4. $[A, B^n]_{\text{PB}} = n[A, B]_{\text{PB}}B^{n-1}$
5. $[A, f(B)]_{\text{PB}} = [A, B]_{\text{PB}} \frac{\partial f}{\partial B}$
6. $[L_i, L_j]_{\text{PB}} = \epsilon_{ijk} L_k$
7. $[L_i, \mathbf{L}^2]_{\text{PB}} = 0$
8. $[L_i, x_j]_{\text{PB}} = \epsilon_{ijk} x_k$
9. $[L_i, p_j]_{\text{PB}} = \epsilon_{ijk} p_k$

$$\begin{aligned}
[L_i, L_j]_{\text{PB}} &= [\epsilon_{iab}x_ap_b, \epsilon_{jcd}x_cp_d]_{\text{PB}} = \epsilon_{iab}\epsilon_{jcd} \left(x_a \underbrace{[p_b, x_c]_{\text{PB}}}_{=-\delta_{bc}} p_d + x_c \underbrace{[x_a, p_d]_{\text{PB}}}_{=\delta_{ad}} p_b \right) \\
&= \epsilon_{iab}\epsilon_{jca}x_cp_b - \epsilon_{iab}\epsilon_{jbd}x_ap_d = (\delta_{bj}\delta_{ic} - \delta_{bc}\delta_{ij})x_cp_b - (\delta_{ij}\delta_{ad} - \delta_{id}\delta_{aj})x_ap_d \\
&= x_ip_j - \delta_{ij}(\mathbf{x} \cdot \mathbf{p}) + \delta_{ij}(\mathbf{x} \cdot \mathbf{p}) - x_jp_i = x_ip_j - x_jp_i = \epsilon_{ijk}L_k
\end{aligned}$$

1.4.6 Hamilton-Jacobi Equation

$$\boxed{\frac{\partial S}{\partial t} + H(q, \frac{\partial H}{\partial q}; t) = 0}$$

$$\begin{aligned}
\delta S &= \int_{t_0}^t \mathcal{L}(q, \dot{q}; t) dt = \int_{t_0}^t \left[\frac{\partial \mathcal{L}}{\partial q} \delta q + \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q \right) - \frac{d}{dt} \left(\frac{\partial \mathcal{L}}{\partial \dot{q}} \right) \delta q \right] \\
&= \int_{t_0}^t \underbrace{\left(\frac{\partial \mathcal{L}}{\partial q} - \frac{d}{dt} \frac{\partial \mathcal{L}}{\partial \dot{q}} \right)}_{=0 \text{ (physical path)}} \delta q dt + \underbrace{\frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q(t) - \frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q(t_0)}_{=0 \text{ (fixed)}} = \frac{\partial \mathcal{L}}{\partial \dot{q}} \delta q = p \delta q
\end{aligned}$$

$$\begin{cases} \delta S = p \delta q \\ \delta S[q(t), t] = \frac{\partial S}{\partial q} \delta q \end{cases} \implies p = \frac{\partial S}{\partial q}, \begin{cases} \frac{dS}{dt} = \frac{\partial S}{\partial t} + \frac{\partial S}{\partial q} \dot{q} \\ \frac{dS}{dt} = \mathcal{L} = p\dot{q} - H \end{cases} \implies \frac{\partial S}{\partial t} + H = 0$$

1.4.7 Phase Space (Dynamics)

phase point $(q, p) \longrightarrow$ state

$$\int \rho(q, p; t) d^D q d^D p = 1$$

ρ : probability

$$\boxed{\frac{\partial \rho}{\partial t} = [H, \rho]_{\text{PB}}} \text{ Liouville Theorem}$$

$$0 = \frac{d\rho}{dt} = \frac{\partial \rho}{\partial t} + [\rho, H]_{\text{PB}} \xrightarrow{QM} \frac{\partial \hat{\rho}}{\partial t} = \frac{1}{i\hbar} [\hat{H}, \hat{\rho}]$$

1.4.8 Regular Lagrangian

$$\left| \frac{\partial^2 \mathcal{L}}{\partial q_i \partial q_j} \right| \neq 0 \implies \begin{cases} \mathcal{L} = T - V \\ H = T + V \end{cases}$$

2 Classical Electrodynamics

2.1 Vector Analysis

$$\mathbf{v} = v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + v_3 \mathbf{e}_3 = \sum_i v_i \mathbf{e}_i = v_i \mathbf{e}_i \text{ (Einstein convention), } i = 1, 2, 3$$

2.1.1 Generalized Stokes' Theorem

$$\boxed{\int_{\Omega} d\omega = \int_{\partial\Omega} \omega}$$

1. Newton-Leibniz formula

$$\int_a^b df = f(b) - f(a)$$

2. Green's theorem

$$\oint f dx + g dy = \iint \left(\frac{\partial g}{\partial x} - \frac{\partial f}{\partial y} \right) dx dy$$

3. Stokes' theorem

$$\begin{aligned} \oint \mathbf{F} \cdot d\mathbf{l} &= \iint (\nabla \times \mathbf{F}) \cdot d\mathbf{S} \\ \Rightarrow \oint \psi d\mathbf{l} &= \iint d\mathbf{S} \times \nabla \psi \end{aligned}$$

4. Gauss-Ostrogradsky's theorem

$$\begin{aligned} \oiint \mathbf{F} \cdot d\mathbf{S} &= \iiint (\nabla \cdot \mathbf{F}) dV \\ \Rightarrow \begin{cases} \oiint \psi d\mathbf{S} = \iiint (\nabla \psi) dV \\ \oiint d\mathbf{S} \nabla \mathbf{A} = \iiint (\nabla \times \mathbf{A}) dV \end{cases} \end{aligned}$$

2.1.2 Helmholtz's theorem

For all continuous differentiable \mathbf{F} ,

$$\boxed{\mathbf{F} = \mathbf{F}_{\perp} + \mathbf{F}_{\parallel}}$$

1. transverse component

$$\nabla \cdot \mathbf{F}_{\perp} = 0 \quad \& \quad \mathbf{F}_{\perp} = \nabla \times \mathbf{A}$$

2. longitudinal component

$$\nabla \times \mathbf{F}_{\parallel} = 0 \quad \& \quad \mathbf{F}_{\parallel} = -\nabla \phi$$

For magnetic field \mathbf{B}

$$\nabla \cdot \mathbf{B} \equiv 0$$

For electric field \mathbf{E}

$$\begin{cases} \nabla \times \mathbf{E} = 0 & \text{for static electric field} \\ \nabla \times \mathbf{E} \neq 0 & \text{for AC electric field} \end{cases}$$

2.2 Experimental Laws and Maxwell's Equations

2.2.1 Coulomb's Law

$$\begin{aligned} \mathbf{E}(\mathbf{x}) &= \int \frac{\rho(\mathbf{x}')(\mathbf{x} - \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^3} dV' = -\nabla\phi, \quad \phi = \int \frac{\rho(\mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|} dV' \\ \implies \boxed{\nabla \cdot \mathbf{E} = 4\pi\rho(\mathbf{x})} & \quad \text{Gauss' Law} \end{aligned}$$

2.2.2 Biot-Savart's Law

$$\mathbf{B} = \int \frac{\mathbf{j}(\mathbf{x}') \times (\mathbf{x} - \mathbf{x}')}{|\mathbf{x} - \mathbf{x}'|^3} dV' = \nabla \times \mathbf{A}(\mathbf{x})$$

3 Fundamental Postulates (Axioms)

4 SOME IS

1. State
2. Observable (Operation)
3. Measurement
4. Evolution
5. Identical Particals and Symmetrization

4.1 State

system \longleftrightarrow state vector

ket: $|\psi\rangle$

bra: $\langle\psi|$

$$(|\psi\rangle)^\dagger = \langle\psi|$$

$$[(|\psi\rangle)^\dagger]^\dagger = (\langle\psi|)^\dagger = |\psi\rangle$$

\dagger : adjoint or Hermitian conjugate

4.2 Observable

physical quantity \longleftrightarrow observable \longleftrightarrow operation $\hat{A} \begin{cases} \text{linear} \\ \text{Hermitian} \end{cases}$

4.3 Measurement

4.3.1 Measurement + Outcomes

$$\hat{A} |\psi\rangle = A_n |\psi_n\rangle \text{ (Eigenvalue equation)}$$

\hat{A} : operation (q-number); A_n : eigenvalue (c-number); ψ_n : eigenstate

4.3.2 Probability of Measurement Outcomes

$$|\psi\rangle = \sum_n c_n |\psi_n\rangle, \quad c_n = \langle \psi_n | \psi \rangle$$

$$P(A_n) = |c_n|^2 = |\langle \psi_n | \psi \rangle|^2$$

$$\langle \hat{A} \rangle = \langle \psi | \hat{A} | \psi \rangle = \sum_n P(A_n) A_n = \sum_n |\langle \psi_n | \psi \rangle|^2 A_n$$

4.3.3 State Collapse

$$|\psi\rangle \xrightarrow{\hat{A}\text{-measurement}} |\psi_n\rangle$$

4.4 Evolution (Dynamics)

4.4.1 Schrödinger Picture

state: change; observable: no change

$$\begin{cases} \boxed{i\hbar \frac{d}{dt} |\psi(t)\rangle = \hat{H} |\psi(t)\rangle} & \text{Schrödinger Equation} \\ |\psi(t=0)\rangle = |\psi_0\rangle \end{cases}$$

where $\hat{H} = \hat{T} + \hat{V} = \frac{\hat{\mathbf{p}}^2}{2m} + \hat{V}(\hat{\mathbf{x}})$

representation: coordinate system

$\hat{\mathbf{x}}$ -representation (position):

$$\hat{\mathbf{x}} |\mathbf{x}\rangle = \mathbf{x} |\mathbf{x}\rangle, \quad |\psi\rangle \longrightarrow \langle \mathbf{x} | \psi \rangle = \psi(\mathbf{x}) \rightarrow \text{wave-function}$$

$$\implies \boxed{i\hbar \frac{\partial}{\partial t} \psi(\mathbf{x}, t) = -\frac{\hbar^2}{2m} \nabla^2 \psi(\mathbf{x}, t) + \hat{V}(\mathbf{x}, t) \psi(\mathbf{x}, t)}$$

$\hat{\mathbf{p}}$ -representation (momentum): $\hat{\mathbf{p}} |\mathbf{p}\rangle = \mathbf{p} |\mathbf{p}\rangle$

4.4.2 Heisenberg Picture

state: no change; observable: change

$$\boxed{\frac{d\hat{F}}{dt} = \frac{\partial \hat{F}}{\partial t} + \frac{1}{i\hbar} [\hat{F}, \hat{H}]} \quad \text{Heisenberg Equation}$$

4.5 Identical Particles and Symmetrization

4.5.1 Identical Particles

$$\begin{cases} \text{boson: } s = 0, 1, 2, \dots & \text{Bose-Einstein Statistics} \\ \text{fermion: } s = \frac{1}{2}, \frac{3}{2}, \dots & \text{Fermi-Dirac Statistics} \end{cases} \xrightarrow[\text{reduce}]{\text{classical}} \text{Maxwell-Boltzmann Statistics}$$

4.5.2 Symmetrization

$$\begin{cases} \text{Boson: symmetric} \\ \text{Femion: anti-symmetric} \end{cases}$$

5 State Vector

5.1 Euclidean Space \mathbb{R}^3

$$\begin{aligned} \mathbf{v} &= v_1 \mathbf{e}_1 + v_2 \mathbf{e}_2 + v_3 \mathbf{e}_3 \\ \mathbf{e}_1 &= \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \quad \mathbf{e}_2 = \begin{pmatrix} 0 \\ 1 \\ 0 \end{pmatrix}, \quad \mathbf{e}_3 = \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \\ \mathbf{v} &= \begin{pmatrix} v_1 \\ v_2 \\ v_3 \end{pmatrix}, \quad \mathbf{v}^T = (v_1 \quad v_2 \quad v_3) \end{aligned}$$

1. vector addition: $\mathbf{v}_1 + \mathbf{v}_2 = \mathbf{v}_2 + \mathbf{v}_1$
2. scalar multiplication: $\begin{cases} a(\mathbf{v}_1 + \mathbf{v}_2) = a\mathbf{v}_1 + a\mathbf{v}_2 \\ (a + b)\mathbf{v} = a\mathbf{v} + b\mathbf{v} \end{cases} \quad \forall a, b \in \mathbb{R}$
3. inner product (dot product): $\mathbf{v}_1^T \mathbf{v}_2 = \mathbf{v}_1 \cdot \mathbf{v}_2 = \mathbf{v}_2 \cdot \mathbf{v}_1 = \mathbf{v}_2^T \mathbf{v}_1$
4. dyadic: $\mathbf{v}_1 \mathbf{v}_2 \neq \mathbf{v}_2 \mathbf{v}_1$
5. basis: $\mathbf{e}_i \cdot \mathbf{e}_j = \delta_{ij} = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases} \longrightarrow \text{orthonormal} \quad \mathbf{e}_i \mathbf{e}_i = \mathbf{I} = \begin{pmatrix} 1 & & \\ & 1 & \\ & & 1 \end{pmatrix}$
6. $\mathbf{v}_1 \cdot \mathbf{v}_2 = r \in \mathbb{R}$
 $\mathbf{v} \cdot \mathbf{v} = r \in \mathbb{R}, r \geq 0$ (equality iff $\mathbf{v} = \mathbf{0}$)

5.2 State Vector Space \mathbb{H} (Hilbert Space)

$$|\psi\rangle = \begin{pmatrix} \vdots \\ \vdots \\ \vdots \end{pmatrix}, \quad \langle\psi| = (\cdots \quad \cdots \quad \cdots)$$

1. vector addition: $|\psi_1\rangle + |\psi_2\rangle = |\psi_2\rangle + |\psi_1\rangle$
2. scalar multiplication: $\begin{cases} c(|\psi_1\rangle + |\psi_2\rangle) = c|\psi_1\rangle + c|\psi_2\rangle \\ (c + c')|\psi\rangle = c|\psi\rangle + c'|\psi\rangle \end{cases} \quad \forall c \in \mathbb{C}$
3. inner product: $\langle\psi_1|\psi_2\rangle = (\langle\psi_2|\psi_1\rangle)^*$
4. outer product: $|\psi\rangle\langle\phi| \neq |\phi\rangle\langle\psi|$

$$(|\psi\rangle\langle\phi|)^\dagger = (\langle\phi|)^\dagger(|\psi\rangle)^\dagger = |\phi\rangle\langle\psi|$$

5. basis: for Hermitian operation $\begin{cases} \hat{A}|\psi_n\rangle = A_n|\psi_n\rangle & \text{discrete} \\ \hat{A}|a\rangle = a|a\rangle & \text{continuous} \end{cases}$

$$\text{discrete: } \begin{cases} \langle\psi_n|\psi_m\rangle = \delta_{nm} = \begin{cases} 1, & n = m \\ 0, & n \neq m \end{cases} \\ \sum_n |\psi_n\rangle\langle\psi_n| = \hat{I} \end{cases}$$

$$\text{continuous: } \begin{cases} \langle a|a'\rangle = \delta(a - a') = \begin{cases} \infty, & a = a' \\ 0, & a \neq a' \end{cases} \\ \int |a\rangle\langle a| da = \hat{I} \end{cases}$$

6. $\langle\psi_1|\psi_2\rangle = c \in \mathbb{C}$
 $(\langle\psi_1|\psi_2\rangle)^* = \langle\psi_2|\psi_1\rangle$
 $\langle\psi|\psi\rangle = \|\psi\|^2 = r \in \mathbb{R}, r \geq 0$ (equality iff $|\psi\rangle = \mathbf{0}$)

6 Operator

$$\hat{A}|\phi\rangle = |\psi\rangle, \quad \hat{A}^{-1}|\psi\rangle = |\phi\rangle$$

6.1 Linear Operator

$$\forall |\psi_1\rangle, |\psi_2\rangle \in \mathbb{H}, a, b \in \mathbb{C}, \hat{A}(a|\psi_1\rangle + b|\psi_2\rangle) = a\hat{A}|\psi_1\rangle + b\hat{A}|\psi_2\rangle$$

then, \hat{A} is a linear operator

$$\hat{A}(a|\psi_1\rangle + b|\psi_2\rangle) = a^*\hat{A}|\psi_1\rangle + b^*\hat{A}|\psi_2\rangle$$

then, \hat{A} is an anti-linear operation

6.1.1 Two Kinds of Linear Operators

$$\begin{cases} \text{Hermitian operator: } \hat{A}^\dagger = \hat{A} \\ \text{Unitary operator: } \hat{U}^\dagger = \hat{U}^{-1} \end{cases}$$

6.1.2 Rules

$$\forall |\psi\rangle, \text{ if } \hat{A}|\psi\rangle = \hat{B}|\psi\rangle \iff \hat{A} = \hat{B}$$

$$1. \hat{A} + \hat{B} = \hat{B} + \hat{A}$$

$$\text{def: } \forall |\psi\rangle, \hat{A}|\psi\rangle + \hat{B}|\psi\rangle = (\hat{A} + \hat{B})|\psi\rangle$$

$$2. (\hat{A} + \hat{B}) + \hat{C} = \hat{A} + (\hat{B} + \hat{C})$$

$$3. \hat{A}\hat{B} \neq \hat{B}\hat{A}$$

$$\text{def commutator: } [\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} \longleftrightarrow [A, B]_{\text{PB}} \longleftrightarrow \frac{1}{i\hbar} [\hat{A}, \hat{B}]$$

$$4. \text{ adjoint (Hermitian conjugate): } (\hat{A}|\phi\rangle)^\dagger = \langle\phi| \hat{A}^\dagger \iff \langle\phi| \hat{A}^\dagger |\psi\rangle = (\langle\psi| \hat{A} |\phi\rangle)^* \\ \implies \text{Hermitian operation: } \hat{A}^\dagger = \hat{A} \text{ (Hermitian matrix)} \implies A_{ij} = A_{ji}^*$$

6.1.3 Properties

$$1. \text{ Identity: } \forall |\psi\rangle, \exists \hat{I}, \hat{I}|\psi\rangle = |\psi\rangle$$

$$2. \text{ Inverse: } \hat{A}|\psi\rangle = |\phi\rangle \implies |\psi\rangle = \hat{A}^{-1}|\phi\rangle, \hat{A}\hat{A}^{-1} = \hat{A}^{-1}\hat{A} = \hat{I} \\ (\hat{A}_1\hat{A}_2\cdots\hat{A}_n)^{-1} = \hat{A}_n^{-1}\cdots\hat{A}_2^{-1}\hat{A}_1^{-1}$$

$$3. (\hat{A}_1\hat{A}_2\cdots\hat{A}_n)^\dagger = \hat{A}_n^\dagger\cdots\hat{A}_2^\dagger\hat{A}_1^\dagger$$

7 Overview of QM

ROSE P... M

8 Roles: State Vector, Operator

8.0.1 Operator

$$\begin{cases} \text{Observable} \\ \text{Transformation} \end{cases} \quad \begin{cases} \text{Hermitian: } \hat{A}^\dagger = \hat{A} \\ \text{Unitary: } \hat{A}^\dagger = \hat{A}^{-1} \end{cases}$$

8.0.2 State

$$\begin{cases} \text{ket } |\psi\rangle \text{ (vector): } (|\psi\rangle)^\dagger = \langle\psi| \\ \text{bra } \langle\psi| \text{ (dual vector): } (\langle\psi|)^\dagger = |\psi\rangle \end{cases}$$

8.0.3 Eigenspectrum

$$\begin{cases} \text{discrete (bound state)} \\ \text{continuous (unbound state)} \end{cases}$$

8.1 Properties

Wave Particle Duality

8.2 Problems

8.2.1 Eigenvalue Problem

$$\hat{A}|\psi\rangle = \lambda|\psi\rangle \begin{cases} \hat{A}|u_n\rangle = A_n|u_n\rangle \begin{cases} \langle u_n|u_m\rangle = \delta_{nm} \\ \sum_n |u_n\rangle\langle u_n| = \hat{I} \end{cases} \\ \hat{A}|a\rangle = a|a\rangle \begin{cases} \langle a|a'\rangle = \delta(a-a') \\ \int da |a\rangle\langle a| = \hat{I} \end{cases} \end{cases}$$

8.2.2 Evolution Problem

$$\begin{cases} i\hbar \frac{d}{dt} |\psi(t)\rangle = \hat{H} |\psi(t)\rangle \text{ Schrödinger eq.} \\ \frac{d\hat{F}}{dt} = \frac{\partial \hat{F}}{\partial t} + \frac{1}{i\hbar} [\hat{F}, \hat{H}] \text{ Heisenberg eq.} \end{cases}$$

8.3 Picture

8.3.1 Schrödinger Picture

$$\frac{d\hat{F}}{dt} = \frac{\partial \hat{F}}{\partial t} \equiv 0$$

8.3.2 Heisenberg Picture

$$\frac{d}{dt} |\psi(t)\rangle \equiv 0$$

8.4 Particles

$$\begin{cases} \text{Boson: } s=\text{integer} \longrightarrow \text{Bose-Einstein Statistics} \\ \text{Femion: } s=\text{half-integer} \longrightarrow \text{Fermi-Dirac Statistics} \end{cases} \xrightarrow{\hbar \rightarrow 0} \text{Maxwell-Boltzmann Statistics}$$

8.5 Perturbation Theories

$$\begin{cases} \text{Non-degenerate} \\ \text{Degenerate} \end{cases}$$

8.6 Angular Momentum and Magnetic Moment

$$\begin{cases} \text{Orbital Angular Momentum} \\ \text{Spin Angular Momentum} \end{cases} \iff \begin{cases} \text{Orbital Magnetic Moment} \\ \text{Intrinsic Magnetic Moment} \end{cases}$$

8.7 Mechanics

$$\begin{cases} \text{Wave Mechanics: wavefunction} \longleftrightarrow \text{state vector} \\ \text{Matrix Mechanics: matrix forms of} \begin{cases} \text{state vector} \\ \text{operator: } n \times n \text{ matrix} \end{cases} \end{cases}$$

8.8 Theorem of Hermitian Operator

$$\hat{A}^\dagger = \hat{A}, \quad \hat{A} |\psi_n\rangle = A_n |\psi_n\rangle \quad (\text{non-degenerate, discrete})$$

1. eigenvalue $A_n \in \mathbb{R}$
2. eigenstate \longrightarrow orthonormal and complete basis

$$\begin{aligned} \langle \psi_n | \hat{A}^\dagger | \psi_n \rangle &= (\langle \psi_n | \hat{A} | \psi_n \rangle)^* = A_n^* \langle \psi_n | \psi_n \rangle = \langle \psi_n | \hat{A} | \psi_n \rangle = A_n \langle \psi_n | \psi_n \rangle \\ &\implies A_n^* = A_n \implies A_n \in \mathbb{R} \end{aligned}$$

$$\begin{cases} \hat{A} |\psi_n\rangle = A_n |\psi_n\rangle \\ \hat{A} |\psi_m\rangle = A_m |\psi_m\rangle \end{cases} \implies \langle \psi_n | \hat{A}^\dagger | \psi_m \rangle = (\langle \psi_m | \hat{A} | \psi_n \rangle)^* = A_n^* (\langle \psi_m | \psi_n \rangle)^* = A_n^* \langle \psi_n | \psi_m \rangle$$

$$\begin{aligned} \langle \psi_n | \hat{A} | \psi_m \rangle &= A_m \langle \psi_n | \psi_m \rangle \implies (A_n - A_m) \langle \psi_n | \psi_m \rangle = 0 \\ n \neq m &\implies A_n \neq A_m \implies \langle \psi_n | \psi_m \rangle = 0 \implies |\psi_n\rangle \perp |\psi_m\rangle \end{aligned}$$

Remark: Degenerate,

$$\hat{A} |\psi_n^i\rangle = A_n |\psi_n^i\rangle \quad (i = 1, 2, \dots, \underset{\text{degeneracy}}{g_n}) \implies |\psi_n^i\rangle \neq c |\psi_n^j\rangle \quad (\text{linear independent})$$

$$|\psi_n^i\rangle \xrightarrow{\text{Gram-Schmidt Orthogonalization}} |\phi_n^i\rangle \left\{ \begin{array}{l} \langle \phi_n^i | \phi_m^j \rangle = \delta_{nm} \delta_{ij} \\ \sum_{n,i} |\phi_n^i\rangle \langle \phi_n^i| = \hat{1} \end{array} \right.$$

9 Wave Mechanics & Matrix Mechanics

9.1 Wave Mechanics: wavefunction

$$\text{wavefunction} \left\{ \begin{array}{l} \text{matter wave (Schrödinger)} \\ \text{probabilistic wave (Born): } |\Psi(\mathbf{x}, t)|^2 \sim \text{probability density} \end{array} \right.$$

wavefunction is the projection of state vector in representation
basis

$$|\psi\rangle = \hat{1} |\psi\rangle = \left\{ \begin{array}{l} \left(\sum_n |u_n\rangle \langle u_n| \right) |\psi\rangle = \sum_n c_n |u_n\rangle, \quad c_n = \langle u_n | \psi \rangle \quad \text{discrete} \\ \left(\int da |a\rangle \langle a| \right) |\psi\rangle = \int da \psi(a) |a\rangle, \quad \psi(a) = \langle a | \psi \rangle \quad \text{continuous} \end{array} \right.$$

c_n is the wavefunction in $\{|u_n\rangle\}$ -representation, $\psi(a)$ is the wavefunction in $\{|a\rangle\}$ -representation.

$$|\psi\rangle = \sum_n c_n |\psi_n\rangle, \quad c_n = \langle \psi_n | \psi \rangle, \quad \hat{A} |\psi\rangle = A_n |\psi_n\rangle$$

$$\text{continuous} \left\{ \begin{array}{l} \mathbf{x}\text{-representation: } \hat{\mathbf{x}} |\mathbf{x}\rangle = \mathbf{x} |\mathbf{x}\rangle \left\{ \begin{array}{l} \langle \mathbf{x} | \mathbf{x}' \rangle = \delta(\mathbf{x} - \mathbf{x}') \\ \int d^3x |\mathbf{x}\rangle \langle \mathbf{x}| = \hat{1} \end{array} \right. \\ \mathbf{p}\text{-representation: } \hat{\mathbf{p}} |\mathbf{p}\rangle = \mathbf{p} |\mathbf{p}\rangle \left\{ \begin{array}{l} \langle \mathbf{p} | \mathbf{p}' \rangle = \delta(\mathbf{p} - \mathbf{p}') \\ \int d^3p |\mathbf{p}\rangle \langle \mathbf{p}| = \hat{1} \end{array} \right. \end{array} \right.$$

Energy-representation, Fork-state representation (occupation number representation)

$$\begin{aligned} |\psi\rangle &= \left(\int d^3x |\mathbf{x}\rangle \langle \mathbf{x}| \right) |\psi\rangle = \int d^3x \psi(\mathbf{x}) |\mathbf{x}\rangle \\ &= \left(\int d^3p |\mathbf{p}\rangle \langle \mathbf{p}| \right) |\psi\rangle = \int d^3p \tilde{\psi}(\mathbf{p}) |\mathbf{p}\rangle \end{aligned}$$

$$\psi(\mathbf{x}) = \langle \mathbf{x} | \psi \rangle \quad \tilde{\psi}(\mathbf{p}) = \langle \mathbf{p} | \psi \rangle$$

$$\begin{aligned} \tilde{\psi}(\mathbf{p}) &= \langle \mathbf{p} | \psi \rangle = \langle \mathbf{p} | \hat{1} | \psi \rangle = \langle \mathbf{p} | \left(\int d^3x |\mathbf{x}\rangle \langle \mathbf{x}| \right) | \psi \rangle = \int d^3x \langle \mathbf{p} | \mathbf{x} \rangle \psi(\mathbf{x}) \\ &= \int d^3x \psi(\mathbf{x}) \psi_{\mathbf{p}}^*(\mathbf{x}) \quad \text{where } \psi_{\mathbf{p}}(\mathbf{x}) = \langle \mathbf{x} | \mathbf{p} \rangle = \frac{1}{(2\pi\hbar)^{\frac{3}{2}}} e^{-i\mathbf{p}\cdot\mathbf{x}/\hbar} \\ &\implies \text{in } \mathbf{x}\text{-representation, } \hat{\mathbf{x}} = \mathbf{x}, \quad \hat{\mathbf{p}} = -i\hbar \frac{\partial}{\partial \mathbf{x}} \end{aligned}$$

$$\implies \begin{cases} i\hbar \frac{d}{dt} |\psi(t)\rangle = \hat{H} |\psi(t)\rangle & \xrightarrow{\int d^3x |\mathbf{x}\rangle \langle \mathbf{x}| = \hat{1}} i\hbar \frac{\partial}{\partial t} \Psi(\mathbf{x}, t) = -\frac{\hbar^2}{2m} \nabla^2 \Psi(\mathbf{x}, t) + V(\mathbf{x}) \Psi(\mathbf{x}, t) \\ \hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + V(\hat{\mathbf{x}}) \end{cases}$$

$$\text{where } \begin{cases} \Psi(\mathbf{x}, t) = \langle \mathbf{x} | \psi(t) \rangle \\ \hat{\mathbf{x}} = \mathbf{x}, \quad \hat{\mathbf{p}} = -i\hbar \nabla \end{cases}$$

$$\text{in } \mathbf{p}\text{-representation, } \hat{\mathbf{x}} = +i\hbar \frac{\partial}{\partial \mathbf{p}}, \quad \hat{\mathbf{p}} = \mathbf{p}$$

$$\text{Schrödinger eq. } \xrightarrow{\int d^3p |\mathbf{p}\rangle \langle \mathbf{p}| = \hat{1}} i\hbar \frac{\partial}{\partial t} \tilde{\psi}(\mathbf{p}, t) = \frac{\mathbf{p}^2}{2m} \tilde{\psi}(\mathbf{p}, t) + V \left(i\hbar \frac{\partial}{\partial \mathbf{p}} \right) \tilde{\psi}(\mathbf{p}, t)$$

$$\text{where } \begin{cases} \tilde{\psi}(\mathbf{p}) = \langle \mathbf{p} | \psi(t) \rangle \\ \hat{\mathbf{p}} = \mathbf{p}, \quad \hat{\mathbf{x}} = +i\hbar \frac{\partial}{\partial \mathbf{p}} \end{cases}$$

9.2 Matrix Mechanics: State Vector

$$|\psi\rangle = \hat{1} |\psi\rangle = \sum_n c_n |u_n\rangle = \begin{pmatrix} c_1 \\ c_2 \\ \vdots \\ c_n \\ \vdots \end{pmatrix} = (c_1 \quad c_2 \quad \cdots \quad c_n \quad \cdots)^T$$

$$\hat{A} = \mathbf{I} \hat{A} \mathbf{I} = \left(\sum_n |u_n\rangle \langle u_n| \right) \hat{A} \left(\sum_m |u_m\rangle \langle u_m| \right) = \sum_{n,m} \underbrace{\langle u_n | \hat{A} | u_m \rangle}_{=A_{nm}} (|u_n\rangle \langle u_m|) = \begin{pmatrix} & A_{nm} \end{pmatrix}$$

9.3 Matrix Mechanics: Hermitian Conjugate

if the operator \hat{A} can be written as a matrix,

$$\langle \psi | \hat{A}^\dagger | \phi \rangle = \left(\langle \phi | \hat{A} | \psi \rangle \right)^* \implies \hat{A}^\dagger = \left(\hat{A}^T \right)^*$$

$$\text{Pf. Let } \begin{cases} |\psi\rangle = \sum_n c_n |u_n\rangle \\ |\phi\rangle = \sum_n d_n |u_n\rangle \end{cases},$$

$$\begin{aligned} \langle\psi|\hat{A}^\dagger|\phi\rangle &= \left(\sum_n c_n^* \langle u_n|\right) \hat{A}^\dagger \left(\sum_m d_m |u_m\rangle\right) = \sum_{n,m} c_n^* d_m \langle u_n|\hat{A}^\dagger|u_m\rangle = \sum_{n,m} c_n^* d_m (\hat{A}^\dagger)_{nm} \\ (\langle\phi|\hat{A}|\psi\rangle)^* &= \left[\left(\sum_n d_n^* \langle u_n|\right) \hat{A} \left(\sum_m c_m |u_m\rangle\right)\right]^* = \left(\sum_{n,m} d_n^* c_m \langle u_n|\hat{A}|u_m\rangle\right)^* \\ &= \sum_{n,m} d_n c_m^* A_{nm}^* = \sum_{n,m} d_m c_n^* A_{mn}^* = \sum_{n,m} c_n^* d_m A_{mn}^* \implies (\hat{A}^\dagger)_{nm} = (\hat{A})_{mn}^* = (\hat{A}^T)_{nm}^* \end{aligned}$$

10 Commutation

10.0.1 Definition

$$[\hat{A}, \hat{B}] = \hat{A}\hat{B} - \hat{B}\hat{A} \quad (1)$$

10.0.2 Compare

$$\begin{cases} \text{q-number : } \hat{A}\hat{B} \neq \hat{B}\hat{A} & (\text{QM}) \\ \text{c-number : } AB = BA & (\text{CM}) \end{cases}$$

$$[A, B]_{\text{PB}} = \sum_{\alpha} \left(\frac{\partial A}{\partial q_{\alpha}} \frac{\partial B}{\partial p_{\alpha}} - \frac{\partial A}{\partial p_{\alpha}} \frac{\partial B}{\partial q_{\alpha}} \right) \quad (2)$$

Classical Canonical Commutator

10.0.3 Relation

$$\boxed{[A, B]_{\text{PB}} \longleftrightarrow \frac{1}{i\hbar} [\hat{A}, \hat{B}]} \quad (3)$$

11 Properties of Commutations

$$[\hat{A}, \hat{B}] \neq 0 \implies \text{Uncertainty Relation}$$

$$\Delta\hat{A} \cdot \Delta\hat{B} \geq \frac{|\langle [\hat{A}, \hat{B}] \rangle|}{2} \quad (4)$$

$$[\hat{A}, \hat{B}] = 0 \implies \begin{matrix} \text{Common} \\ \text{Simultaneous} \end{matrix} \text{ Eigenstates}$$

1. **Linearity**

$$[a\hat{A}, b\hat{B} + c\hat{C}] = ab[\hat{A}, \hat{B}] + ac[\hat{A}, \hat{C}] \quad (5)$$

2. **Leibniz rule**

$$\begin{cases} [\hat{A}, \hat{B}\hat{C}] = [\hat{A}, \hat{B}]\hat{C} + \hat{B}[\hat{A}, \hat{C}] \\ [\hat{A}\hat{B}, \hat{C}] = [\hat{A}, \hat{C}]\hat{B} + \hat{A}[\hat{B}, \hat{C}] \end{cases} \quad (6)$$

3. **Power related**

$$[\hat{A}, \hat{B}^n] = n[\hat{A}, \hat{B}]\hat{B}^{n-1} \quad (7)$$

provided that $[\hat{B}, [\hat{A}, \hat{B}]] = 0$

Pf. $[\hat{A}, \hat{B}] = 1[\hat{A}, \hat{B}]\hat{B}^{1-1}$ ($n = 1$). Suppose $[\hat{A}, \hat{B}^k] = k[\hat{A}, \hat{B}]\hat{B}^{k-1}$, then when $n = k + 1$, one has $[\hat{A}, \hat{B}^{k+1}] = [\hat{A}, \hat{B}^k]\hat{B} + \hat{B}^k[\hat{A}, \hat{B}] = (k + 1)[\hat{A}, \hat{B}]\hat{B}^k \Rightarrow [\hat{A}, \hat{B}^n] = n[\hat{A}, \hat{B}]\hat{B}^{n-1}$

4.

$$[\hat{A}, f(\hat{B})] = [\hat{A}, \hat{B}] \frac{\partial f}{\partial \hat{B}} \quad (8)$$

provided that $[\hat{B}, [\hat{A}, \hat{B}]] = 0$

$$\begin{aligned} \text{Pf. } [\hat{A}, f(\hat{B})] &= \left[\hat{A}, \sum_n \frac{f^{(n)}(0)}{n!} \hat{B}^n \right] = \sum_n \frac{f^{(n)}(0)}{n!} [\hat{A}, \hat{B}^n] = \sum_n \frac{f^{(n)}(0)}{(n-1)!} \hat{B}^{n-1} [\hat{A}, \hat{B}] \\ &= \sum_n \frac{f^{(n)}(0)}{n!} [\hat{A}, \hat{B}] \hat{B}^n = [\hat{A}, \hat{B}] \frac{\partial f}{\partial \hat{B}} \end{aligned}$$

5. **Jacobi identity** \rightarrow Lie Algebra

$$[A, [B, C]_{\text{PB}}]_{\text{PB}} + [B, [C, A]_{\text{PB}}]_{\text{PB}} + [C, [A, B]_{\text{PB}}]_{\text{PB}} = 0 \quad (\text{CM}) \quad (9)$$

$$\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) + \mathbf{B} \times (\mathbf{C} \times \mathbf{A}) + \mathbf{C} \times (\mathbf{A} \times \mathbf{B}) = 0 \quad (\text{VA}) \quad (10)$$

$$\mathbf{F}_{\mu\nu, \lambda} + \mathbf{F}_{\nu\lambda, \mu} + \mathbf{F}_{\lambda\mu, \nu} = 0 \quad (\text{CED}) \quad (11)$$

$$[\hat{A}, [\hat{B}, \hat{C}]] + [\hat{B}, [\hat{C}, \hat{A}]] + [\hat{C}, [\hat{A}, \hat{B}]] = 0 \quad (\text{QM}) \quad (12)$$

12 Fundamental Commutation Relations

$$\text{CM} \begin{cases} [x_i, p_j]_{\text{PB}} = \delta_{ij} \\ [x_i, x_j]_{\text{PB}} = [p_i, p_j]_{\text{PB}} = 0 \end{cases} \longleftrightarrow \text{QM} \begin{cases} [\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij} \\ [\hat{x}_i, \hat{x}_j] = [\hat{p}_i, \hat{p}_j] = 0 \end{cases} \quad (13)$$

Pf. in \mathbf{x} -representation, $\hat{\mathbf{x}} = \mathbf{x}$, $\hat{\mathbf{p}} = -i\hbar\nabla \implies \hat{x} = x$, $\hat{p}_x = -i\hbar\frac{\partial}{\partial x}$. $\forall\psi(x)$,

$$\begin{aligned} [\hat{x}, \hat{p}_x]\psi(x) &= (\hat{x}\hat{p}_x - \hat{p}_x\hat{x})\psi(x) = -i\hbar\left(x\frac{\partial}{\partial x} - \frac{\partial}{\partial x}x\right)\psi(x) = -i\hbar\left[x\frac{\partial\psi}{\partial x} - \frac{\partial}{\partial x}(x\psi)\right] \\ &= i\hbar\psi(x) \implies [\hat{x}, \hat{p}_x] = i\hbar. \end{aligned}$$

Similarly, $[\hat{y}, \hat{p}_y] = [\hat{z}, \hat{p}_z] = i\hbar$. $[\hat{x}_i, \hat{p}_j] = 0$, ($i \neq j$).

13 Commutation Relations of Orbital Angular Momentum

$$\hat{\mathbf{L}} = \hat{\mathbf{x}} \times \hat{\mathbf{p}} \iff \hat{L}_i = \varepsilon_{ijk}\hat{x}_j\hat{p}_k \quad (14)$$

$$\implies \boxed{[\hat{L}_i, \hat{L}_j] = i\hbar\varepsilon_{ijk}\hat{L}_k \iff \hat{\mathbf{L}} \times \hat{\mathbf{L}} = i\hbar\hat{\mathbf{L}}} \quad (15)$$

$$\implies [\hat{L}_i, \hat{\mathbf{L}}^2] = 0 \implies [\hat{L}_z, \hat{\mathbf{L}}^2] = 0, z: \text{quantum axis} \quad (16)$$

$$\begin{cases} [\hat{L}_i, \hat{x}_j] = i\hbar\varepsilon_{ijk}\hat{x}_k \iff \hat{\mathbf{L}} \times \hat{\mathbf{x}} + \hat{\mathbf{x}} \times \hat{\mathbf{L}} = 2i\hbar\hat{\mathbf{x}} \\ [\hat{L}_i, \hat{p}_j] = i\hbar\varepsilon_{ijk}\hat{p}_k \iff \hat{\mathbf{L}} \times \hat{\mathbf{p}} + \hat{\mathbf{p}} \times \hat{\mathbf{L}} = 2i\hbar\hat{\mathbf{p}} \end{cases} \quad (17)$$

$$\forall \text{ vector } \hat{\mathbf{v}}, [\hat{L}_i, \hat{v}_j] = i\hbar\varepsilon_{ijk}\hat{v}_k \iff \hat{\mathbf{L}} \times \hat{\mathbf{v}} + \hat{\mathbf{v}} \times \hat{\mathbf{L}} = 2i\hbar\hat{\mathbf{v}} \quad (18)$$

$$\text{for } V(\mathbf{x}) = V(r), [\hat{L}_i, V(r)] = 0 \quad (19)$$

$$\text{for } \hat{T} = \frac{\hat{\mathbf{p}}^2}{2m}, [\hat{L}_i, \hat{T}] = 0 \quad (20)$$

$$\text{for } \hat{H} = \hat{T} + V(r), [\hat{L}_i, \hat{H}] = [\hat{\mathbf{L}}^2, \hat{H}] = 0 \quad (21)$$

$$\text{Pf. } (\hat{\mathbf{L}} \times \hat{\mathbf{L}})_i = \varepsilon_{ijk}\hat{L}_j\hat{L}_k = \frac{1}{2}\varepsilon_{ijk}\hat{L}_j\hat{L}_k - \frac{1}{2}\varepsilon_{ijk}\hat{L}_k\hat{L}_j = \frac{1}{2}\varepsilon_{ijk}[\hat{L}_j, \hat{L}_k]$$

$$= \frac{1}{2}\varepsilon_{ijk}(i\hbar)\varepsilon_{jkl}\hat{L}_l = i\hbar\delta_{il}\hat{L}_l = i\hbar\hat{L}_i \implies \hat{\mathbf{L}} \times \hat{\mathbf{L}} = i\hbar\hat{\mathbf{L}}$$

$$\begin{aligned} [\hat{L}_i, \hat{L}_j] &= i\hbar\varepsilon_{ijk}\hat{L}_k \implies [\hat{L}_i, \hat{\mathbf{L}}^2] = [\hat{L}_i, \hat{L}_j\hat{L}_j] = [\hat{L}_i, \hat{L}_j]\hat{L}_j + \hat{L}_j[\hat{L}_i, \hat{L}_j] \\ &= i\hbar\varepsilon_{ijk}(\hat{L}_k\hat{L}_j + \hat{L}_j\hat{L}_k) = 0 \end{aligned}$$

$$[\hat{L}_i, \hat{x}_j] = [\varepsilon_{iab}\hat{x}_a\hat{p}_b, \hat{x}_j]$$

$$[\hat{L}_i, V(r)] = [\hat{L}_i, \hat{x}_j]\frac{\partial V}{\partial x_j} = i\hbar\varepsilon_{ijk}\hat{x}_k\frac{dV}{dr}\frac{\hat{x}_j}{r} = \left(\frac{i\hbar}{r}\frac{dV}{dr}\right)\varepsilon_{ijk}\hat{x}_k\hat{x}_j = 0$$

$$[\hat{L}_i, \hat{\mathbf{p}}^2] = [\hat{L}_i, \hat{p}_j\hat{p}_j] = [\hat{L}_i, \hat{p}_j]\hat{p}_j + \hat{p}_j[\hat{L}_i, \hat{p}_j] = i\hbar\varepsilon_{ijk}(\hat{p}_k\hat{p}_j + \hat{p}_j\hat{p}_k) = 0$$

14 Eigenvalue Problem

15 Observables

1. Position (canonical)

$$\hat{\mathbf{x}} = \begin{cases} \mathbf{x} & \mathbf{x}\text{-representation} \\ +i\hbar \frac{\partial}{\partial \mathbf{p}} & \mathbf{p}\text{-representation} \end{cases} \quad (22)$$

2. Momentum (canonical)

$$\hat{\mathbf{p}} = \begin{cases} -i\hbar \frac{\partial}{\partial \mathbf{x}} & \mathbf{x}\text{-representation} \\ \mathbf{p} & \mathbf{p}\text{-representation} \end{cases} \quad (23)$$

$$\text{Fundamental Commutation Relation } \begin{cases} [\hat{x}_i, \hat{p}_j] = i\hbar \delta_{ij} \\ [\hat{x}_i, \hat{x}_j] = [\hat{p}_i, \hat{p}_j] = 0 \end{cases}$$

3. Orbital Angular Momentum

$$\hat{\mathbf{L}} = \hat{\mathbf{x}} \times \hat{\mathbf{p}} \text{ or } \hat{L}_i = \varepsilon_{ijk} \hat{x}_j \hat{p}_k$$

$$\text{Commutation Relations } [\hat{L}_i, \hat{L}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k \implies [\hat{L}_i, \hat{\mathbf{L}}^2]$$

Remark $\hat{\mathbf{L}} \times \hat{\mathbf{L}} \neq 0$, but $\hat{\mathbf{x}} \times \hat{\mathbf{p}} = -\hat{\mathbf{p}} \times \hat{\mathbf{x}}$

4. Intrinsic Angular Momentum (No classical correspondence) - SPIN

$$\begin{aligned} [\hat{S}_i, \hat{S}_j] &= i\hbar \varepsilon_{ijk} \hat{S}_k \iff \hat{\mathbf{S}} \times \hat{\mathbf{S}} = i\hbar \hat{\mathbf{S}} \\ [\hat{L}_i, \hat{S}_j] &= 0 \end{aligned}$$

5. Hamiltonian

$$\hat{H} = \hat{T} + \hat{V} = \frac{\hat{\mathbf{p}}^2}{2m} + \hat{V}(\mathbf{x}) \quad \begin{cases} \hat{T} : & \mathbf{p}\text{-dependent only} \\ \hat{V} : & \mathbf{x}\text{-dependent only} \end{cases}$$

6. Energy

$$\begin{cases} \hat{E} = i\hbar \frac{\partial}{\partial t} & \text{wavefunctions} \\ \hat{E} = i\hbar \frac{d}{dt} & \text{state vector} \end{cases}$$

Remark In non-relativistic QM, time t is NOT observable.

16 Eigenvalue & Eigenstates

1. Position

$$\begin{aligned}\hat{\mathbf{x}} |\mathbf{x}'\rangle &= \mathbf{x}' |\mathbf{x}'\rangle \\ \text{eigenvalue eq.} \\ \begin{cases} \langle \mathbf{x}' | \mathbf{x}'' \rangle = \delta(\mathbf{x}' - \mathbf{x}'') & (\text{ON}) \\ \int d^3x |\mathbf{x}\rangle \langle \mathbf{x}| = \hat{1} & (\text{RI}) \end{cases} \\ \langle \mathbf{x} | \hat{\mathbf{x}} | \mathbf{x}' \rangle &= \mathbf{x}' \delta(\mathbf{x} - \mathbf{x}')\end{aligned}$$

In $\hat{\mathbf{x}}$ -representation, wavefunction of $|\mathbf{x}\rangle$ is $\langle \mathbf{x}' | \mathbf{x} \rangle = \delta(\mathbf{x} - \mathbf{x}')$, and eigenvalue eq. is

$$\hat{\mathbf{x}} \delta(\mathbf{x} - \mathbf{x}') = \mathbf{x}' \delta(\mathbf{x} - \mathbf{x}')$$

2. Momentum

$$\begin{aligned}\hat{\mathbf{p}} |\mathbf{p}'\rangle &= \mathbf{p}' |\mathbf{p}'\rangle \\ \begin{cases} \langle \mathbf{p}' | \mathbf{p}'' \rangle = \delta(\mathbf{p}' - \mathbf{p}'') & (\text{ON}) \\ \int d\mathbf{p} |\mathbf{p}\rangle \langle \mathbf{p}| = \mathbf{I} & (\text{RI}) \end{cases}\end{aligned}$$

In $\hat{\mathbf{p}}$ -representation, wavefunction of $|\mathbf{p}\rangle$ is $\langle \mathbf{p}' | \mathbf{p} \rangle = \delta(\mathbf{p} - \mathbf{p}')$, and eigenvalue eq. is

$$\hat{\mathbf{p}} \delta(\mathbf{p} - \mathbf{p}') = \mathbf{p}' \delta(\mathbf{p} - \mathbf{p}')$$

Remark eigenspectrum: set of eigenvalues of observable

3. Relation

$$\psi(\mathbf{x}) = \psi_{\mathbf{p}}(\mathbf{x}) \tilde{\psi}(\mathbf{p}), \quad \psi_{\mathbf{p}}(\mathbf{x}) = \langle \mathbf{x} | \mathbf{p} \rangle$$

$$|\psi\rangle = \hat{I} |\psi\rangle = \begin{cases} \int d\mathbf{x} |\mathbf{x}\rangle \langle \mathbf{x} | \psi \rangle = \int d\mathbf{x} |\mathbf{x}\rangle \psi(\mathbf{x}) \\ \int d\mathbf{p} |\mathbf{p}\rangle \langle \mathbf{p} | \psi \rangle = \int d\mathbf{p} |\mathbf{p}\rangle \tilde{\psi}(\mathbf{p}) \end{cases} = \begin{cases} \psi(\mathbf{x}) & \hat{\mathbf{x}}\text{-representation} \\ \tilde{\psi}(\mathbf{p}) & \hat{\mathbf{p}}\text{-representation} \end{cases}$$

$$\begin{aligned}|\psi\rangle &= \int d\mathbf{p} |\mathbf{p}\rangle \tilde{\psi}(\mathbf{p}) = \int d\mathbf{x} |\mathbf{x}\rangle \left(\langle \mathbf{x} | \mathbf{p} \rangle \tilde{\psi}(\mathbf{p}) \right) = \int d\mathbf{x} |\mathbf{x}\rangle \psi(\mathbf{x}) \\ \implies \psi(\mathbf{x}) &= \langle \mathbf{x} | \mathbf{p} \rangle \tilde{\psi}(\mathbf{p}) = \psi_{\mathbf{p}}(\mathbf{x}) \tilde{\psi}(\mathbf{p})\end{aligned}$$

$$\begin{aligned}\hat{\mathbf{p}} |\mathbf{p}\rangle &= \mathbf{p} |\mathbf{p}\rangle \implies \int d\mathbf{x} |\mathbf{x}\rangle \langle \mathbf{x} | \hat{\mathbf{p}} | \mathbf{p} \rangle = \mathbf{p} \int d\mathbf{x} |\mathbf{x}\rangle \langle \mathbf{x} | \mathbf{p} \rangle \quad (\hat{\mathbf{p}} = -i\hbar \frac{\partial}{\partial \mathbf{x}}) \\ \implies -i\hbar \frac{\partial}{\partial \mathbf{x}} \psi_{\mathbf{p}}(\mathbf{x}) &= \mathbf{p} \psi_{\mathbf{p}}(\mathbf{x})\end{aligned}$$

$$-i\hbar \frac{d}{dx} \psi_p(x) = p \psi_p(x) \implies \psi_p(x) = N \exp\left(\frac{i}{\hbar} p x\right) \implies \psi_{\mathbf{p}}(\mathbf{x}) = N \exp\left(\frac{i}{\hbar} \mathbf{p} \cdot \mathbf{x}\right)$$

17 Spin and Pauli Matrices

17.1 Stern-Gerlach Experiment

17.1.1 Meanings

1. Quantization of Spatial Orientation of Angular Momentum
2. Intrinsic Angular Momentum (Spin)
3. S-G experiment \longrightarrow Rabi oscillation \longrightarrow Ramsey (atom clock, Hydrogen maser)
 \longrightarrow Quantum measurement

17.1.2 Analysis

1. (Silver) neutral atom beam \implies no Lorentz force
2. Magnetic field is inhomogeneous

$$\mathbf{F} = -\nabla W_m = \boldsymbol{\mu} \cdot \nabla \mathbf{B} \quad (24)$$

$$W_m = -\boldsymbol{\mu} \cdot \mathbf{B} \quad (25)$$

$\boldsymbol{\mu}$: magnetic dipole moment

3. Deflection: two values $\begin{cases} \text{up} & 50\% \\ \text{down} & 50\% \end{cases}$
4. furnace: low-temperature $\longrightarrow \mathbf{v}$ is small $\longrightarrow \mathbf{p}$ is small $\longrightarrow \mathbf{L}$ is small $\xrightarrow{\boldsymbol{\mu}_L \propto \mathbf{L}}$ $\boldsymbol{\mu}_L$ is small \implies Intrinsic freedom?

17.1.3 History

1921-1922 Stern, Gerlach - S-G experiment

1924 Pauli - Two-valuedness not described classically \longrightarrow Pauli exclusion principle

1925 Kronig - Self-rotation of electron (unpublished)

1925 Uhlenbeck, Goudsmit - Self-rotation

1927 Pauli - Pauli matrices, Pauli equation (wavefunction is a spinor with 2-component)

1928 Dirac - Relativistic QM, Dirac equation (4-component spinor)

17.1.4 Cascaded S-G Experiment

$$\text{Ag} \xrightarrow{\uparrow \mathbf{B}} \mu_z \rightarrow \begin{cases} +\mu_B & \text{up } 50\% \\ -\mu_B & \text{down } 50\% \end{cases} \quad \mu_B = \frac{e\hbar}{2mc} \quad (\text{Bohr magneton})$$

$$\text{Ag} \longrightarrow \mu_x \rightarrow \begin{cases} |\mu_{x,+}\rangle \rightarrow \mu_z \rightarrow \begin{cases} +\mu_B & 50\% \\ -\mu_B & 50\% \end{cases} \\ |\mu_{x,-}\rangle \rightarrow \mu_z \rightarrow \begin{cases} +\mu_B & 50\% \\ -\mu_B & 50\% \end{cases} \end{cases}$$

18 Spin Angular Momentum

18.0.1 Significance

1. Spin is an Intrinsic Angular Momentum
2. Spin is a signature to distinguish two families

$$\begin{aligned}
 (a) \quad & \begin{cases} s = 0, 1, 2, \dots \text{ (integer)} & \text{Bosons} \\ s = \frac{1}{2}, \frac{3}{2}, \dots \text{ (half-integer)} & \text{Fermion} \end{cases} \\
 (b) \quad \text{Boson} & \begin{cases} \text{Statistics: Bose-Einstein statistics} \\ \text{Permutation: } \psi_S(\dots, x_i, \dots, x_j, \dots) = \psi_S(\dots, x_j, \dots, x_i, \dots) \\ & \text{unchanged} \end{cases} \\
 (c) \quad \text{Fermion} & \begin{cases} \text{Statistics: Fermi-Dirac statistics} \\ \text{Permutation: } \psi_A(\dots, x_i, \dots, x_j, \dots) = -\psi_A(\dots, x_j, \dots, x_i, \dots) \\ & \text{opposite sign} \end{cases} \\
 (d) \quad \text{Quantum Statistics} & \begin{cases} \text{Bose-Einstein} \\ \text{Fermi-Dirac} \end{cases} \xrightarrow{\hbar \rightarrow 0} \text{Maxwell-Boltzmann statistics}
 \end{aligned}$$

$$[\hat{S}_i, \hat{S}_j] = i\hbar \varepsilon_{ijk} \hat{S}_k \iff \hat{\mathbf{S}} \times \hat{\mathbf{S}} = i\hbar \hat{\mathbf{S}} \quad (26)$$

$$\implies [\hat{\mathbf{S}}^2, \hat{S}_i] = 0 \implies \begin{cases} \hat{\mathbf{S}}^2 |s, m_s\rangle = \hbar^2 s(s+1) |s, m_s\rangle \\ \hat{S}_z |s, m_s\rangle = \hbar m_s |s, m_s\rangle \end{cases} \quad \text{common eigenstates: } |s, m_s\rangle$$

$$\text{Dirac eq.} \longleftrightarrow s = \frac{1}{2} \quad (\text{electron \& positron, proton, neutron, } \dots) \quad (27)$$

$$\left| s = \frac{1}{2}, m_s = +\frac{1}{2} \right\rangle = |+\rangle = |\uparrow\rangle = |0\rangle = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (28)$$

$$\left| s = \frac{1}{2}, m_s = -\frac{1}{2} \right\rangle = |-\rangle = |\downarrow\rangle = |1\rangle = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (29)$$

$$\begin{cases} |0\rangle \langle 0| + |1\rangle \langle 1| = \mathbf{I} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \\ \langle 0|0\rangle = \langle 1|1\rangle = 1; \langle 1|0\rangle = \langle 0|1\rangle = 0 \end{cases}$$

$$\implies \text{In } \hat{S}_z\text{-representation} \begin{cases} \hat{S}_z |0\rangle = +\frac{\hbar}{2} |0\rangle \\ \hat{S}_z |1\rangle = -\frac{\hbar}{2} |1\rangle \end{cases} \implies \hat{\mathbf{S}} = \frac{\hbar}{2} \hat{\boldsymbol{\sigma}} \implies \begin{cases} \hat{\boldsymbol{\sigma}}_z |0\rangle = |0\rangle \\ \hat{\boldsymbol{\sigma}}_z |1\rangle = -|1\rangle \end{cases}$$

$$\implies \hat{\boldsymbol{\sigma}}_z = |0\rangle \langle 0| - |1\rangle \langle 1| = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

$$\hat{\mathbf{S}}^2 = \frac{3}{4} \hbar^2 \mathbf{I} \implies \hat{\boldsymbol{\sigma}}^2 = 3\mathbf{I} \implies [\hat{\boldsymbol{\sigma}}_i, \hat{\boldsymbol{\sigma}}_j] = 2i\varepsilon_{ijk} \hat{\boldsymbol{\sigma}}_k$$

19 Algebraic Method to Eigenvalue Problem of Angular Momentum

20 Review

Fundamental Commutations

$$[\hat{x}_i, \hat{p}_j] = i\hbar\delta_{ij} \quad [\hat{x}_i, \hat{x}_j] = [\hat{p}_i, \hat{p}_j] = 0 \quad (30)$$

20.1 Orbital Angular Momentum

20.1.1 Definition of Orbital AM

$$\hat{\mathbf{L}} = \hat{\mathbf{x}} \times \hat{\mathbf{p}} \quad \text{or} \quad \hat{L}_i = \varepsilon_{ijk} \hat{x}_j \hat{p}_k \quad (31)$$

20.1.2 Commutation Relations of Orbital AM

$$[\hat{L}_i, \hat{L}_j] = i\hbar\varepsilon_{ijk} \hat{L}_k \iff \hat{\mathbf{L}} \times \hat{\mathbf{L}} = i\hbar\hat{\mathbf{L}} \quad (32)$$

$$[\hat{L}^2, \hat{L}_i] = 0 \implies \text{Common Eigenstates} \quad (33)$$

20.1.3 Eigenvalue Equation

$$\begin{cases} \hat{\mathbf{L}}^2 |l, m_l\rangle = \hbar^2 l(l+1) |l, m_l\rangle \\ \hat{L}_z |l, m_l\rangle = \hbar m_l |l, m_l\rangle \end{cases} \quad m_l = \underbrace{-l, -l+1, \dots, l-1, l}_{2l+1} \quad (34)$$

20.2 Intrinsic Angular Momentum (Spin)

Generalize to Intrinsic AM (Spin):

$$[\hat{S}_i, \hat{S}_j] = i\hbar\varepsilon_{ijk} \hat{S}_k \iff \hat{\mathbf{S}} \times \hat{\mathbf{S}} = i\hbar\hat{\mathbf{S}} \quad (35)$$

$$[\hat{S}^2, \hat{S}_i] = 0 \implies \text{Common Eigenstates} \quad (36)$$

20.2.1 Eigenvalue Equation

$$\begin{cases} \hat{\mathbf{S}}^2 |s, m_s\rangle = \hbar^2 s(s+1) |s, m_s\rangle \\ \hat{S}_z |s, m_s\rangle = \hbar m_s |s, m_s\rangle \end{cases} \quad m_s = \underbrace{-s, -s+1, \dots, s-1, s}_{2s+1} \quad (37)$$

$$s = \frac{1}{2} \implies |s, m_s\rangle = \begin{cases} |+\rangle = |0\rangle = \begin{pmatrix} 1 & 0 \end{pmatrix}^T \\ |-\rangle = |1\rangle = \begin{pmatrix} 0 & 1 \end{pmatrix}^T \end{cases} \quad (38)$$

20.3 Pauli Matrix

$$\hat{\mathbf{S}} = \frac{\hbar}{2} \hat{\boldsymbol{\sigma}} \quad (39)$$

$$\begin{cases} \hat{\mathbf{S}}^2 = \frac{3}{4} \hbar^2 \hat{1}_2 \implies \hat{\boldsymbol{\sigma}}^2 = 3 \hat{1}_2 \\ \hat{S}_z^2 = \frac{1}{4} \hbar^2 \hat{1}_2 \implies \hat{\sigma}_z^2 = \hat{1}_2 \end{cases} \quad (40)$$

$$\text{Cascaded SGE} \implies \begin{cases} |\mu_x, +\rangle = c_1 |0\rangle + c_2 |1\rangle \\ |c_1|^2 = |c_2|^2 = \frac{1}{2} \end{cases} \implies \hat{\sigma}_x^2 = \hat{\sigma}_y^2 = \hat{1}_2$$

$$\hat{\sigma}_i = \hat{1}_2 \quad (41)$$

$$\{\hat{\sigma}_i, \hat{\sigma}_j\} = 2\delta_{ij} \hat{1}_2 \quad (42)$$

$$[\hat{\sigma}_i, \hat{\sigma}_j] = 2i\epsilon_{ijk} \hat{\sigma}_k \quad (43)$$

$$\hat{\sigma}_i \hat{\sigma}_j = \delta_{ij} \hat{1}_2 + i\epsilon_{ijk} \hat{\sigma}_k \quad (44)$$

$$\text{tr}(\hat{\sigma}_i) = 0 \quad \text{tr}(\hat{\sigma}_i \hat{\sigma}_j) = 2\delta_{ij} \quad (45)$$

$$\hat{S}_{\pm} = \hat{S}_x \pm i\hat{S}_y = \frac{\hbar}{2} (\hat{\sigma}_x \pm i\hat{\sigma}_y) = \hbar \hat{\sigma}_{\pm} \quad (46)$$

$$\begin{cases} \hat{\sigma}_x = \hat{\sigma}_+ + \hat{\sigma}_- \\ \hat{\sigma}_y = -i(\hat{\sigma}_+ - \hat{\sigma}_-) \end{cases} \quad (47)$$

20.3.1 Ladder / Raising & Lowering / Transition Operator

$$\hat{\sigma}_+ = |0\rangle \langle 1| \quad \& \quad \hat{\sigma}_- = |1\rangle \langle 0| \quad (48)$$

20.3.2 Pauli Matrices ($S_z(\sigma_z)$ -representation)

$$\hat{\sigma}_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \hat{\sigma}_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix} \quad \hat{\sigma}_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (49)$$

21 Arbitrary Angular Momentum

$$\hat{\mathbf{J}} = \hat{\mathbf{L}} + \hat{\mathbf{S}} \quad (50)$$

$$[\hat{J}_i, \hat{J}_j] = i\hbar \epsilon_{ijk} \hat{J}_k \quad (51)$$

OAM & Spin are in DIFFERENT State Vector Spaces:

$$[\hat{L}_i, \hat{S}_j] \equiv 0 \quad (52)$$

$$[\hat{J}_i, \hat{J}_j] = [\hat{L}_i + \hat{S}_i, \hat{L}_j + \hat{S}_j] = [\hat{L}_i, \hat{L}_j] + [\hat{S}_i, \hat{S}_j] = i\hbar \epsilon_{ijk} (\hat{L}_k + \hat{S}_k) = i\hbar \epsilon_{ijk} \hat{J}_k$$

21.1 Ladder Operator

21.1.1 Definition of Ladder Operators

$$\hat{J}_{\pm} = \hat{J}_x \pm i\hat{J}_y \quad (53)$$

$$\left[\hat{J}_i, \hat{\mathbf{J}}^2 \right] = 0 \implies \begin{cases} \hat{\mathbf{J}}^2 |j, m_j\rangle = \hbar j(j+1) |j, m_j\rangle \\ \hat{J}_z |j, m_j\rangle = \hbar m_j |j, m_j\rangle \end{cases} \quad j \geq 0, \quad m_j = \underbrace{-j, -j+1, \dots, j-1, j}_{(2j+1)\text{-fold (degeneracy)}} \quad (54)$$

$$\begin{cases} j: \text{azimuthal quantum number} \\ m_j: \text{magnetic quantum number} \end{cases}$$

21.1.2 Effect of Ladder Operators on Common Eigenstates

$$\hat{J}_{\pm} |j, m_j\rangle = \hbar \sqrt{j(j+1) - m_j(m_j \pm 1)} |j, m_j \pm 1\rangle \quad (55)$$

21.1.3 Restriction

$$\hat{J}_+ |j, j\rangle = \hat{J}_- |j, -j\rangle = 0 \quad (56)$$

21.1.4 Commutations

1. $\left[\hat{J}_i, \hat{J}_j \right] = i\hbar \varepsilon_{ijk} \hat{J}_k$
2. $\left[\hat{\mathbf{J}}^2, \hat{J}_i \right] = 0$
3. $\left[\hat{\mathbf{J}}^2, \hat{J}_{\pm} \right] = 0$
4. $\left[\hat{J}_z, \hat{J}_{\pm} \right] = \pm \hbar \hat{J}_{\pm}$

$$\left[\hat{J}_z, \hat{J}_{\pm} \right] = \left[\hat{J}_z, \hat{J}_x \pm i\hat{J}_y \right] = \left[\hat{J}_z, \hat{J}_x \right] \pm i \left[\hat{J}_z, \hat{J}_y \right] = i\hbar \hat{J}_y \pm i(-i\hbar) \hat{J}_x = \pm \hbar \hat{J}_{\pm}$$

$$5. \hat{J}_+ \hat{J}_- = \hat{\mathbf{J}}^2 - \hat{J}_z^2 + \hbar \hat{J}_z$$

$$\hat{J}_+ \hat{J}_- = (\hat{J}_x + i\hat{J}_y)(\hat{J}_x - i\hat{J}_y) = \hat{J}_x^2 + \hat{J}_y^2 - i \left[\hat{J}_x, \hat{J}_y \right] = \hat{\mathbf{J}}^2 - \hat{J}_z^2 + \hbar \hat{J}_z$$

$$6. (\hat{J}_{\pm})^{\dagger} = \hat{J}_{\mp}$$

\hat{J}_{\pm} are NOT Hermitian Operators, but \hat{J}_+ & \hat{J}_- are mutually Hermitian Conjugates.

21.2 Eigenvalue Equation of Angular Momentum Operator

$$\begin{cases} \hat{\mathbf{J}}^2 |\lambda, m\rangle = \hbar^2 \lambda |\lambda, m\rangle \\ \hat{J}_z |\lambda, m\rangle = \hbar m |\lambda, m\rangle \end{cases} \quad (57)$$

$$[\hat{\mathbf{J}}^2, \hat{J}_\pm] = 0 \implies \hat{J}_\pm \hat{\mathbf{J}}^2 |\lambda, m\rangle = \hat{\mathbf{J}}^2 \hat{J}_\pm |\lambda, m\rangle \implies \lambda \hbar^2 (\hat{J}_\pm |\lambda, m\rangle) = \hat{\mathbf{J}}^2 (\hat{J}_\pm |\lambda, m\rangle) \quad (58)$$

$$\begin{aligned} [\hat{J}_\pm, \hat{J}_z] &= \pm \hbar \hat{J}_\pm \implies \hat{J}_z \hat{J}_\pm = \hat{J}_\pm \hat{J}_z \pm \hbar \hat{J}_\pm \\ \implies \hat{J}_z (\hat{J}_\pm |\lambda, m\rangle) &= (\hat{J}_\pm \hat{J}_z \pm \hbar \hat{J}_\pm) |\lambda, m\rangle = m \hbar \hat{J}_\pm |\lambda, m\rangle \pm \hbar \hat{J}_\pm |\lambda, m\rangle \\ &= (m \pm 1) \hbar (\hat{J}_\pm |\lambda, m\rangle) \end{aligned} \quad (59)$$

$\implies \hat{J}_\pm |\lambda, m\rangle$ is an eigenstate of \hat{J}_z , with eigenvalue of $(m \pm 1)\hbar$.

$$\implies \hat{J}_\pm |\lambda, m\rangle = C_\pm |\lambda, m \pm 1\rangle \quad (60)$$

$$\implies \hat{J}_\pm^n |\lambda, m\rangle = D_\pm |\lambda, m \pm n\rangle \quad (61)$$

$$\begin{cases} \langle \hat{\mathbf{J}}^2 \rangle = \langle \lambda, m | \hat{\mathbf{J}}^2 | \lambda, m \rangle = \lambda \hbar^2 \\ \langle \hat{J}_z^2 \rangle = \langle \lambda, m | \hat{J}_z^2 | \lambda, m \rangle = m^2 \hbar^2 \end{cases} \quad (62)$$

$$\langle \hat{\mathbf{J}}^2 \rangle \geq \langle \hat{J}_z \rangle \implies \lambda \geq m^2 \geq 0 \quad (63)$$

$\exists m_0$ (minimal), $\exists N$ (integer), $m_0 + N$ is the maximal

$$\hat{J}_- |\lambda, m_0\rangle = 0 \quad \hat{J}_+ |\lambda, m_0 + N\rangle = 0 \quad (64)$$

$$0 = \hat{J}_+ \hat{J}_- |\lambda, m_0\rangle = (\hat{\mathbf{J}}^2 - \hat{J}_z^2 + \hbar \hat{J}_z) |\lambda, m_0\rangle = \hbar^2 (\lambda - m_0^2 + m_0) \quad (65)$$

$$0 = \hat{J}_- \hat{J}_+ |\lambda, m_0 + N\rangle = (\hat{\mathbf{J}}^2 - \hat{J}_z^2 - \hbar \hat{J}_z) |\lambda, m_0 + N\rangle = \hbar^2 [\lambda - (m_0 + N)^2 - (m_0 + N)] \quad (66)$$

$$\begin{cases} \lambda - m_0^2 + m_0 = 0 \\ \lambda - (m_0 + N)^2 - (m_0 + N) = 0 \end{cases} \implies \begin{cases} m_0 = -\frac{N}{2} \\ m_0 + N = \frac{N}{2} \\ \lambda = -\frac{N}{2} \left(\frac{N}{2} + 1 \right) \end{cases} \quad (67)$$

let $j = \frac{N}{2}$, $\lambda = j(j+1)$, $m_j = \underbrace{-j, -j+1, \dots, j-1, j}_{2j+1}$

$$\implies \boxed{\begin{cases} \hat{\mathbf{J}}^2 |j, m_j\rangle = j(j+1) \hbar^2 |j, m_j\rangle \\ \hat{J}_z |j, m_j\rangle = m_j \hbar |j, m_j\rangle \end{cases}} \quad (68)$$

$$\hat{J}_\pm |j, m_j\rangle = C_\pm |j, m_j \pm 1\rangle \quad (69)$$

$$\langle j, m_j | j', m_{j'} \rangle = \delta_{jj'} \delta_{m_j m_{j'}} \quad (\hat{J}_\pm)^\dagger = \hat{J}_\mp \quad (70)$$

$$\implies 1 = \langle j, m_j \pm 1 | j, m_j \pm 1 \rangle = \langle j, m_j | \hat{J}_\mp \hat{J}_\pm | j, m_j \rangle = |C_\pm|^2 \langle j, m_j | j, m_j \rangle \quad (71)$$

$$\hat{J}_\mp \hat{J}_\pm = (\hat{J}_x \mp i\hat{J}_y)(\hat{J}_x \pm i\hat{J}_y) = \hat{J}_x^2 + \hat{J}_y^2 \pm i[\hat{J}_x, \hat{J}_y] = \hat{\mathbf{J}}^2 - \hat{J}_z^2 \mp \hbar \hat{J}_z \quad (72)$$

$$\begin{aligned} \implies \langle j, m_j | \hat{J}_\mp \hat{J}_\pm | j, m_j \rangle &= \langle j, m_j | (\hat{\mathbf{J}}^2 - \hat{J}_z^2 \mp \hbar \hat{J}_z) | j, m_j \rangle \\ &= \hbar^2 [j(j+1) - m_j^2 \mp m_j] \langle j, m_j | j, m_j \rangle = |C_\pm|^2 \langle j, m_j | j, m_j \rangle \\ \implies |C_\pm|^2 &= [j(j+1) - m_j(m_j \pm 1)] \hbar^2 \end{aligned} \quad (73)$$

for simplicity, let $C_\pm \in \mathbb{R}$

$$\implies C_\pm = \hbar \sqrt{j(j+1) - m_j(m_j \pm 1)} \quad (74)$$

$$\implies \boxed{\hat{J}_\pm | j, m_j \rangle = \hbar \sqrt{j(j+1) - m_j(m_j \pm 1)} | j, m_j \pm 1 \rangle} \quad (75)$$

22 Commutation Relations between Angular Momentum Operators and Hamiltonian

$$\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + V(r) + \xi(r) \hat{\mathbf{L}} \cdot \hat{\mathbf{S}} \quad (76)$$

$V(r)$: central potential, $\xi(r) \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}$: $\hat{\mathbf{L}}\text{-}\hat{\mathbf{S}}$ coupling \longleftrightarrow fine structure

$$[\hat{L}_i, \hat{p}_j] = i\hbar \varepsilon_{ijk} \hat{p}_k \implies [\hat{L}_i, \hat{\mathbf{p}}^2] = [\hat{L}_i, \hat{p}_j] \hat{p}_j + \hat{p}_j [\hat{L}_i, \hat{p}_j] = i\hbar \varepsilon_{ijk} (\hat{p}_k \hat{p}_j + \hat{p}_j \hat{p}_k) = 0 \quad (77)$$

$$[\hat{S}_i, \hat{\mathbf{p}}^2] = 0 \implies [\hat{J}_i, \hat{\mathbf{p}}^2] = 0 \quad (78)$$

$$[\hat{L}_i, V(r)] = 0 \quad \& \quad [\hat{S}_i, V(r)] = 0 \implies [\hat{J}_i, V(r)] = 0 \quad (79)$$

$$[\hat{L}_i, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] = [\hat{L}_i, \hat{L}_j \hat{S}_j] = i\hbar \varepsilon_{ijk} \hat{L}_k \hat{S}_j \neq 0 \quad (80)$$

$$[\hat{\mathbf{L}}^2, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] = [\hat{L}_i \hat{L}_i, \hat{L}_j \hat{S}_j] = [\hat{L}_i, \hat{L}_j] \hat{L}_i \hat{S}_j + \hat{L}_i [\hat{L}_i, \hat{L}_j] \hat{S}_j = i\hbar \varepsilon_{ijk} (\hat{L}_k \hat{L}_i + \hat{L}_i \hat{L}_k) \hat{S}_j = 0 \quad (81)$$

similarly, $[\hat{S}_i, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] \neq 0$, but $[\hat{\mathbf{S}}^2, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] = 0$. futher, $[\hat{J}_i, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] = 0$, $[\hat{\mathbf{J}}^2, \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}] = 0$.

$$\hat{\mathbf{L}}^2, \hat{\mathbf{S}}^2, \hat{\mathbf{J}}^2, \hat{J}_i \text{ commutes with } \hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + V(r) + \xi(r) \hat{\mathbf{L}} \cdot \hat{\mathbf{S}}$$

23 Addition of Angular Momenta

assume $\hat{\mathbf{J}}_1, \hat{\mathbf{J}}_2, \hat{\mathbf{J}} = \hat{\mathbf{J}}_1 + \hat{\mathbf{J}}_2$, and Commutation Relations:

$$[\hat{J}_{\alpha i}, \hat{J}_{\beta j}] = i\hbar \delta_{\alpha\beta} \varepsilon_{ijk} \hat{J}_{\alpha k} \quad (\alpha, \beta = 1, 2) \quad (82)$$

$$\left[\hat{\mathbf{J}}_\alpha^2, \hat{J}_{\alpha i} \right] \quad (\alpha = 1, 2) \quad (83)$$

Eigenvalue Equations:

$$\begin{cases} \hat{\mathbf{J}}_1^2 |j_1, m_1\rangle = j_1(j_1 + 1)\hbar^2 |j_1, m_1\rangle \\ \hat{J}_{1z} |j_1, m_1\rangle = m_1\hbar |j_1, m_1\rangle \end{cases} \quad (84)$$

$$\begin{cases} \hat{\mathbf{J}}_2^2 |j_2, m_2\rangle = j_2(j_2 + 1)\hbar^2 |j_2, m_2\rangle \\ \hat{J}_{2z} |j_2, m_2\rangle = m_2\hbar |j_2, m_2\rangle \end{cases} \quad (85)$$

$$\left[\hat{J}_i, \hat{J}_j \right] = i\hbar \epsilon_{ijk} \hat{J}_k \quad (86)$$

24 Uncertainty Principle

24.1 Heisenberg Uncertainty Principle

$\forall |\psi\rangle$,

$$\underbrace{\Delta \hat{x}^2 \cdot \Delta \hat{p}^2}_{\text{variance}} \geq \frac{\hbar^2}{4} \quad \text{or} \quad \underbrace{\Delta \hat{x} \cdot \Delta \hat{p}}_{\text{root-mean-square}} \geq \frac{\hbar}{2} \quad (87)$$

24.2 Generalized Uncertainty Principle

$\forall |\psi\rangle$, if $[\hat{A}, \hat{B}] \neq 0$, then

$$\Delta \hat{A}^2 \cdot \Delta \hat{B}^2 \geq \frac{|\langle [\hat{A}, \hat{B}] \rangle|^2}{4} \quad \text{or} \quad \Delta \hat{A} \cdot \Delta \hat{B} \geq \frac{|\langle [\hat{A}, \hat{B}] \rangle|}{2} \quad (88)$$

Variance:

$$\Delta \hat{A}^2 = \langle (\hat{A} - \langle \hat{A} \rangle)^2 \rangle = \langle \hat{A}^2 \rangle - \langle \hat{A} \rangle^2 \quad (89)$$

root-mean-square:

$$\Delta \hat{A} = \sqrt{\Delta \hat{A}^2} = \sqrt{\langle \hat{A}^2 \rangle - \langle \hat{A} \rangle^2} \quad (90)$$

24.2.1 Essential Mathematical Tools

1. If $\hat{F}^\dagger = \hat{F}$ (Hermitian), then eigenvalue of \hat{F} is REAL
2. If $\hat{F}^\dagger = -\hat{F}$ (skew-Hermitian), then \hat{F} is IMAGINARY
3. Cauchy-Schwarz Inequality

$$(a) \quad \left(\sum_n a_n^2 \right) \left(\sum_n b_n^2 \right) \geq \left(\sum_n a_n b_n \right)^2, \text{ equality iff } a_n = k b_n$$

$$(b) \quad |\mathbf{u}|^2 |\mathbf{v}|^2 \geq |\mathbf{u} \cdot \mathbf{v}|^2, \text{ equality iff } \mathbf{u} = k \mathbf{v}$$

$$(c) \quad \boxed{\langle \psi | \psi \rangle \langle \phi | \phi \rangle \geq |\langle \psi | \phi \rangle|^2, \text{ equality iff } |\psi\rangle = \lambda |\phi\rangle}$$

24.2.2 Define Deviation Operator

$$\hat{\sigma}_A = \hat{A} - \langle \hat{A} \rangle \quad \& \quad \hat{\sigma}_B = \hat{B} - \langle \hat{B} \rangle \quad (91)$$

$$\implies \begin{cases} \Delta \hat{A}^2 = \langle \hat{\sigma}_A^2 \rangle \\ \Delta \hat{B}^2 = \langle \hat{\sigma}_B^2 \rangle \end{cases} \quad (92)$$

24.2.3 Properties of $\hat{\sigma}_A, \hat{\sigma}_B$

1. $\hat{\sigma}_A^\dagger = \hat{\sigma}_A, \hat{\sigma}_B^\dagger = \hat{\sigma}_B$ (Hermitian)
2. $\hat{\sigma}_A^\dagger \hat{\sigma}_A = \hat{\sigma}_A^2, \hat{\sigma}_B^\dagger \hat{\sigma}_B = \hat{\sigma}_B^2$
3. $[\hat{\sigma}_A, \hat{\sigma}_B] = [\hat{A}, \hat{B}]$
4. $\hat{\sigma}_A \hat{\sigma}_B = \frac{1}{2}[\hat{\sigma}_A, \hat{\sigma}_B] + \frac{1}{2}\{\hat{\sigma}_A, \hat{\sigma}_B\}$
5. $[\hat{\sigma}_A, \hat{\sigma}_B]^\dagger = -[\hat{\sigma}_A, \hat{\sigma}_B]$ (skew-Hermitian)
6. $\{\hat{\sigma}_A, \hat{\sigma}_B\}^\dagger = \{\hat{\sigma}_A, \hat{\sigma}_B\}$ (Hermitian)
7. $\Delta \hat{A}^2 = \langle \hat{\sigma}_A^2 \rangle, \Delta \hat{B}^2 = \langle \hat{\sigma}_B^2 \rangle$

24.2.4 Proof of GUP

$$\begin{aligned} \text{def: } \hat{\sigma}_A |\psi\rangle &= |\psi_A\rangle \quad \& \quad \hat{\sigma}_B |\psi\rangle = |\psi_B\rangle \\ \langle \hat{\sigma}_A^2 \rangle &= \langle \psi | \hat{\sigma}_A \hat{\sigma}_A | \psi \rangle = \langle \psi | \hat{\sigma}_A^\dagger \hat{\sigma}_A | \psi \rangle = \langle \psi_A | \psi_A \rangle \quad \& \quad \langle \hat{\sigma}_B^2 \rangle = \langle \psi_B | \psi_B \rangle \\ \implies \langle \hat{\sigma}_A^2 \rangle \langle \hat{\sigma}_B^2 \rangle &= \langle \psi_A | \psi_A \rangle \langle \psi_B | \psi_B \rangle \geq |\langle \psi_A | \psi_B \rangle|^2 = |\langle \psi | \hat{\sigma}_A \hat{\sigma}_B | \psi \rangle|^2 = |\langle \hat{\sigma}_A \hat{\sigma}_B \rangle|^2 \\ \langle \hat{\sigma}_A \hat{\sigma}_B \rangle &= \frac{1}{2} \langle [\hat{\sigma}_A, \hat{\sigma}_B] \rangle + \frac{1}{2} \langle \{\hat{\sigma}_A, \hat{\sigma}_B\} \rangle \\ \implies |\langle \hat{\sigma}_A \hat{\sigma}_B \rangle|^2 &\geq \frac{1}{4} |\langle [\hat{\sigma}_A, \hat{\sigma}_B] \rangle|^2 = \frac{1}{4} |\langle [\hat{A}, \hat{B}] \rangle|^2 \implies \text{GUP} \end{aligned}$$

24.3 Energy-Time Uncertainty Relation

$$\Delta E \cdot \Delta T \geq \frac{\hbar}{2} \quad (93)$$

24.3.1 Ehrenfest Theorem

$$\begin{aligned} [\hat{A}, \hat{H}] &\neq 0, \quad \frac{\partial \hat{A}}{\partial t} = 0, \\ \frac{d}{dt} \langle \hat{A} \rangle &= \frac{1}{i\hbar} \langle [\hat{A}, \hat{H}] \rangle \end{aligned} \quad (94)$$

24.3.2 Proof

$$\implies \Delta \hat{A} \cdot \Delta \hat{H} \geq \frac{1}{2} \left| \langle [\hat{A}, \hat{H}] \rangle \right| = \frac{\hbar}{2} \left| \frac{d}{dt} \langle \hat{A} \rangle \right| \implies \Delta \hat{H} \frac{\Delta \hat{A}}{\left| \frac{d}{dt} \langle \hat{A} \rangle \right|} = \frac{\hbar}{2}$$

$$\text{define } \Delta \hat{H} = \Delta E, \Delta T = \frac{\Delta \hat{A}}{\left| \frac{d}{dt} \langle \hat{A} \rangle \right|}$$

$$\implies \Delta E \cdot \Delta T \geq \frac{\hbar}{2}$$

24.4 Minimum Uncertainty State

$\frac{\hbar}{2}$: quantum limit (Heisenberg limit) \implies Minimum Uncertainty State

$$\exists |\psi\rangle_{\min}, \Delta \hat{A} \cdot \Delta \hat{B} = \frac{1}{2} \left| \langle [\hat{A}, \hat{B}] \rangle \right|$$

$|\psi\rangle_{\min}$: Minimum Uncertainty State

24.4.1 MUS of \hat{x} & \hat{p}

$$\begin{cases} \hat{\sigma}_x = \hat{x} - \langle \hat{x} \rangle \\ \hat{\sigma}_p = \hat{p} - \langle \hat{p} \rangle \end{cases}, \text{ let } |\phi\rangle = (\hat{\sigma}_x - i\lambda \hat{\sigma}_p) |\psi\rangle \text{ (testing state)}$$

$$\begin{aligned} \langle \phi | \phi \rangle &= \langle \psi | (\hat{\sigma}_x + i\lambda \hat{\sigma}_p) (\hat{\sigma}_x - i\lambda \hat{\sigma}_p) |\psi\rangle = \langle \psi | (\hat{\sigma}_x^2 - i\lambda \hat{\sigma}_x \hat{\sigma}_p + i\lambda \hat{\sigma}_p \hat{\sigma}_x + \lambda^2 \hat{\sigma}_p^2) |\psi\rangle \\ &= \langle \psi | \hat{\sigma}_x^2 |\psi\rangle - i\lambda \langle \psi | [\hat{\sigma}_x, \hat{\sigma}_p] |\psi\rangle + \lambda^2 \langle \psi | \hat{\sigma}_p^2 |\psi\rangle = \langle \hat{\sigma}_x^2 \rangle + \lambda \hbar + \lambda^2 \langle \hat{\sigma}_p^2 \rangle \\ |\psi\rangle = 0 &\implies \langle \phi | \phi \rangle = 0 \implies \langle \hat{\sigma}_p^2 \rangle \lambda^2 + \hbar \lambda + \langle \hat{\sigma}_x^2 \rangle = 0 \end{aligned} \tag{95}$$

$$\Delta = \hbar^2 - 4 \langle \hat{\sigma}_x^2 \rangle \langle \hat{\sigma}_p^2 \rangle \lambda^2 \leq 0 \implies \langle \hat{\sigma}_x^2 \rangle \langle \hat{\sigma}_p^2 \rangle \lambda^2 \geq \frac{\hbar^2}{4}$$

$$\implies \Delta \hat{x}^2 \cdot \Delta \hat{p}^2 \geq \frac{\hbar^2}{4} \quad \text{or} \quad \Delta \hat{x} \cdot \Delta \hat{p} \geq \frac{\hbar}{2}$$

$$\Delta = 0 \implies \begin{cases} \lambda = -\frac{2}{\hbar} \Delta \hat{x}^2 \\ |\phi\rangle = \left(\hat{\sigma}_x + \frac{2i}{\hbar} \Delta \hat{x}^2 \hat{\sigma}_p \right) |\psi\rangle = 0 \end{cases} \tag{96}$$

$$\langle x | (\hat{x} - \langle \hat{x} \rangle) - i\lambda \left(-i\hbar \frac{d}{dx} - \langle \hat{p} \rangle \right) |\psi\rangle = (x - \langle \hat{x} \rangle) \psi(x) - \lambda \hbar \frac{d}{dx} \psi(x) + i\lambda \langle \hat{p} \rangle \psi(x)$$

25 Quantum Dynamics

25.1 Schrödinger Equation

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = \hat{H} |\psi(t)\rangle \quad (97)$$

where $\hat{H} = \frac{\hat{\mathbf{p}}^2}{2m} + V(\hat{\mathbf{x}})$

25.1.1 Remark

1. Pure State
2. Closed System $\implies \langle \psi(t) | \psi(t) \rangle = 1$
3. Non-Relativistic QM
4. Initial Value Problem (IVP): $|\psi(t = t_0)\rangle = |\psi(t_0)\rangle$
5. Schrödinger Picture: $\frac{d\hat{A}}{dt} \equiv 0$

25.1.2 Representation of Position

In \mathbf{x} -representation: $\hat{\mathbf{p}} = -i\hbar \nabla$, $\langle \mathbf{x} | \psi(t) \rangle = \psi(\mathbf{x})$

$$i\hbar \frac{d}{dt} |\psi(t)\rangle = \hat{H} |\psi(t)\rangle \implies i\hbar \frac{\partial}{\partial t} \psi(\mathbf{x}, t) = \left(-\frac{\hat{\mathbf{p}}^2}{2m} + V(\hat{\mathbf{x}}) \right) \psi(\mathbf{x}, t) \quad (98)$$

1. In general, $\frac{\partial \hat{A}}{\partial t} \equiv 0$ & $\frac{\partial \hat{H}}{\partial t} \equiv 0$. Unless there is external influence.
2. $\langle \psi(t) | \psi(t) \rangle = 1 \iff \int d^3x |\psi(\mathbf{x}, t)|^2 = 1$
3. $\psi(\mathbf{x}, t = t_0) = \psi(\mathbf{x}, t_0)$

25.2 Time Evolution Operator

$$|\psi(t)\rangle = \hat{U}(t, t_0) |\psi(t_0)\rangle \quad (99)$$

$$\text{Schrödinger eq.} \implies i\hbar \frac{d}{dt} [\hat{U}(t, t_0) |\psi(t_0)\rangle] = \hat{H} [\hat{U}(t, t_0) |\psi(t_0)\rangle] \quad (100)$$

$$\implies i\hbar \frac{d}{dt} \hat{U}(t, t_0) = \hat{H} \hat{U}(t, t_0) \quad (101)$$

25.2.1 TEO for Conservative System

If the system is conservative, that is $\frac{\partial \hat{H}}{\partial t} \equiv 0$.

$$\implies \frac{d\hat{U}}{\hat{U}} = -\frac{i}{\hbar} \hat{H} dt \implies \boxed{\hat{U}(t, t_0) = e^{-i\hat{H}(t-t_0)/\hbar}}$$

If $\frac{\partial \hat{H}}{\partial t} \neq 0$, then

$$\begin{aligned} \hat{U}(t, t_0) &= \hat{1} - \frac{i}{\hbar} \int dt' \hat{U}(t', t_0) \\ &= \hat{1} - \frac{i}{\hbar} (\text{Dyson Series}) \end{aligned} \tag{102}$$

25.2.2 Properties of $\hat{U}(t, t_0)$

1. $\hat{U}^\dagger(t, t_0) = \hat{U}^{-1}(t, t_0 = \hat{U}(t_0, t))$ or $\hat{U}^\dagger(t, t_0)\hat{U}(t, t_0) = \hat{1}$
2. $\hat{U}(t, t_1)\hat{U}(t_1, t_0) = \hat{U}(t, t_0) \xrightarrow{\text{expand}} \hat{U}(t, t_n)\hat{U}(t_n, t_{n-1}) \cdots \hat{U}(t_2, t_1)\hat{U}(t_1, t_0) = \hat{U}(t, t_0)$
3. $\hat{U}(t, t_0) = \hat{U}(t - t_0) = \hat{U}(\tau)$ ($\tau \stackrel{\text{def}}{=} t - t_0$)

26 title