

A resolved view of star formation, gas cycling and chemical enrichment in Virgo Cluster spirals.

Rongjun Huang

Supervisors: Luca Cortese & Luke Davies

ICRAR/UWA

Feb 26, 2026



International
Centre for
Radio
Astronomy
Research




Curtin University



THE UNIVERSITY OF
WESTERN
AUSTRALIA

Galaxy evolution is regulated by the baryon cycle

The Baryon Cycle




- Gas inflow → fuels star formation
- Star formation → builds stars + metal synthesis
- Feedback → drives outflows + recycling

Bathtub/Gas-regulator Model

Star formation Rate (SFR) = current growth rate (“how fast building now”)

Stellar Mass (M_*) = integrated past star formation (“how much built up”)



Gas-phase metallicity (Z_{gas}) = enrichment vs dilution (“how processed the gas is”)




How do we measure them?

- **SFR**. Estimated by $H\alpha$: recent Star Formation, timescale $\sim 10\text{Myr}$
- **M_*** . Estimated by stellar continuum: Spectral Energy Distribution (SED) fitting, Mass-to-Light ratio (M/L)
- **Z_{gas}** . Commonly referred to the oxygen abundance, $12+\log(\text{O}/\text{H})$: measured by emission lines or physical models

Global SFMS (SFR vs M_*)

Gas-star formation cycle


- SFR tracks gas accretion and stellar mass build up
- $\Rightarrow$ **Star-Forming Main Sequence (SFMS):** $\text{SFR} \propto M_*$ with small scatter and evolve with redshift



Global MZR (Z_{gas} vs M_*)

Metal enrichment


- Dilution by inflow; metal removal via outflows.
- Deeper potentials (higher M_*) $\Rightarrow$ better metal retention $\Rightarrow$ **Mass-Metallicity Relation (MZR)**: Z_{gas} increases with M_* (steep at low mass, saturating at high mass, and evolve with cosmic time).



Global FMR (interplay between M_* , SFR and Z_{gas})


Fundamental Metallicity Relation (FMR)

- A 3D surface: $Z_{\text{gas}}(M_*, \text{SFR})$
- “Fundamental”: smaller scatters than SFMS and MZR; roughly invariant with redshift to $z \sim 2-3$
- Interpretation: Adding SFR as a second dependence of MZR.




Global (Integrated) $\Rightarrow$ Local (Resolved)

Integrated galaxy-wide




Spatially resolved scales




Require Integral Field Unit (IFU) data

Integral Field Unit (IFU) & Datacube



Resolved scaling relations

- **rSFMS (bottom left)**: Σ_{SFR} correlates with Σ_*
- **rMZR (bottom right)**: Z_{gas} rises with Σ_* ; steep at low-mass end, saturation at high-mass
- **rFMR (top)**: A 3D surface of $Z_{\text{gas}}(\Sigma_*, \Sigma_{\text{SFR}})$, but **debated!**
 - Some IFU surveys found no clear signal of secondary dependence on Σ_{SFR} (different Z_{gas} diagnostics and sample selection)
 - Still an anti-correlation between $12 + \log(\text{O}/\text{H})$ and Σ_{SFR} at fixed Σ_* ?





International
Centre for
Radio
Astronomy
Research

My PhD goals

Answer the following questions at ~ 100 pc scales:

- Spatially resolved properties: Σ_* , Σ_{SFR} , Z_{gas} , etc.
- Do resolved scaling relations exist, and are they universal?
- Origin: are these relations driven by local self-regulation or global/environmental control

Understanding this requires resolved spectroscopy across an environmentally processed spiral population



International
Centre for
Radio
Astronomy
Research

Observational gap

Global relations: well-established in past two decades

Resolved relations (IFU, past decade):

- kpc scales ($\gtrsim$ kpc): MaNGA, SAMI, CALIFA, MUSE-Wide, GASP, MAGPI, etc.
- kpc scales (~ 100 pc)
 - { Field galaxies: PHANGS, MAD.
 - { Cluster galaxies: ??? (this work)



International
Centre for
Radio
Astronomy
Research

This Work. MAUVE Survey (MUSE + ALMA)

- 40 late-type Virgo galaxies; VLT/MUSE provides ~ 100 pc datacubes; nGIST products: Voronoi binning, SFH, emission line...
- Complementary ALMA (molecular gas, down to ~ 60 pc; also legacy from VERTICO survey).
- Targets span several infall phases (first infall $\Rightarrow$ pericentre $\Rightarrow$ post-pericentre), ideal for environmental sequence tests.
- Survey current status:
 - MAUVE-MUSE: 26/40 fully delivered (14 used in my first project).
 - MAUVE-ALMA: 34/40 fully delivered



MAUVE


Multiphase Astrophysics to Unveil the Virgo Environment




Project 1. rMZR's secondary dependence on Σ_{SFR}



1. An anti-correlation between Z_{gas} and Σ_{SFR} at fixed Σ_* ?
2. What causes this secondary dependence?



MaNGA, $\gtrsim$ kpc (Sánchez-Menguiano+2019)





MAGPI, $\gtrsim$ kpc (Koller+2024)



International
Centre for
Radio
Astronomy
Research


Pipeline (`Mass.py`): Stellar Mass Map






International
Centre for
Radio
Astronomy
Research

Pipeline (`SFR+Z.py`): SFR & [O/H] Maps




14 MAUVE galaxies: Σ_{SFR} and [O/H] (O3N2-M13) maps (SF Spaxels)



rMZR and Σ_{SFR} -- Z_{gas} relation

- Left: rMZR, color-coding by Σ_{SFR}
- Right: Σ_{SFR} -- Z_{gas} relation, color-coding by Σ_*
 - Low- Σ_* end: anti-correlated;
 - High- Σ_* end: correlated.





International
Centre for
Radio
Astronomy
Research

$\Delta\Sigma_{\text{SFR}}$, ΔZ_{gas} , Local Moran correlation and Σ_* maps

Top panels.

In each Σ_* bin (bottom right), for each spaxel i , we compute:

$$\begin{cases} \Delta \log_{10} \Sigma_{\text{SFR}}(i) \equiv \log_{10} \Sigma_{\text{SFR}}(i) - \langle \log_{10} \Sigma_{\text{SFR}} \rangle_m, \\ \Delta [\text{O}/\text{H}](i) \equiv [\text{O}/\text{H}](i) - \langle [\text{O}/\text{H}] \rangle_m, \end{cases}$$


Bottom left.

Local Moran-like correlation Map

Red: correlation

Blue: anti-correlation


Both correlated and anti-correlated Hii regions are presented everywhere



Mass-dependent reversal

Spearman coefficient ρ
between $\Delta[\text{O}/\text{H}]$ and
 $\Delta\log\Sigma_{\text{SFR}}$, as a function
of $\log\Sigma_*$

Low- Σ_* : anti-correlated
High- Σ_* : correlated
Inversion point:
 $\log(\Sigma_*/M_\odot\text{kpc}^{-2}) \sim 7.5$
– 8.0



What drives this $\Delta\Sigma_{\text{SFR}}--\Delta Z_{\text{gas}}$ (anti-)correlation?




Molecular gas mass and gas-phase metal mass continuity equations:

$$\frac{d\Sigma_g(t)}{dt} = \Sigma_\Phi(t) - (1 - R)\Sigma_{\text{SFR}}(t) - \Sigma_{\text{out}}(t), \quad (1)$$

$$\frac{d\Sigma_Z(t)}{dt} = y(1 - R)\Sigma_{\text{SFR}}(t) - [(1 - R)\Sigma_{\text{SFR}}(t) + \Sigma_{\text{out}}(t)]Z(t) + \Sigma_\Phi(t)Z_\Phi, \quad (2)$$

Spatially resolved gas-regulator model




Changes of SFR vs Changes of GSR

$$\delta Z(t) \propto \delta \log_{10}(\Sigma_{\text{SFR}}(t)) - \delta \log_{10}(\Sigma_{\Phi}(t))$$

Star-formation-driven
variability: change of
 Σ_{SFR} in log scales

Gas-supply-driven
variability: change of Σ_{Φ}
in log scales

Star-formation-driven variability dominates $\Rightarrow \uparrow \Sigma_{\text{SFR}}$
(metal yields) with smaller changes in $\Sigma_{\Phi} \Rightarrow \uparrow Z_{\text{gas}}$.
Gas-supply-driven variability dominates $\Rightarrow \uparrow \Sigma_{\text{SFR}}$ with
with larger changes in Σ_{Φ} (Z_{Φ} dilution) $\Rightarrow \downarrow Z_{\text{gas}}$.






Summary of Project 1

1. **Derive** my pipelines and analyse 14 MAUVE—MUSE galaxies
2. **Confirm** the rMZR's secondary dependence on Σ_{SFR} at ~ 100 pc scales
3. **Detect** a mass-dependent inversion in residual $\Delta Z_{\text{gas}} - \Delta \Sigma_{\text{SFR}}$ coupling
4. **Show** that this flip-sign is spatially varying in the local Moran-like map
5. **Set up** a spatially resolved gas-regulator model and **interpret** the flip-sign as a transition from gas supply/dilution-dominated to star formation/enrichment-dominated regulation

Project 2: 26 MAUVE–MUSE galaxies

1. Double our sample sizes: $14 \Rightarrow 26$ MAUVE–MUSE galaxies
2. Beyond Project 1: T_e -based direct measurement of metallicity, ionization conditions, electron density, diffused ionised gas (DIG) contamination, more selection criteria, etc.






International
Centre for
Radio
Astronomy
Research

Project 3: Adding MAUVE—ALMA

Closing the baryon-cycle loop with molecular gas (fuel+regulator)

Molecular gas maps provide the direct handle on fuel availability and cycling

Measurement of Σ_{gas} (and potentially GSR, Σ_{ϕ})



Rongjun Huang (黄镭钧)¹, Luca Cortese¹, Barbara Catinella¹, Luke J. M. Davies¹,
Toby Brown², Andrei Ristea^{3,4}, Alessandro Boselli⁵, Andrew J. Battisti^{1,6}, Vicente Villanueva^{7,8},
Kristine Spekkens⁹, Sara L. Ellison¹⁰, Daniel A. Dale¹¹, Sabine Thater¹², Amirnezhad Amiri^{13,14}

[illegible]



International
Centre for
Radio
Astronomy
Research

End

Thank you !
Questions?

