


20260317 Make specMask_GAS_narrow10

The default spectral mask for GAS module is specMask_GAS:

```
# This file defines spectral ranges that are masked during the
analysis of the
# SFH module. Note that all masks are defined in restframe
wavelength ranges.
# If you intend to mask sky-lines, please use the keyword
'sky' in the comment
# column.
#
# Lambda [Angst.]    Width [Angst.]    Comment
5577.00              30              sky
5890.00              30              sky
5896.00              30              sky
5890.00              30              NaD
5896.00              30              NaD
6300.00              30              sky
6363.00              30              sky
```

So clearly we need to adding the skyline masks for $> 6800\text{\AA}$. Also, as mentioned before, the skyline masks at $6300\text{\AA}$ and $6300\text{\AA}$ may be too wide. For $[\text{S III}]\lambda 6312\text{\AA}$, it is coincidentally shifted to the edge of $6363\text{\AA}$ skyline mask. So in `ppxf`, it seems to fit some "illusion". For example, below is a crazy bright $[\text{S III}]$ spaxel, but the observed spectrum is actually flat.



Below is how i create the mask:

1. Change sky masks at 6300 and 6363 from width 30 to width 10.
2. Copy all sky masks with `Lambda > 6800` from `specMask_KIN_narrow10`.
3. Keep target fitting regions unmasked for lines:
 - ArIII 7135.79
 - OII 7318.92, 7319.99, 7329.67, 7330.73
 - HPa11 8862.89
 - HPa10 9015.30
 - SIII 9068.60
4. Fitting regions use width 20 Angstrom and redshift range z in $[-0.001, 0.01]$.
5. Any new sky mask that overlaps these fitting regions (even partially) is dropped.

