
A
Died by day 730 (3,459)
Survival at 730 not confirmed (14,361)
Alive at 730, no later record (1,476)
CV hospitalization, days 1–730 (1,048)
CV hospitalization before index (1,281)
No follow-up after day 730 (358)
Surgically correctable secondary cause (808)
Death within 90 days (1,510)
Coverage ended before day 730 (11,445)
Association analysis
Adults with hypertension and a later BP reading n = 50,946
Eligible at index n = 48,628
At-index cohort (coverage through day 730), n = 37,183
Landmark population (primary), n = 15,200 664 CV hospitalizations
survival confirmed by day 730 only
Day-730-information sample, n = 7,649 318 CV hospitalizations
No systolic reading >140 mmHg (16,527)
Prediction analysis
First systolic reading >140 mmHg, n = 32,101
Any data after the prediction date, n = 31,390 (competing-event sample)
Prediction sample n = 30,942; 10,899 with no recorded BP in 90 days
No data after the prediction date (711)
Data ended within 90 days (254)
Died within 90 days (194)
B
Index
Landmark (day 730)
Covariates
Trajectory classes
Gap in recorded contact
Outcome follow-up
Prediction window (90 days)
−365
0
365
730
1095
1460
Days from index blood pressure reading