

1 ORDINARY DIFFERENTIAL EQUATIONS (ODE)

8th Aug 2024 12:47pm

Differential equation (DE)

An equation involving derivative(s) of dependent variable with respect to independent variable(s) is called differential equation.

e.g: $x \frac{dy}{dx} + y = 0$

- **Order of DE:** Order of highest order derivative of dependent variable, with respect to independent variable involved in the equation.
- **Degree of DE:** If differential equation is a polynomial equation of derivative, the highest power (positive integral index) of highest order derivative is its degree.

ODE

A differential equation involving derivatives of dependent variable with respect to only one independent variable is called ODE

e.g:

$$3 \left(\frac{d^2 y}{dx^2} \right)^4 + \left(\frac{dy}{dx} \right)^5 = 0$$

This has degree 4 and order 2

1st order differential equation

General form : $f(x, y, y') = 0$

or $y' = g(x, y)$

e.g: $y^2 = \sqrt{x^2 + y^2}$

1st Order Linear Differential equation

General form:

$$y' + P(x)y = Q(x) \quad \text{where functions } P \& Q \text{ are continuous in some interval } I$$

General solution:

$$yF = \int FQdx + C$$

$$\text{where } F = e^{\int Pdx}$$

F is also known as Integration factor

1 solve $y' + 2xy = x$

Answer

Given equation is an 1st order LDE comparing the general form: $y' + P(x)y = Q(x)$ we get: $P = 2x, Q = x$

\therefore Integrating factor $F = e^{\int P \cdot dx} = e^{\int 2x \cdot dx} = e^{x^2}$

$$\begin{aligned}\therefore \text{Solution: } y \cdot F &= \int FQdx + C \\ ye^{x^2} &= \int e^{x^2} x dx + C\end{aligned}$$

let $t = x^2$

$$\therefore 2x = \frac{dt}{dx} \quad \text{or} \quad xdx = \frac{1}{2}dt$$

$$\begin{aligned}\therefore \int e^{x^2} x dx &= \int \frac{1}{2}e^t dt \\ &= \frac{1}{2}e^t = \frac{1}{2}e^{x^2}\end{aligned}$$

$$\begin{aligned}\therefore ye^{x^2} &= \frac{1}{2}e^{x^2} + C \\ \therefore y &= e^{-x^2} \left(\frac{1}{2}e^{x^2} + C \right)\end{aligned}$$

$$\boxed{y = \frac{1}{2} + e^{-x^2}C}$$

2 Solve $\frac{dy}{dx} + 2y \tan(x) = \sin(x)$

Answer

Here $P(x) = 2 \tan(x), Q(x) = \sin(x)$

$$\begin{aligned}\text{Integrating factor (IF) } F &= e^{\int P \cdot dx} \\ &= e^{\int 2 \tan(x) dx} \\ &= e^{2 \cdot \ln |\sec(x)|} \\ &= (e^{\ln |\sec(x)|})^2 \\ &= |\sec(x)|^2 \equiv \sec^2(x)\end{aligned}$$

\therefore Solution:

$$\begin{aligned}y \cdot \sec^2(x) &= \int \sin(x) \cdot \sec^2(x) dx + C \\ &= \int \sec(x) \tan(x) dx + C \\ &= \sec(x) + C \\ \therefore y &= \frac{\sec(x)}{\sec^2(x)} + \frac{C}{\sec^2(x)} \\ y &= \cos(x) + \cos^2(x) \cdot C\end{aligned}$$

3 Find solution of initial value problem:

$$x^2 y' - xy = x^4 \cos(2x), \quad y(\pi) = 2\pi$$

Answer

$$x^2 y' - xy = x^4 \cos(2x) \\ \therefore y' - x^{-1}y = x^2 \cos(2x)$$

$$P(x) = -x^{-1}, Q(x) = x^2 \cos(2x) \\ \therefore \text{I.F} = e^{\int p(x) \cdot dx} = e^{-\int x^{-1} dx} \\ = e^{-\ln(x)} = x^{-1} = \frac{1}{x}$$

General Solution:

$$y \cdot IF = \int Q \cdot IF \cdot dx + C \\ y \cdot \frac{1}{x} = \int x^2 \frac{1}{x} \cos(2x) \cdot dx + C \\ = \int x \cos(2x) dx + C \\ = \frac{x \sin(2x)}{2} - \frac{1}{2} \int \sin(2x) dx + C \\ = \frac{x \sin(2x)}{2} + \frac{\cos(2x)}{4} + C \\ \therefore y = \frac{x^2 \cdot \sin(2x)}{2} + \frac{x \cos(2x)}{4} + xC \quad (1)$$

It is given that $y(\pi) = 2\pi$

$$\therefore 2\pi = \frac{\pi^2 \cdot \overbrace{\sin(2\pi)}^0}{2} + \frac{\pi \cdot \cos(2\pi)}{4} + \pi \cdot C \\ \therefore 2 = \frac{1}{4} + c \\ \Rightarrow c = 2 - \frac{1}{4} = \frac{7}{4}$$

Substitute in Eq.(1) to get specific solution:

$$\boxed{y = \frac{x^2 \sin(2x)}{2} + \frac{x \cos(2x)}{4} + \frac{7}{4}x}$$

4 Solve $y' - 2xy = 2$

Answer

$$\text{Let } P(x) = -2x, Q(x) = 2x$$

$$I.F = e^{\int -2x \cdot dx} = e^{-x^2}$$

$$\therefore \text{Solution: } y \cdot e^{-x^2} = \int 2x \cdot e^{-x^2} dx + c$$

$$\text{Let } -x^2 = t \Rightarrow dx = \frac{dt}{-2x}$$

$$\therefore \int 2xe^{-x^2} dx + C = - \int e^t dt = -e^t + C \\ = -e^{-x^2} + C$$

$$\therefore y \cdot e^{-x^2} = -e^{-x^2} \cdot C$$

$$\boxed{\therefore y = -1 + e^{x^2} \cdot C}$$

5 Solve $xy' - 2y = -x$

Answer

$$\begin{aligned}xy' - 2y &= -x \\ \therefore y' - \frac{2}{x} \cdot y &= -1 \\ \therefore P(x) &= -\frac{2}{x}, Q(x) = -1 \\ IF &= e^{\int P x} = e^{\int -2/x \cdot dx} \\ &= e^{-2 \ln(x)} = x^{-2}\end{aligned}$$

\therefore Solution:

$$\begin{aligned}y \cdot x^{-2} &= \int x^{-2} \times (-1) \cdot dx + C \\ &= \frac{x^{-1}}{-1} \times -1 + C \\ &= \frac{1}{x} + C \\ \therefore \boxed{y} &= \boxed{x + Cx^2}\end{aligned}$$

6 Solve $xy' + 2y = \frac{\cos(x)}{x}$

Answer

$$\begin{aligned}xy' + 2y &= \frac{\cos(x)}{x} \\ y' + \frac{2}{x} \cdot y &= \frac{\cos(x)}{x^2} \\ P(x) &= \frac{2}{x}, Q(x) = \frac{\cos(x)}{x^2} \\ IF &= e^{\int P \cdot dx} = e^{\int 2/x \cdot dx} \\ &= e^{\ln(x^2)} = x^2\end{aligned}$$

Solution: $y \cdot IF = \int IF \cdot Q \cdot dx + C$

$$\begin{aligned}y \cdot x^2 &= \int x^2 \cdot \frac{\cos(x)}{x^2} \cdot dx + C \\ &= \int \cos(x) dx + C \\ y \cdot x^2 &= \sin(x) + C \\ \boxed{y} &= \boxed{\frac{\sin(x)}{x^2} + \frac{C}{x^2}}\end{aligned}$$

7 Solve $y' + \frac{2y}{x} = \frac{4}{x}$. where $y(1) = 6$

Answer

$$\begin{aligned}P(x) &= \frac{2}{x}, Q(x) = \frac{4}{x} \\ IF &= e^{\int P dx} = e^{\int 2/x \cdot dx} = x^2\end{aligned}$$

\therefore Solution:

$$\begin{aligned}y \cdot x^2 &= \int x^2 \cdot \frac{4}{x} dx + C = 2x^2 + C \\ \therefore y &= 2 + \frac{C}{x^2}\end{aligned}$$

Given $y(1) = 6$

$$\therefore 6 = 2 + \frac{C}{1} \Rightarrow C = 4$$

$$y = 2 + \frac{4}{x^2}$$

Variable Separable Equation

A differential equation of the form $m(x, y)dx + n(x, y)dy = 0$ is a variable separable equation if it can be expressed in the form: $f(x)dx + g(y)dy = 0$

1 Solve : $\frac{dy}{dx} = \frac{y}{x}$

Answer

$$\begin{aligned}\frac{dy}{dx} &= \frac{y}{x} \\ \Rightarrow dy \cdot x &= dx \cdot y \\ \Rightarrow \frac{dx}{x} &= \frac{dy}{y}\end{aligned}$$

Integrate both sides:

$$\begin{aligned}\int \frac{dx}{x} &= \int \frac{dy}{y} \\ \ln(y) &= \ln(x) + \ln(C) \\ &= \ln(xC) \\ \Rightarrow \boxed{y} &= \boxed{x C}\end{aligned}$$

2 Solve: $(y + 2)dx + y(x + 4)dy = 0$

Answer

Divide by $(y + 2)(x + 4)$

$$\frac{1}{x + 4}dx + \frac{y}{y + 2}dy = 0$$

Integrating both sides:

$$\begin{aligned}\int \frac{dx}{x + 4} + \int \frac{y}{y + 2}dy &= \ln(C) & [\ln(C) \text{ is used to make further steps easier}] \\ \int \frac{dx}{x + 4} + \int \frac{y + 2 - 2}{y + 2}dy &= \ln(C) \\ \int \frac{dx}{x + 4} + \int \left[\frac{y + 2}{y + 2} + \frac{-2}{y + 2} \right] dy &= \ln(C) \\ \int \frac{dx}{x + 4} + \int \left[1 + \frac{-2}{y + 2} \right] dy &= \ln(C)\end{aligned}$$

$$\ln(x + 4) + y - 2\ln(y + 2) = \ln(C)$$

$$\begin{aligned}y &= \ln(C) + 2\ln(y + 2) - \ln(x + 4) \\ \therefore \boxed{y} &= \boxed{\ln \left[C \cdot \frac{(y + 2)^2}{x + 4} \right]}\end{aligned}$$

3 solve $3x \sin(y) \cdot dx + (x^2 + 1) \cdot \cos(y) \cdot dy = 0$

Answer

Divide by $\sin(y) \cdot (x^2 + 1)$

$$\therefore \frac{x}{x^2 + 1} dx + \frac{\cos(y)}{\sin(y)} dy = 0$$

Integrate both sides:

$$\int \frac{x}{x^2 + 1} dx + \int \frac{\cos(y)}{\sin(y)} dy = \ln(C)$$

$$\text{Let } t = x^2 + 1 \Rightarrow dx = \frac{dt}{2x}$$

$$u = \sin(y) \Rightarrow du = dy \cos(y)$$

$$\therefore \int \frac{dt}{2t} + \int \frac{du}{u} = \ln(C)$$

$$\frac{1}{2} \ln(t) + \ln(u) = \ln(C)$$

$$\frac{1}{2} \ln(x^2 + 1) + \ln(\sin(y)) = \ln(C)$$

$$\ln(\sin(y)) = \ln \left[\frac{C}{(x^2 + 1)^2} \right]$$

$$\sin(y) = \frac{C}{(x^2 + 1)^2}$$

or

$$y = \sin^{-1} \left[\frac{C}{(x^2 + 1)^2} \right]$$

4 Solve $\tan(\theta)dr + 2r \cdot d\theta = 0$

Answer

$$\frac{dr}{2r} + \frac{d\theta}{\tan(\theta)} = 0$$

$$\int \frac{dr}{2r} + \underbrace{\int \frac{d\theta}{\tan(\theta)}}_{\cot(\theta)d\theta} = \ln(C)$$

$$\frac{1}{2} \ln(r) + \ln(\sin(\theta)) = \ln(C)$$

$$\ln(\sqrt{r}) + \ln(\sin(\theta)) = \ln(C)$$

$$\ln(\sqrt{r}) = \ln \left[\frac{C}{\sin(\theta)} \right]$$

$$\sqrt{r} = \frac{C}{\sin(\theta)}$$

$$\Rightarrow r = \frac{C}{\sin^2(\theta)}$$

$$r = C \cdot \csc^2(\theta)$$

5 Solve $4xy \, dx + (x^2 + 1) \, dy = 0$

Answer

$$4xy \, dx + (x^2 + 1) \, dy = 0$$

$$\Rightarrow \frac{4x}{x^2 + 1} dx + \frac{dy}{y} = 0$$

$$2 \int \frac{2x}{x^2 + 1} dx + \int \frac{dy}{y} = \ln(C)$$

$$2 \ln(x^2 + 1) + \ln(y) = \ln(C)$$

$$y = \frac{C}{(x^2 + 1)^2}$$

Homogeneous Differential Equation

An differential equation that can be reduced into the form: $\frac{dy}{dx} = F\left(\frac{y}{x}\right)$ is called homogerious differential equation. This can be solved by putting $y = vx$ and hence reducing to variable separable form.

1 Solve $2xy \cdot \frac{dy}{dx} - y^2 + x^2 = 0$

Answer

$$2xy \frac{dy}{dx} - y^2 + x^2 = 0$$

$$\Rightarrow \frac{dy}{dx} = \frac{y^2 - x^2}{2xy}$$

Put $y = vx \therefore v = \frac{y}{x}$
 \therefore Equation becomes:

$$v + x \cdot \frac{dv}{dx} = \frac{v^2x^2 - x^2}{2x^2v}$$

$$v + x \cdot \frac{dv}{dx} = \frac{v^2 - 1}{2v}$$

$$\therefore x \cdot \frac{dv}{dx} = \frac{v^2 - 1}{2v} - v = \frac{v^2 - 1 - 2v^2}{2v}$$

$$x \cdot \frac{dv}{dx} = \frac{-(1 + v^2)}{2v}$$

$$\frac{2v}{(1 + v^2)} \cdot dv = \frac{dx}{x}$$

$$\therefore - \int \frac{2v}{(1 + v^2)} dv = \int \frac{dx}{x} + \ln(c)$$

$$= -\ln(v^2 + 1) = \ln(x) + \ln(c)$$

$$x \equiv \ln(v^2 + 1) = -\ln(x) + \ln(c)$$

$$\therefore v^2 + 1 = \frac{c}{x}$$

$$\frac{y^2}{x^2} + 1 = \frac{c}{x} \Rightarrow y^2 + x^2 = cx$$

$$\text{let } y = vx, \quad \therefore v = \frac{y}{x}$$

$$\therefore \frac{dy}{dx} = 1 + \frac{vx}{x} = 1 + v$$

$$= v + x \cdot \frac{dv}{dx} = 1 + v \Rightarrow x \cdot \frac{dv}{dx} = 1$$

$$1 \Rightarrow \frac{dx}{x} = dv$$

$$\therefore \int \frac{dx}{x} = \int dv + C$$

$$= \ln(x) = v + c$$

$$y = \ln\left(\frac{x}{D}\right) \cdot x$$

Bernoülli's Differential Equation

A differential equation of form $y' + p(x)y = Q(x)y^n, n \in \mathbb{R}/\{0, 1\}$ called Bernocilli's Differential equation

Method to solve:

i.) Divide by y^n

$$y^{-n} \cdot y' + p(x)y^{1-n} = Q(x) - (1)$$

$$2. \text{ put } z \leq y^{1-n}, \therefore \frac{dz}{dx} = (1-n)y^{-n} \cdot \frac{dy}{dx} \Rightarrow \underbrace{y^{-n} \cdot \frac{dy}{dx} = \frac{1}{1-n} \cdot \frac{dz}{dx}}_{(2)}$$

3. Substitute (2) in (1):

$$(1) \rightarrow \frac{1}{1-n} \cdot \frac{dz}{dx} + P(x) \cdot z = Q(x)$$

$$\Rightarrow z' + (1-n)P(x) \cdot z = (1-n)Q(x)$$

This is FLDE. in dependent variable z

\therefore Solution:

$$Z \cdot (I \cdot F) = \int (1-n)Q(x) \cdot IF \cdot dx + C, IF = e^{\int (1-n) \cdot P(x) \cdot dx}$$

Solve following:

$$a) y' + 2y = y^2$$

A. Divide by y^2 :

$$y^{-2} \cdot y' + 2y^{-1} = 1$$

$$\text{put } z = y^{-1} \quad \therefore z \frac{dz}{dx} = -y^{-2} \cdot \frac{dy}{dx}$$

\therefore (1) becomes:

$$-\frac{dz}{dx} + 2z = 1 \Rightarrow \frac{dz}{dx} - 2z = -1$$

This is FLDE.

$$\therefore P(x) = -2, \quad Q(x) = -1$$

$$IF = e^{\int P dx} = e^{\int -2 \cdot dx} = e^{-2x}$$

\therefore General Solution:

$$\therefore z \cdot e^{-2x} = \int -e^{-2x} dz + c/2$$

$$zx e^{-2x} = \frac{1}{2} e^{-2x} + \frac{c}{2}$$

$$\text{now } z = y^{-1}$$

$$\therefore y^{-1} \cdot e^{-2x} = \frac{1}{2} e^{-2x} + \frac{c}{2}$$

$$\therefore y = \frac{2}{1 + e^{2x} \cdot c}$$

Divide by y^4 :

$$y^{-4} \cdot \frac{dy}{dx} - y^{-3} \cdot \tan(x) = \sec(x) - (1)$$

$$\text{let } z = y^{-3}, \quad \frac{dz}{dx} = -3 \cdot y^{-4} \cdot \frac{dy}{dx}$$

multiply θ by -3 & substitute $\frac{dz}{dx}$

$$+3, \quad z \tan(x) = -3 \sec(x)$$

$$\begin{aligned}\frac{dz}{dx} \\ P(x) = 3 \tan(x) \quad Q(x) = -3 \sec(x) \\ I : F = e^{\int P \cdot dx} = e^{3 \int \tan x \cdot dx} = e^{\ln(\sec^3(x))} = \sec^3(x)\end{aligned}$$

multiply (2) by I.F.

$$\sec^3(x) \cdot \frac{dz}{dx} + 3 \cdot \tan(x) \cdot \sec^3(x)z = -3 \sec^4(x)$$

Apply reverse product rule: $uv' + vu' = (4v)'$

$$\Rightarrow \frac{d}{dx} (\sec^3(x) \cdot z) = -3 \sec^4(x)$$

$$\int sc.^4 \rightarrow \int se^2 \cdot sc^2$$

Integrate both sides: $\rightarrow (1 + t^2) sc^2$

$$\sec^3(x) \cdot z = -3 \int \sec^4(x) \cdot dx + c \quad (x)$$

$$= \int (1 + \infty) dt$$

$$= -3 \left[\tan(x) + \frac{\tan^3(x)}{3} \right] + c$$

$$\Rightarrow \frac{\sec^3(x)}{y^3} = -3 \tan(x) = \tan^3(x) + c$$

$$\therefore y = \frac{1}{\sec^5(x) \cdot \sqrt[3]{-3 \tan(x) - \tan^3(x) + c}}$$

$$d) \frac{dy}{dx} + \tan(x) \tan(y) = \cos(x) \cdot \sec(y)$$

A) This is B.D.E.

Divide by $\sec(y)$

$$\cos(y) \frac{dy}{dx} + \tan(x) \sin(y) = \cos(x) \quad - (1)$$

$$\text{let } z = \sin(y). \quad \frac{dz}{dx} = \cos(y) \cdot \frac{dy}{dx}.$$

Substitute this in (1)

$$\frac{dz}{dx} + \tan(x) \cdot z = \cos(x)$$

$$\text{let IF} = e^{\int \tan(x) \cdot dx} = e^{\ln(\sec(x))} = \sec(x)$$

multiply both sides by T.F.

$$\sec(x) \cdot \frac{dz}{dx} + \tan(x) \cdot \sec(x) \cdot z = 1$$

$$\equiv \frac{d}{dx} (\sec(x) \cdot z) = 1 \Rightarrow \sec(x) \cdot z = x + c \Rightarrow \sec(x) \cdot \sin(y) = x + c.$$

$$\therefore y = \sin^{-1}(\cos(x) \cdot (x + c))$$

$$d) \frac{dy}{dx} + x \cdot \sin(2y) = x^3 \cdot \cos^2(y)$$

Divide by $\cos^2(y)$:

$$\sec^2(y) \cdot \frac{dy}{dx} + \frac{x \cdot 2 \sin(y) \cos(y)}{\cos^2(y)} = x^3$$

$$\sec^2(y) \frac{dy}{dx} + 2x \cdot \tan(y) = x^3 - 10$$

$$\text{let } z = \tan(y) \quad \frac{dz}{dx} = \sec^2(y) \cdot \frac{dy}{dx}$$

substitute this in (1):

$$\frac{dz}{dx} + 2x \cdot z = x^3$$

$$\text{let I.F } e^{\int 2x \cdot dx} = e^{x^2}, \text{ demultiply both sides by I.F } e^{x^2} \cdot \frac{dz}{dx} + z \cdot 2x \cdot e^{x^2} = x^3 e^{x^2}$$

$$\Rightarrow \frac{d}{dx} (e^{x^2} \cdot z) = x^3 e^{x^2}$$

$$\begin{aligned}
 e^{x^2} z &= \int x^3 \cdot e^{x^2} dx + C \quad \longrightarrow \quad \begin{array}{l} \text{let } t = x^2, \quad dt = 2x \cdot dx \\ \downarrow \\ \frac{1}{2} \int t \cdot e^t dt \end{array} \\
 &= \frac{1}{2} e^{x^2} (x^2 - 1) + C \\
 &= \tan(y) \cdot e^{x^2} = \frac{1}{2} e^{x^2} (x^2 - 1) + C \quad \begin{array}{l} = \frac{1}{2} (t \cdot \int e^t dt - \int \int e^t dt \cdot dt) \\ = \frac{1}{2} (t \cdot e^t - e^t) = \frac{1}{2} e^t (t - 1) \end{array} \\
 y &= \tan^{-1} \left(\frac{1}{2} (x^2 - 1) + e^{-x^2} \cdot C \right)
 \end{aligned}$$

- 1.) $\frac{dy}{dx} + y = xy^3$ 2.) $\frac{dy}{dx} - y \tan(x) = y^2 \cdot \sec(x)$
 3.) $\frac{dy}{dx} + \frac{y}{x} = \frac{y^2}{x} \ln(x)$
 4.) $\frac{dy}{dx} + xy = x^3 y^3$

Partial Differentiation

If $z = f(x, y)$ it can be differentiated partially w.r.t. x or y . $\frac{\partial z}{\partial x} = \lim_{\Delta x \rightarrow 0} \frac{f(x+\Delta x, y) - f(x, y)}{\Delta x}$, Here we treat y as constant $z_x = \frac{\partial z}{\partial x}$

e.g.: $z(x, y) = x^2 + y^2 + 2xy$,

$$z_y = \frac{\partial z}{\partial x} \frac{\partial z}{\partial y} = 2x + 0 + 2y = 0$$

or w.r.t. y by: $\frac{\partial z}{\partial y} = \lim_{\Delta y \rightarrow 0} \frac{f(x, y+\Delta y) - f(x, y)}{\Delta y}$ [we treat x constant] $\frac{\partial z}{\partial y} = 0 + 2y + 2x$

find $\frac{\partial f}{\partial x}$ & $\frac{\partial f}{\partial y}$ in following:

a) $f(x, y) = x^3 + 3x^2y + xy^3$

A) $\frac{\partial f}{\partial x} = 3x^2 + 6xy + y^3$, $\frac{\partial f}{\partial y} = 0 + 3x^2 + 3xy^2$

b) $f(x, y) = 2x \cos(y) + 3x^2y$

A) $\frac{\partial f}{\partial x} = 2 \cos(y) + 6xy$ $\frac{\partial f}{\partial y} = -2x \sin(y) + 3x^2$

C)

$$f(x, y) = x \tan^{-1} \left(\frac{y}{x} \right)$$

$$f_x = \frac{1}{1 + y^2/x^2} \cdot \frac{\partial}{\partial x} \cdot \left(\frac{y}{x} \right) = \frac{y}{x^2} \cdot \frac{x^2}{x^2 + y^2} \cdot -\frac{1}{x^2} = \frac{-y}{x^2 + y^2}$$

$$\Delta f_y = \frac{1}{1 + y^2/x^2} \cdot \frac{\partial}{\partial y} \left(\frac{y}{x} \right) = \frac{x^2}{x^2 + y^2} \cdot \frac{1}{x} = \frac{x}{x^2 + y^2}$$

d) $f(x, y) = x^3 - x^2 \sin(y) - y$

$$f_x = 3x^2 - 2x \sin(y), \quad f_y = -x^2 \cos(y) - 1$$

Higher Order Partial Derivative

- $\frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right) = f_{xx}$

- $\frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right) = f_{yy}$

- $\frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right) = f_{xy}$

- $\frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right) = f_{yx}$

- If $f(x, y)$ is continuous function, $f_{xy} = f_{yx}$

1. find I & II order partial derivatives of

a) $t = x^2 y$

A) $f_x = 2xy$, $f_y = x^2$, $f_{xy} = 2x$, $f_{yx} = 2x$, $f_{xx} = 2y$, $f_{yy} = 0$

b) $x^3 f(x, y) = x^3 \sin(y)$

$$f_x = 3x^2 \sin(y), f_{xx} = 6x \sin(y), f_{yx} = 3x^2 \cos(y)$$

$$f_y = x^3 \cos(y), f_{yy} = -x^3 \sin(y), f_{xy} = -3x^2 \sin(y)$$

Differentials

If $z = f(x, y)$, dz, dx, dy are known as differentials. in z, x, y respectively.

$$dz = \frac{\partial z}{\partial x} dx + \frac{\partial z}{\partial y} dy$$

1.) find the differentials in f of if $f = \frac{x^3}{3} - xy^2$

$$df = \frac{df}{\partial x} \frac{\partial f}{\partial x} dx + \frac{\partial f}{\partial y} dy$$

$$df = (x^2 - y^2) dx - 2xy dy$$

Exact Differential Equation

- A Differential equation of the form $M(x, y)dx + N(x, y)dy = 0$ is said to be exact differential equation.

such that $\frac{\partial \mu}{\partial x} = m(x, y)$, & $\frac{\partial \mu}{\partial y} = N(x, y)$

ie, $Mdx + Ndy = \frac{\partial \mu}{\partial x} dx + \frac{\partial \mu}{\partial y} dy = d\mu$

$$\therefore \text{Solution is } \int d\mu \Rightarrow \mu(x, y) = c$$

Method to find exact or not

- If Diff. eqn. is exact then $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

eg: $(1 - x) \cdot dx - (1 + y)dy = 0$

$$M = 1 - x, \quad N = -(1 + y)$$

$$\frac{\partial M}{\partial y} = 0, \quad \frac{\partial N}{\partial x} = 0 \Rightarrow \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \quad \therefore \text{Eqn. is exact.}$$

Method to solve E.D.E

Solution: $\int M dx + \int [\text{Terms in } N \text{ not containing } x] dy = c$

Solve $(1-x)dx - (1+y)dy = 0$

A) Solution: $\int (1-x)dx + \int -(1+y)dy = c/2$ (soy)

$$\begin{aligned} x - \frac{x^2}{2} - y - \frac{y^2}{2} &= c/2 \\ \Rightarrow x - y &= \frac{x^2 + y^2}{2} + c/2 \\ 2(x - y) &= x^2 + y^2 + c \end{aligned}$$

2. $(3x^2 + 4xy) dx + (2x^2 + 2y) dy = 0$

$$\begin{aligned} M &= 3x^2 + 4xy, & \frac{\partial M}{\partial y} &= 4x \\ N &= 2x^2 + 2y, & \frac{\partial N}{\partial x} &= 4x \end{aligned} \quad \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \Rightarrow \text{eqn. is Exact}$$

Solution: $\int M \cdot dx + \int [\text{terms in } N \text{ not containing } x] dy = c$

$$\begin{aligned} \int (3x^2 + 4xy) dx + \int 2y \cdot dy &= c \\ x^3 + 2x^2y + y^2 &= c \end{aligned}$$

3. Why condition for exactness is $\frac{\partial M}{\partial x} = \frac{\partial N}{\partial y}$?

A) for E.D.E. $\exists u(x, y) : \frac{\partial u}{\partial y} = N(x, y) \cdot \frac{\partial u}{\partial x} = m(x, y)$

consider $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial N}{\partial x}, \quad \frac{\partial^2 u}{\partial y \partial x} = \frac{\partial M}{\partial y}$

since $u(x, y)$ represents a family of curve and

it is continuous, $\frac{\partial^2 u}{\partial x \partial y} = \frac{\partial^2 u}{\partial y \partial x} \Rightarrow \frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$

2. Anodian way]:

$$m = 3x^2 + 4xy, (1) \quad N = 2x^2 + 2y - (2)$$

$$\frac{\partial M}{\partial y} = 4x$$

$$\rightarrow \cdot m \quad 0, 4xy + \Psi(x)$$

$$= 4xy + \psi(x) - 15$$

comparing (1) & (3) $\psi(x) = 3x^2$

4. $(2x \cos(y) + 3x^2y) dx + (x^3 - x^2 \sin y - y) dy = 0$

$$\begin{aligned} M &= 2x \cos(y) + 3x^2y & N &= x^3 - x^2 \sin(y) - y \\ \frac{\partial M}{\partial y} &= -2x \sin(y) + 3x^2, & \frac{\partial N}{\partial x} &= 3x^2 - 2x \sin(y) \\ \frac{\partial M}{\partial y} &= \frac{\partial N}{\partial x} \Rightarrow \text{eqn. is EDE} \end{aligned}$$

Solution: $\int (2x \cos(y) + 3x^2y) dx + \int -y dy \pm c$

$$x^2 \cos(y) + x^3y - \frac{y^2}{2} = c$$

HW:

1. Solve:

- a) $(5x^4 + 3x^2y^2 - 2xy^3) dx + (2x^3y - 3x^2y^2 - 5y^4) dy = 0$
 b) $(2xy + y - \tan(y))dx + (x^2 - x \tan^2(y) + \sec^2(y) + 2y) dy = 0$

2. (another uy).

$$M = 3x^2 + 4xy, N = 2x^2 + 2y$$

Define $f(x, y)$ f $f_x = m.f_y = N$

Then solution is given by $f(x, y) = c_1$

1. Integrate f_x with respect to x to find $f(x, y)$:

$$f(x, y) = \int (3x^2 + 4yx) dx = x^3 + 2x^2y + \psi(y)$$

Differentiate this curty to find $\psi(y)$:

$$\partial f_y = 2x^2 + \frac{d\psi}{dy}$$

Substitute $f_y = N$, (by def.)

$$2x^2 + \frac{d\psi}{dy} = 2x^2 + 2y \Rightarrow \frac{d\psi}{dy} = 2y$$

Integrate $d\psi$ sixy urty: $\psi(y) = y^2$

substitute $\psi(y)$ in $f(x, y)$:

$$f(x, y) = x^3 + y^2 + 2x^2y$$

The Solution is $f(x, y) = c$:

$$\begin{aligned} &: \frac{x^3 + y^2 + 2x^2y}{2} = c \\ x^4 + y^2 + 2x^2y &= x^4 - x^3 + c \\ \Rightarrow (y + x^2)^2 &= x^4 - x^3 + c \\ y + x^2 &= \pm \sqrt{x^4 - x^3 + c} \\ y &= -x^2 \pm \sqrt{x^4 - x^3 + c} \end{aligned}$$

? $\frac{dy^4}{dx} + \frac{xy}{1-x^2} = x\sqrt{y}$
 divide by \sqrt{y} .

$$y^{-1/2} \frac{dy}{dx} + \frac{xy^{1/2}}{1-x^2} = x \quad (1)$$

let $z = y^{1/2}$, $\frac{dz}{dx} = \frac{1}{2}y^{-1/2} \frac{dy}{dx}$
 \therefore (1) become:

$$\frac{dz}{dx} + \underbrace{\frac{dx}{2(1-x^2)}}_p \cdot z = \frac{1}{2}x.$$

$$\text{let } f = e^{\frac{1}{e} \int \frac{x}{1-x^2} dx} = (1-x^2)^{-1/4} \quad -\frac{1}{2} \int \frac{x}{1-x^2} dx \quad t = 1-x^2 \quad dt = dx \cdot (-2x)$$

$$s : \mathbb{Z} \cdot F = \int Q \cdot F \cdot dx + c = -2x \cdot dx$$

$$\begin{aligned} \Psi \cdot (1-x^2)^{-1/4} &= \int \frac{1}{2}x \cdot (1-x^2)^{-1/4} dx + c \quad -\frac{1}{4} \int \frac{dt}{t} \Rightarrow \Rightarrow n(t) \\ &= \frac{1}{2} \int x \cdot (1-x^2)^{-1/4} dx + c \quad - (2) \end{aligned}$$

$$\therefore \text{Solution is: } y \cdot F = \int Q \cdot F \cdot dx + c$$

$$\text{let } t = 1-x^2, dt = -2x dx$$

\therefore (2) becomes:

$$\begin{aligned} y \cdot (1-x^2)^{-1/4} &= -\frac{1}{4} \int t^{-1/4} dt + c \\ &= -\frac{1}{4}x \frac{t^{3/4}}{-1/4+1} = -\frac{1}{3} \cdot t^{3/4} + c \\ z \cdot (1-x^2)^{-1/4} &= -\frac{1}{3} (1-x^2)^{3/4} + c \\ z &= -\frac{1}{3} (1-x^2) + c \cdot (1-x^2)^{1/4} \end{aligned}$$

$$z = \sqrt{y}$$

$$\therefore y = \left[\sqrt[4]{c \cdot (1-x^2)} - \frac{1}{3} (1-x^2) \right]^2$$

$$\begin{aligned} \frac{dy}{dx} + x \cdot \sin(2y) &= x^3 \cdot \cos^2(y) \\ \equiv \frac{dy}{dx} + x \cdot 2 \cdot \sin(y) \cos(y) &= x^3 \cdot \cos^2(y) \end{aligned}$$

Divide by $\cos^2(y)$:

$$\sec^2(y) \cdot \frac{dy}{dx} + 2x \cdot \tan(y) = x^3$$

$$\text{let } z = \tan(y) = \frac{dz}{dx} = \sec^2(y) \cdot \frac{dy}{dx}.$$

$$\frac{dz}{dx} + 2x \cdot z = x^3$$

I.F = $e^{\sqrt{2x}} = e^{x^2}$, & multiply by it:

$$\begin{aligned}
e^{x^2} \frac{dz}{dx} + 2x \cdot e^{x^2} \cdot z &= x^3 \cdot e^{x^2} \\
\Rightarrow e^{x^2} \cdot z &= \int x^3 \cdot e^{x^2} dx + c \\
&= \frac{1}{2} e^{x^2} (x^2 - 1) + c \\
\therefore \tan(y) &= \frac{1}{2} (x^2 - 1) + e^{-x^2} \cdot c
\end{aligned}$$

3. $\frac{dy}{dx} + y \tan(x) = y^3 \cdot \sec(x)$

EDE -Hw-1

$$\begin{aligned}
&\underbrace{(5x^4 + 3x^2y^2 - 2xy^3)}_M dx + \underbrace{(2x^3y - 3x^2y^2 - 5y^4)}_N dy = 0 \\
M_{xy} &= 6x^2y - 6xy^2, N_y = 6x^2y - 6xy^2 \\
M_y &= N_y \quad \therefore \text{EDE}
\end{aligned}$$

\therefore So ln: $\int_{\text{cor}} m dx + \int (N \text{ } \cancel{dx}) dy = C$

$$= x^5 - y^5 + x^3y^2 - x^2y^3 = c$$

29/9

$$(3x^2 + 6xy^2) dx + (6x^2y + 4y^3) dy = 0$$

Solve:

a) $[\cos(x) \tan(y) + \cos(x + y)] dx + [\sin(x) \cdot \sec^2(y) + \cos(x + y)] dy = 0$

A) $M = \cos(x) \tan(y) + \cos(x + y) \cdot \frac{\partial M}{\partial y} = \cos(x) \cdot \sec^2(y) - \sin(x + y)$

$$N = \sin(x) \sec^2(y) + \cos(x + y), \frac{\partial N}{\partial x} = \sec^2(y) \cdot \cos(x) - \sin(x + y)$$

$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x} \Rightarrow$ Equation is exact.

\therefore Solution is:

$$\begin{aligned}
&\int_{y \cdot \text{cons}} M \cdot dx + \underbrace{\int (\text{terms in } N \text{ not containing } x) dy}_0 = c \\
&\tan(y) \int \cos(x) dx + \int \cos(x + y) dx = c \\
&\tan(y) \cdot \sin(x) + \sin(x + y) = c
\end{aligned}$$

1. $(y \cos(x) + 1) dx + \sin(x) dy = 0$

2. $(\sec(x) \tan(x) \tan(y) - e^x) dx + (\sec(x) \sec^2(y) dy = 0$

Linear D.E with Constant Coeffis

It is eqn of form.

$$a_0 \cdot \frac{d^n y}{dx^n} + a_1 \frac{d^{n-1} y}{dx^{n-1}} + \cdots + a_{n-1} \frac{dy}{dx} + a_n y = \phi(x)$$

where $a_i \in \mathbb{R}$,

If $\phi(x) = 0$: to solve this we have to change the equation to symbolic form. is $(a_0 D^n + Da, D^{n-1} + \cdots) s = 0$
It's Auxillary equation is: $(a_0 m^n + a_1 m^{n-1} + \cdots) = 0$

From the auxillary equation we get the roots, m_1, m_2, \dots Now we proceed by following rules. (which depends on nature of roots.

Roots	complimentary f_x
1 Roots are Rd equal $m_1 = m_2$ $m_1 \neq m_2$	$(c_1 + c_2 x) e^{m_1 x}$ $c_1 e^{m_1 x} + c_2 e^{m_2 x}$
2) $m_1 = m_2 = m_3$ 3) $m_1 \neq m_2 \neq m_3$.	$(c_1 + c_2 x + c_3 x^2) e^{m_1 x}$ $(c_1 e^{m_1 x} + c_2 e^{m_2 x} + c_3 e^{m_3 x})$
4) $m_1 = m_2 \neq m_3$ 4) $\mathbb{I} : \alpha \pm i\beta$	$(c_1 + c_2 2e^{m_1 x} + c_3 e^{m_3 x})$ $e^{\alpha x} (c_1 \cos(\beta x) + c_2 \sin(\beta x))$

- From the nature of roots, we get complimentary function, Hence the Solution is:

$$y = C \cdot F$$

1. Solve $\frac{d^2 y}{dx^2} + \frac{f}{dx} + 6y = 0$

Symbolic form: $(D^2 + 5D + 6)y = 0 \Rightarrow (D + 3)(D + 2) = 0$

\therefore roots are: $m = -3, -2$

Real & distinct.

\therefore complimentary function is : $c_1 \cdot e^{m \cdot x} + c_2 \cdot e^{m_2 x}$

$$= c_1 e^{-2x} + c_2 e^{-3x}$$

\therefore Solution is: $y = c_1 e^{-2x} + c_2 e^{-3x}$

2 Solve $(D^3 + 1)y = 0$

$\rightarrow D^3 = -1 \Rightarrow$ roots are: $-1, \frac{1}{2} \pm \frac{\sqrt{2}}{2}i$

usang: $(a + b)(a^2 - ab + b^2)$

$$\rightarrow (D + 1)(D^2 - D + 1) = 0$$

$$\Rightarrow D + 1 = 0 \Rightarrow \text{root} = -1$$

$$D^2 - D + 1 = 0 \Rightarrow \text{root} = \frac{1 \pm \sqrt{3}i}{2}$$

CF : $e^{1/2x} \left(c_1 \cos\left(\frac{\sqrt{3}}{2}x\right) + c_2 \sin\left(\frac{\sqrt{3}}{2}x\right) \right) + c_3 \cdot e^{-x}$

To find particular seder integral $\frac{\phi(e)}{5}$

Case

$$I : \phi(x) = e^{ax}, \text{ put } D = a$$

$$\text{e.g: } \frac{d^2y}{dx^2} - 13\frac{dy}{dx} + 12y = e^{-2x}$$

$$\begin{aligned} &\rightarrow \underbrace{(D^2 - 12D + 12)}_{=0 \rightarrow \text{roots}} y = e^{-2x} \\ &\therefore C \cdot F = c_1 e^x + c_2 e^{12x} \end{aligned}$$

$$\therefore \text{Porticular integral : } PI = \frac{e^{-2x}}{D^2 - 13D + 12}, D = -2,$$

$$\Rightarrow \frac{e^{-2x}}{4 + 26 + 12} = \frac{e^{-2x}}{42}$$

$$\therefore \text{Solution: } y = CF + PI$$

$$\begin{aligned} &= c_1 e^x + c_2 e^{12x} + \frac{e^{-2x}}{42} \\ 6D^2y - D_y - 2y &= e^{4x}, \quad 6 \end{aligned}$$

$$\therefore \text{Auy-fx} = 6D^2 - D - 2, \text{ rooks} = \frac{+1 \pm \sqrt{1 + 4 \times 6 \times 2}}{12} = \frac{1 \pm 7}{12} \Rightarrow \frac{2}{3} - \frac{1}{2}$$

$$\therefore CF = c_1 e^{2/3x} + c_2 e^{1/2x}$$

$$PI = \frac{e^{4x}}{6D^2 - D - 2} = \frac{e^{4x}}{6 \times 16 - 4 - 2} = \frac{e^{4x}}{90}$$

\therefore Solution:

$$\begin{aligned} y &= CF + PF \\ &= c_1 e^{\frac{2}{3}x} + c_2 e^{\frac{1}{2}x} + \frac{e^{4x}}{90} \\ y &= c_1 \sqrt[3]{e^x}^2 + c_2 \sqrt{e^x} + e^{4x}/90 \end{aligned}$$

Particular Integral

case 2 : $\phi(x) = \cos(ax)$ or $\sin(ax)$, put $D^2 = a - a^2$?. Solve $(0^2 + 4)y = \cos(3x)$

Aux. $f_x = D^2 + 4$, roots $= \pm 2i$

$$\therefore CF = e^{ox} (c_1 \cdot \sin(2x) + c_2 \cdot \cos(2x)) = c_1 \cdot \sin(2x) + c_2 \cdot \cos(2x)$$

$$PI = \frac{\cos(3x)}{D^2 + 4} = \frac{\cos(3x)}{-9 + 4} = \frac{\cos(3x)}{-5}$$

$$\therefore y = CF + PI$$

$$= c_1 \sin(2x) + c_2 \cdot \cos(2x) - \frac{\cos(3x)}{5}$$

$$\text{II? } (D^2 - 3D + 2)y = \sin(3x)$$

Aux. $f_x : D^2 - 3D + 2 \rightarrow \text{roots} : 1, 2$

$$\therefore CF = c_1 e^x + c_2 e^{2x}$$

$$D^2 = -9$$

$$PI = \frac{\sin(3x)}{D^2 - 3D + 2} =$$

$$\begin{aligned}
&= -\frac{\sin(3x)}{67+3D} = \frac{-\sin(52)}{3D+7} \\
&= \frac{-\sin(3x)(30-7)}{9D^2-49} \\
&= \sin(3x) \cdot (3D-7) \\
&= +81+49 \\
&= \frac{\sin(3x)(3D-7)}{130} \\
&= \frac{1}{130} \left(\frac{3D \cdot \sin(3x)}{\frac{d \sin(x)}{dx}} - 7 \cdot \sin(3x) \right) \\
&= \frac{1}{130} (9 \cos(3x) - 7 \sin(30))
\end{aligned}$$

\therefore Solution: $c_1 e^x + c_2 e^{2x} + \frac{1}{130} (9 \cos(3x) - 7 \sin(3x))$

$$(D^2 - 2D - 8)y = 4 \cos(2x) + e^{4x}$$

Aus. $f_n = D^2 - 2D - 8 \Rightarrow (D-4)(D+2) \Rightarrow$ roots 24, -2,

$$\therefore C \cdot F = c_1 e^{4x} + c_2 e^{-2x}$$

$$\begin{aligned}
PI_1 &= \frac{4 \cos(2x)}{D^2 - 2D - 8} \quad D^2 = -4 \\
&= \frac{4 \cos(2x)}{-4 - 2D - 8} = -\frac{4 \cos(2x)}{-2D + 12} \\
&\Rightarrow \frac{-2 \cos(2x)(D-6)}{(D+6)(D-6)} = \frac{-2 \cos(2x)(D-6)}{D^2 - 36} \\
&= \frac{2 \cos(2x)(D-6)}{40} \\
&= \frac{\cos(2x)(D-6)}{20} \\
&= \frac{D \cdot \cos(2x) - 6 \cos(2x)}{20} \\
&= -\frac{\sin(2x) + 3 \cos(2x)}{10}
\end{aligned}$$

$$\begin{aligned}
PI_2 &= \frac{e^{4x}}{D^2 - 2D - 8} & D = 4 \quad * : 1f f(x) = e^{ax} \\
&= \frac{e^{4x}}{16 - 8 - 8} \quad \frac{1}{f(a)} = 0, \frac{1}{f(D)} e^{ax} = \frac{x}{\phi} \\
&= \frac{e^{4x}}{0} & \frac{1}{0} \quad f(x) = \cos(ax) \& \frac{1}{f((2))} = 0
\end{aligned}$$

$$\begin{aligned}
PI_2 &= \frac{e^{4x}}{(D-4)(D+2)} \quad * \text{ If } f(x) = \sin(ax) d \frac{1}{f(D^2)} = 0 \\
&= \frac{1}{D-4} \times \frac{e^{4x}}{D+2} \quad \frac{1}{f(D)} \sin(ax) = \frac{-x \cdot \cos(ax)}{29} \\
\Rightarrow &= \frac{x e^{4x}}{4+2} = \frac{x e^{4x}}{6}
\end{aligned}$$

\therefore Solution is: $CF + PI_1 + PI_2$

$$= c_1 e^{4x} + c_2 e^{-2x} - \frac{\sin(2x) + 3 \cos(2x)}{10} + \frac{x e^{4x}}{6}$$

$$2. (D^2 - 9)y = 1 + 5e^{4x} + 2e^{3x}$$

A)

$$\text{Ans } F_n = D^2 - 9 = 3, -3$$

$$CF = c_1 e^{3x} + c_2 e^{-3x}$$

$$PI_1 = \frac{e^{0x}}{D^2 - 9} = D^2 = 0$$

$$= \frac{1}{-9} \quad D = 4$$

$$PI_2 = \frac{5e^{4x}}{D^2 - 9} = \frac{5e^{4x}}{7} = \frac{1}{(D-3)} \cdot \frac{2e^{3x}}{(D+3)}$$

$$= \frac{2e^{3x}}{D^2 - 9} = \frac{2xe^{3x}}{6} = \frac{xe^{3x}}{3}$$

$$\therefore \text{Solution: } c_1 e^{3x} + c_2 e^{-3x} - \frac{1}{9} + \frac{5e^{4x}}{7} + \frac{xe^{3x}}{3}$$

$$3. (0^2 + 16)y = \cos(4x)$$

Case 3: $\phi(x) = x^m$

To find PI. $\frac{1}{f(D)}\phi(x)$, take $[f(D)]^{-1}\phi(x)$

→ expand binomially, neglecting higher powers of D , (upto m^{th} power)

$$(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \frac{n(n-1)(n-2)}{3!}x^3 + \dots$$

$$(1+x)^{-1} = 1 - x + x^2 - x^3 + x^4 + \dots$$

$$(1+x)^{-2} = 1 - 2x + 3x^2 - 4x^3 + \dots$$

$$1. (D^2 + D + 1)y = x^2$$

$$\text{Aux. } f(D) = D^2 + D + 1, \text{ roots: } \frac{-1 \pm \sqrt{-3}}{2} = \frac{-1 \pm i\sqrt{3}}{2}$$

$$EF = e^{-\frac{1}{2}x} \left(C_1 \cos(\sqrt{3}x) + C_2 \sin(x\sqrt{3}) \right)$$

$$PI = \frac{x^2}{D^2 + D + 1} = (1 + (D + D^2))^{-1} x^2$$

$$= \left(1 - (D + D^2) + (D + D^2)^2 - (D + D^2)^3 + \dots \right) x^2$$

$$= [1 - D - D^2 + D^2 + 2D^3 + D^4] x^2$$

$$= x^2 - D(x^2) - D^2(x^2) + D^2(x^2) + 2D^3(x^2) + D^4(x^2)$$

$$= x^2 - D(x^2) + 2D^3(x^2) + D^4(x^2)$$

$$= x^2 - 2x + 0 + 0 = x^2 - 2x$$

$$\therefore \text{Solution: } y = cF + PI = e^{-\frac{1}{2}x} (c_1 \cos(x\sqrt{3}) + c_2 \sin(x\sqrt{3})) + x^2 - 2x \rightarrow -p$$

$$(D^2 + 2D + 1)y = 2x + x^2$$

$$- 1$$

$$\therefore F = c_1 e^{-x} + c_2 e^{-x} x$$

$$\begin{aligned}
PI_1 &= \frac{2x}{D^2 + 2D + 1} = (D^2 + 2D + 1)^{-1} (2x) \\
&= (D + 1)^{-2} (2x) \\
&= (1 - 2D + 3D^2) 2x \\
&= 1 - 2D(2x) + 3D^2(2x) \\
&= 2x - 4 + 6 = 2x - 4 \\
PI_2 &= \frac{x^2}{(D + 1)^2} = (D + 1)^{-2} (x^2) \\
&= (1 - 2D + 3D^2) x^2 \\
&= x^2 - 2D(x^2) + 3D^2(x^2) \\
&= x^2 - 4x + 6 = x^2 - 4x + 6
\end{aligned}$$

$$\begin{aligned}
\therefore \text{Solution} &= y = c_1 e^{-x} + x^2 \rightarrow -2x + 2 + c_1 e^{-x} \\
&= e^{-x} (c_1 + c_2 x) + x^2 - 2x + 2
\end{aligned}$$

$$\begin{aligned}
(2D^2 - 5D + 3)y &= \cos(3x) \cos(2x) \\
&= \frac{1}{2} (\cos(5x) - \cos(x)) & C_H C_B \\
2D^2 - 5D + 3 &= \frac{5 \pm \sqrt{25 - 24}}{4} \Rightarrow \frac{3}{2}, 1 & S_1 S_2 = \\
\therefore CF &= C_1 e^{3/2x} + C_2 e^x & S_1 C_2 = \\
PI_1 &= \frac{\frac{1}{2} \cos(5x)}{2 \cdot \frac{3}{2} - 5D + 3} & C_1 = \\
&= \frac{\frac{1}{2} \cos(5x)}{10 - 5D + 3} = -\frac{1}{2} \frac{\cos(5x)}{5D - 13} = \frac{1}{2} \frac{\cos(5x)(5D + 13)}{25D^2 - 169} \\
C_A C_B &= \frac{1}{2} [C(A + B) + C(A - B)]
\end{aligned}$$

$$\begin{aligned}
&= -\frac{1}{2} \frac{\cos(5x)(5D + 13)}{125 - 169} = \frac{\cos(58)(50 + 13)}{88} \\
&= \frac{5D(\cos(5x)) + 13 \cos(5x)}{88} \\
&= \frac{13 \cos(5x) - 25 \sin(5x)}{88} \\
PI_2 &= \frac{1}{2} \frac{\cos(x)}{2D^2 - 50 + 3} \rightarrow
\end{aligned}$$

$$\therefore \text{solution: } C_1 e^{3/2x} + C_2 e^x + \frac{1}{5668} (47 \cos(5x) + 25 \sin(5x))$$

$$(D^2 - 4D + 3)y = \sin(3x) \cos(2x)$$

A 2nd *O.D.E.* has complimentary fo \$ particular integral compl.fn is of form: $Ae^{m_1x} + Be^{m_2x} + Ce^{m_3x} \dots$ where m_1, m_2, m_3 are roots of auzillory f_n . for imaginary roots:

let a_{5bi} be the root,

$$\therefore \text{Solution is: } Ae^{(a+bi)x} + Be^{(a-bi)x}$$

$$\begin{aligned}
&\Rightarrow Ae^{ax}e^{bix} + Be^{ax}e^{-bix} \\
&= e^{ax} (Ae^{bix} + Be^{-bix}) \\
&= e^{ax} (A(\cos(bx) + i\sin(bx)) + B(\cos(bx) - i\sin(bx))) \\
&= e^{ax} ((A+B)\cos(bx) + (A-B)i\sin(bx))
\end{aligned}$$

$$\begin{aligned}
&\rightarrow e^{ax}(c \cdot \cos(bx)) \\
&\sin(3x) \cos(2x)
\end{aligned}$$

$$= \frac{1}{2} \sin(5x) + \frac{1}{2} \sin(-x)$$

$$\text{Aux. } f_n = D^2 - 4D + 3, \text{ roots } = 1, 3$$

$$= (D-1)(D-3)$$

$$\therefore CF = c_1 e^x + c_2 e^{3x}$$

$$PI_1 = \frac{\sin(5x)}{2(D^2 - 4D + 3)}, D^2 = 5$$

$$\begin{aligned}
&\Rightarrow \frac{\sin(5x)}{13-8D} \Rightarrow \frac{\sin(5x)(13+8D)}{169-64D^2} = -\frac{\sin(5x)(13+8D)}{151} \\
&= -\frac{13\sin(5x) - 40\cos(5x)}{151}
\end{aligned}$$

$$PI_2 = \frac{\sin(x)}{2(D^2 - 4D + 3)} \Rightarrow CD^2 = 1$$

$$\Rightarrow \frac{\sin(x)}{5-4D} \Rightarrow \frac{\sin^2(x)(5+4D)}{25-16D^2} = \frac{\sin(x) + 4\cos(x)}{9}$$

$$\therefore \text{Solution: } y = CF + PI_1 + PI_2$$

$$= c_1 e^x + c_2 e^{3x} - \frac{13\sin(5x) + 40\cos(5x)}{91} + \frac{5\cos(x) - 4\sin(x)}{9}$$

$$\text{Case } \forall T - e^{ax} f(x)$$

$$\lambda(D^2 + 30 + 2)y = e^{2x} \sin(x)$$

$$\text{C.F in } y = c_1 e^{-1} + c_2 e^{-2x}$$

$$P.I., = \frac{e^{2x} \sin(x)}{D^2 + 3x + 2} = e^{2x} \cdot \frac{\sin(x)}{D^2 + 3D + 2} \stackrel{D=D+a}{\sin(x)} \cdot e^{2x}$$

$$\frac{\sin(x)}{D^2 + 4D + 4 + 3D + 6 + 2} e^{23}$$

$$= \frac{\sin(x)}{D^2 + 7x + 12} e^{2x}$$

$$D^2 = -(1)$$

$$\rightarrow \frac{\sin(x)}{70 + 11} e^{221}$$

$$\Rightarrow \frac{\sin(x)(7D+11)}{\cos D^2 - 121} e^{22}$$

$$\Rightarrow \frac{\sin(x)(70-11)}{-170} (e^{2x}) \Rightarrow \frac{7\cos(x) - 11\cos\sin(x)}{9} \cdot e^{2x}$$

-Solution.

$$y = c_1 e^{-x} + c_2 e^{-2x} + \frac{11 \sin(x) - 7 \cos(x)}{170} e^{2x}$$

Ho

$$\begin{cases} (D^2 + 4D + 5)y = 12e^{-12} \cdot \cos(x) \\ (D^2 - 2D + 1)y = xe^{2x} \end{cases}$$

e.

$$\therefore CF = c_1 e^{-x} + c_2 x e^x$$

$$PI = e^{2x} \cdot \frac{x}{D^2 - 2D + 1} = D^2 \left\{ \begin{array}{l} \\ ye^{2x} - (D^2 - 2D + 1)x \end{array} \right\}$$

$$\begin{aligned} &\Rightarrow e^{2x} \cdot (D^2 - 2D + 1)^{-1} x \\ &\Rightarrow e^{2x} \cdot (D - 1)^{-2} \cdot x \\ &\Rightarrow e^{2x} \cdot (1 + 2D - 3D^2) x \\ &\Rightarrow e^{2x} \cdot (x + 2) \end{aligned}$$

$$\begin{aligned} &\Rightarrow \frac{x}{(D - 1 + 2)^2} \cdot e^{2x} \\ &\Rightarrow (D + 1)^{-2} x \cdot e^{2x} \\ &\quad (1 + 2D + 3D^2) x \cdot e^{2x} \\ &\Rightarrow (x - 2)e^{2x} \end{aligned}$$

$$\therefore \text{Solution: } y = c_1 e^x + c_2 e^x + (x - 2)e^{2x}$$

$$\begin{aligned}
& (D^3 - 3D_x^2 + 3D - 3) y = x^2 e^x \\
& \Rightarrow (D - 1)^3 \Rightarrow D = 1 \\
& \therefore c = c_1 e^x + c_2 e^x \cdot x + c_2 x^2 e^x \\
& PI = e^x \cdot \frac{x^2}{(Q - 1)^3}, \quad D \rightarrow D + 1 \\
& \Rightarrow e^x \frac{x^2}{D^3} \Rightarrow e^x \cdot (D^{-3}) x^2 \\
& e^x \cdot (D^{-2}) \frac{x^3}{3} = e^x \cdot \left(D^{-1} \cdot \frac{24}{12} \right) = e^x \cdot \frac{25}{650} \\
& D^{-3} + (D + 1 - 1)^{-3} \rightarrow (1 + (D - 1))^3 x^2 \\
& \rightarrow (1 + 3(D - 1) + 3(D - 1)^2) \geq^2 \\
& = x^2 + 3(D - 1)x^2 + 3(D^2 - 2D + 1)x^2 \\
& \Rightarrow x^2 + (3D - 3)x^2 + (3D^2 - 6D + 3)x^2 \\
& \Rightarrow x^2 + -3x^2 + 6 \\
& x^2 + 6x^3 + 6 - 12x \\
& (D^2 - 2D + 1) y = x \cdot e^x \sin(x) \\
& C \cdot f = \text{se}(c_1 + c_2 x) e^x \\
& PI = e^x \cdot \frac{x \sin(x)}{(D - 1)^2} \xrightarrow{D \rightarrow D+1} \frac{1}{2} e^x \cdot \frac{x \sin(x)}{D^2} \\
& = e^x \cdot D^1(x \cdot \sin(x))
\end{aligned} \tag{1}$$

$$\begin{aligned}
& e^x \cdot D^{-4}(-x \cdot \cos(x) + \sin(x)) \\
& e^x \cdot (-x \cdot \sin(x) + \cos(x) + \cos(x)) \\
& \Rightarrow e^x \cdot (x \sin(x) + 2 \cos(x)) \\
& \therefore \text{soln } y = (c_1 + c_2 x - x \sin(x) - 2 \cos(x)) e^x
\end{aligned}$$

Simultaneous LDE

It contains two or more dependent variables (say x, y .. and one independent variable (say t)

1. $\frac{dx}{dt} = 7x - y, \quad \frac{dy}{dt} = 5(x - y)$
A)

$$\frac{dx}{dt} - 2x = y = 0 \rightarrow (D - 7)x + y = 0 \quad (1)$$

$$\frac{dx}{dt} - \frac{dy}{5x} - \frac{dy}{dx} + 5y = 0 \rightsquigarrow (D - 5)x + -(D - 5)y = 0 \quad (2)$$

$$(D - 5) \times 0 :$$

$$(D - 7)(D - 5)x + (D - 5)y = 0$$

$$+ [(D - 5)\dot{x} - (D - 5)y = 0]$$

$$\underbrace{(D^2 - 11D + 30)}_{\text{aux ff}} x = 0$$

$$=$$

$$\therefore \text{ costs of aux ff } z$$

$$5, 6$$

$$\therefore C \cdot F_{12}x = C_1 e^{5t} + C_2 e^{6t}$$

$$D = \frac{d}{db}$$

$$\therefore (1) \text{ becomv.}$$

$$\begin{aligned} (D - 7)(c_1 e^{st} + c_2 e^{6t}) + y &= 0 \\ \Rightarrow 5c_1 e^{st} + 6c_2 e^{6t} - 7c_1 e^{st} - 7c_2 e^{-st} &= -y \\ \Rightarrow y &= 2c_1 e^{st} + c_2 e^{6t} \end{aligned}$$

$$\& \quad x = c_1 e^{5t} + c_2 e^{6t}$$

$$2\frac{dx}{dt} + 2x - 3y = t \quad , \quad \frac{dy}{dt} - 3x + 2y = e^{2t}$$

A)

$$\begin{aligned} (D + 2)x - 3y &= t \\ (D + 2)y - 3x &= e^{2t} \end{aligned} \quad (2)$$

$$(1) \times 3 : 8$$

$$3x(D + 2) - 9y = 3t \quad (3)$$

$$+ (-3x(D + 2) + (D + 2)^2 y = (D + 2)e^{2t}) \quad (9)$$

$$(3) + (4)$$

$$\begin{aligned} y((D + 2)^2 - 4) &= 3t + 4e^{2t} \\ y(D^2 + 4D - 5) &= 3t + 4e^{2t} \\ &= y(D + 5)(D - 1) \cdot \\ \therefore \text{ roots } &= -5, +1 \\ \therefore CF_{\text{an } y} c_1 e^{-5t} + C_2 e^{+t} \end{aligned}$$

$$\begin{aligned}
PI_{y_1} &= \frac{3t}{(D^2 + 4D - 5)} \\
&= -\frac{3}{5} \times \left(1 + \left(\frac{D^2}{-5} - \frac{4}{5}D\right)\right)^{-1} t \\
&= -\frac{3}{5} \times \left(1 - \frac{D^2}{-5} + \frac{4}{5}Dx^2 + \dots\right) t \\
&= -\frac{3}{5} \left(t + 0 + \frac{4}{5}\right) \\
PI_{y_1} &= \frac{4e^{2t}}{(D+5)(D-1)} \\
&\Rightarrow \frac{4}{7}t - \frac{12}{25} \\
&\Rightarrow y = CF_y + PI_{y_1} + PI_{y_2} \\
\therefore y &= C_1e^{-5t} + C_2e^t - \frac{3}{5}t + \frac{4}{7}e^{2t} - \frac{12}{25}
\end{aligned}$$

Substitute in (2):

$$\begin{aligned}
(D+2)y - e^{2t} &= 3x \Rightarrow x = \frac{1}{3} \left[(D+2)y - \frac{t}{2}e^{2t} \right] \\
x &= \frac{1}{3} \left(-5c_1e^{-5t} + c_2e^t - \frac{3}{5} + \frac{8}{7}e^{2t} \right. \\
&\quad \left. + 2c_1e^{-5t} + 2c_2e^t - \frac{6}{5}t + \frac{8}{7}e^{2t} - \frac{24}{25} - e^{2t} \right) \\
&= \frac{1}{3} \left(-3c_1e^{-5t} + 3c_2e^t + \frac{9}{7}e^{2t} - \frac{6}{5}t - \frac{39}{25} \right) \\
x &= -c_1e^{-5t} + c_2e^t + \frac{3}{7}e^{2t} - \frac{2}{5}t - \frac{13}{25}
\end{aligned}$$

3.

$$\begin{aligned}
\frac{dx}{dt} + 2y &= -\sin(t), \\
\frac{dy}{dt} &= 2x + \cos(t)
\end{aligned}$$

A) $\frac{dx}{dt} + 2y = -\sin(t) - 1$

$$\begin{aligned}
\frac{dx}{dt} + 2y &= -\sin(t) - (D)(D-2)x + (D+2)y = \cos(t) - \sin(t) \\
\frac{dy}{dt} - 2x &= \cos(t) \quad (D)
\end{aligned}$$

(1) $\times 2$,

$$2x \cdot 0 + 4y = -2\sin(t)$$

(2) $\times 0$.

$$D^2 \cdot y - 2x \cdot D = D \cdot \cos(t) - (4)$$

(3) + (4):

$$D^2 \cdot y + 4y = -3 \sin(t)$$

$$\frac{d^2 y}{dt^2} + 4y = -3 \sin(t)$$

$$\therefore y e^{4t} = \int e^{4t} x - 3 \sin(t) dt = -3 \int \sin(t) e^{4t} \cdot dt$$

$$\begin{aligned} I &= -3 \int \sin(t) e^{4t} dt \quad I_2 = \cos(t) 4e^{4t} + 4 \underbrace{\int \sin(t) \dots}_{I} \\ &= -3(\sin(t) \times 4e^{4t} - 4 \underbrace{\cos(t) \cdot e^{4t}}_I) = 40 \cos(t) e^{4t} + 4I \\ &= -8 \sin(t) e^{4t} + 32 \cos(t) e^{4t} + 32I \\ \therefore &= \frac{e^{4t}}{4^2 + 1} (4 \sin(t) - \cos(4t)) \end{aligned}$$

$$\therefore = \frac{e^{4t}}{4^2 + 1} (4 \sin(t) - \cos(\theta t))$$

or

$$(p^2 + 4) y = -3 \sin(t)$$

$$\therefore y = (c_1 \cos(2t) + c_2 \sin(2t))$$

$$PI = \frac{-3 \sin(t)}{D^2 + 4}, \quad D^2 = -1$$

$$\Rightarrow -\frac{3}{3} \sin(t) = -\sin(t)$$

$$\therefore y = c_1 \cos(2t) + c_2 \sin(2t) - \sin(t)$$

Substitute this in (2):

$$\Rightarrow x = -e^{2t} (\cos^5(-c_1 + c_2))$$

$$\Rightarrow x = -2e^{2t} (c_1 + c_2) \cos(2t) + (c_2 - c_1) \sin(2E)) - \frac{1}{16} - \cos(t)$$

$$2 = -\frac{1}{2} (\cos(t) - 2c_1 \sin(2t) + 2c_2 \cos(2t) - \sin(t))$$

$$\text{Hew 1. } \frac{dy}{dt} + 2y + x = \sin(t) \quad \frac{dx}{dt} - 4y - 2x = \cos(t) \quad \text{H)}$$

$$Dy + 2y + x = \sin(t)$$

$$Dx - 4y - 2x = \cos(t)$$

$\mathcal{C} \times \bar{\partial} D :$

$$-D^2 y + 2Dy - Dx = -\cos(t)$$

$$\oplus -4y + Dx - 42x = \sin(t)$$

$$-(+D^2 + 2D + 4) y - 2x = \sin(t) - \cos(t)$$

(20) 2

$$\therefore + \frac{2Dy + 4y + 2x = 2 \sin(t)}{4D^2 + 3)}$$

$$- D^2y = 3 \sin(t) - \cos(t)$$

$$\therefore y = \int (\cos(t) - 3 \sin(t)) dt dt$$

$$= \int (\sin(t) + 3 \cos(t)) dt$$

$$y = 3 \sin(t) \cos(t) \quad y = 2 \sin(t) + ct + 2$$

$$\therefore \text{from (0: } 8 \sin(t) + B3 \cos(t) + 266 \sin(t) - 2 \cos(t) + x = \sin(t) \therefore x = -6 \sin(t) - \cos(t) \quad x = -3 \sin(t) - 2 \cos(\theta)$$

$$-2ct + 2c_2 + c$$