

Notes on Generating Functions

Math 239: Introduction to Combinatorics

David Duan, 2019 Winter

Contents

1 Formal Power Series

1.1 Introduction

1.2 Definition

1.2.1 Formal Power Series

1.2.2 Ring of Formal Power Series, Addition and Multiplication

1.3 Inverse

1.3.1 Definition

1.3.2 Theorem: Existence of Inverse

1.4 Composition

1.4.1 Definition

1.4.2 Theorem: Existence of Composition

1.5 Remarks

1.5.1 Proposition: Coefficients after Multiplication

1.5.2 Problem-Solving: Recurrence Relation

2 Generating Series

2.1 Introduction

2.2 Definition

2.2.1 Definition: Generating Series

2.2.2 Remarks: Weight Function

2.2.3 Connections: Generating Series vs. Formal Power Series

2.3 Trivial Examples

2.3.1 k -Subsets

2.3.2 Binary Strings with k 1's

2.3.3 Two Dices

2.3.4 General Binary Strings

2.4 Sum Lemma, Product Lemmas, Geometric Series

2.4.1 The Sum Lemma

2.4.2 The Product Lemma

2.4.3 Geometric Series

2.5 Product Lemma Applications

2.5.1 Binary String Sums

2.5.2 Generalized Strings

2.5.3 Negative Binomial Theorem

2.5.4 Coin Value

1 Formal Power Series

1.1 Introduction

A **formal power series** is a generalization of a polynomial, where the number of terms is allowed to be infinite; this implies giving up the possibility of replacing the variable in the polynomial with an arbitrary number. One way to view a FPS is an *infinite ordered sequence of numbers*. In this sense, the power of the variable are used only to indicate the *order of the coefficients*, so that the coefficient of x^5 is the fifth term in the sequence.

In combinatorics, formal power series provide representations of *numerical sequences* and *multisets*, and for instance allow concise expressions for recursively defined sequences regardless of whether the recursion can be explicitly solved. This is known as the method of **generating functions**.

1.2 Definition

1.2.1 Formal Power Series

Let (a_0, a_1, a_2, \dots) be a sequence of rational numbers¹. Then

$$A(x) = a_0 + a_1x + a_2x^2 + \dots = \sum_{n \geq 0} a_n x^n$$

is called a **formal power series**. We say that a_n is the **coefficient** of x^n in $A(x)$ is write $a_n = [x^n]A(x)$. Note that each a_n must be a finite number for $A(x)$ to be a FPS.

1.2.2 Ring of Formal Power Series, Addition and Multiplication

The set of all formal power series in X with coefficients in a *commutative ring* R form another ring that is written $R[[X]]$, and called the **ring of formal power series** in the variable X over R . For $A(x) = \sum_{n \geq 0} a_n x^n \in R[[x]]$ and $B(x) = \sum_{n \geq 0} b_n x^n \in R[[x]]$, addition and multiplication are well defined:

$$\begin{aligned} A(x) + B(x) &= \sum_{n \geq 0} (a_n + b_n) x^n \\ A(x) \times B(x) &= \sum_{n \geq 0} \left(\sum_{k=0}^n a_k b_{n-k} \right) x^n \end{aligned}$$

The multiplication rule can be derived as follows:

$$\begin{aligned} A(x)B(x) &= \left(\sum_{i \geq 0} a_i x^i \right) \left(\sum_{j \geq 0} b_j x^j \right) \\ &= \sum_{i \geq 0} \sum_{j \geq 0} a_i b_j x^{i+j} && k \leftarrow i, n \leftarrow i + j = k + j \\ &= \sum_{n \geq 0} \left(\sum_{k=0}^n a_k b_{n-k} \right) x^n \end{aligned}$$

1.3 Inverse

1.3.1 Definition

Let $A(x)$ and $B(x)$ be formal power series. We say that $B(x)$ is the **inverse** of $A(x)$ if

$$A(x)B(x) = 1;$$

we denote this by $B(x) = A(x)^{-1}$ or $B(x) = \frac{1}{A(x)}$.

1.3.2 Theorem: Existence of Inverse

A formal power series has an inverse if and only if it has a non-zero constant term.

Proof. Let $A(x) = \sum_{n \geq 0} a_n x^n$ be a formal power series. Then a power series $B(x) = \sum_{n \geq 0} b_n x^n$ is the inverse of $A(x)$ if and only if it satisfies the linear equation $A(x)B(x) = 1$. Suppose to the contrary $[x^0]B(x) = 0$ and $A(x)B(x) = 1$. Then

$$\begin{aligned} 1 &= A(x)B(x) \\ &= b_0 A(x) + b_1 x A(x) + b_2 x^2 A(x) \\ &= 0 A(x) + b_1 x A(x) + b_2 x^2 A(x) \\ &= b_1 x A(x) + b_2 x^2 A(x) \end{aligned}$$

Observe RHS does not have a non-zero constant term, i.e., $[x^0](b_1 x A(x) + b_2 x^2 A(x)) = 0$, because each term of $A(x)$ is multiplied by a non-zero power of x from $B(x)$. This leads to a contradiction as LHS has a non-zero term, namely 1. Similarly, $A(x)$ cannot have an inverse if it does not have a non-zero constant term, which can be proved in a mirror argument. \square

1.4 Composition

1.4.1 Definition

The **composition** of a formal power series $A(x) = \sum_{n \geq 0} a_n x^n$ and $B(x)$ is defined by

$$A(B(x)) = \sum_{n \geq 0} a_n (B(x))^n = a_0 + a_1 B(x) + a_2 (B(x))^2 + \cdots.$$

Unlike polynomials, however, this operations is not always defined. See theorem below.

1.4.2 Theorem: Existence of Composition

Let $A(x)$ and $B(x)$ be formal power series. Then the composition $A(B(x))$ is a formal power series if the constant term of $B(x)$ is equal to zero.

Proof. Let $A(x) = a_0 + a_1 x + a_2 x^2 + \cdots$ and $B(x) = xC(x)$ (so that $[x^0]B(x) = 0$). Then

$$\begin{aligned} A(B(x)) &= A(xC(x)) \\ &= a_0 + a_1 xC(x) + a_2 (xC(x))^2 + \cdots \\ &= a_0 + a_1 xC(x) + a_2 x^2 (C(x))^2 + \cdots \end{aligned}$$

For each $k \geq 0$, note that $a_k x^k (C(x))^k$ is a formal power series (since it is the product of formal power series). Moreover, for each $n < k$, we have $[x^n](a_k x^k (C(x))^k) = 0$. Therefore,

$$\begin{aligned} [x^n]A(B(x)) &= [x^n](a_0 + a_1 xC(x) + a_2 x^2 (C(x))^2 + \cdots) \\ &= [x^n](a_0 + a_1 xC(x) + a_2 x^2 (C(x))^2 + \cdots a_n x^n (C(x))^n). \end{aligned}$$

Now, $a_0 + a_1 xC(x) + a_2 x^2 (C(x))^2 + \cdots a_n x^n (C(x))^n$ is a formal power series (as it is a sum of formal power series), so $[x^n]A(B(x))$ is well-defined. \square

1.5 Remarks

1.5.1 Proposition: Coefficients after Multiplication

Let $A(x) = \sum_{n \geq 0} a_n A(x)$ be any formal power series, then

$$[x^n]x^k A(x) = \begin{cases} [x^{n-k}]A(x) & n \geq k \\ 0 & n < k \end{cases}$$

Proof. To see this, consider

$$A(x) = 1 + 2x + 4x^2 + \cdots = \sum_{n \geq 0} 2^n x^n$$

We recognize that $[x^n]A(x) = 2^n$. Next, we multiply $A(x)$ by x^3 (so $k = 3$ in the formula above):

$$x^3 A(x) = x^3 + 2x^4 + 4x^5 + \cdots = \sum_{n \geq 0} 2^n x^{n+3}$$

It looks as if we shifted all coefficients to the right by a factor of $k = 3$:

Power of x	a_k in $A(x)$	a_k in $x^3 A(x)$	Remarks
x^0	$2^0 = 1$	0	x^0 no longer in $x^3 A(x)$
x^1	$2^1 = 2$	0	x^1 no longer in $x^3 A(x)$
x^2	$2^2 = 4$	0	x^2 no longer in $x^3 A(x)$
x^3	$2^3 = 8$	$2^0 = 1$	x^3 now has coeff that previously belongs to x^0
x^4	$2^4 = 16$	$2^1 = 2$	x^4 now has coeff that previously belongs to x^1
x^5	$2^5 = 32$	$2^2 = 4$	x^5 now has coeff that previously belongs to x^2
...
x^i	2^i	2^{i-3}	x^i now has coeff that previously belongs to x^{i-3}

Hence, when we are extracting the n th coefficient from $x^k A(x)$, if $n < k$, $[x^n]x^k A(x)$ would be 0 because the term x^n no longer exists after $A(x)$ has multiplied with x^k ; otherwise, the coefficients can be seen as shifted towards right for k positions and what we are looking for (i.e., a_n for x^n) was previously belonged to x^{n-k} . \square

Example. Let $A(x) = (1 + 2x)^2 = 1 + 4x + 4x^2$ and $B(x) = \sum_{i \geq 0} 2^i x^i$. Then

$$\begin{aligned}
 [x^n]A(x)B(x) &= [x^n](1 + 4x + 4x^2)B(x) \\
 &= [x^n]B(x) + 4[x^n]xB(x) = 4[x^n]x^2 B(x) \\
 &= 2^n + 4[x^{n-1}]B(x) + 4[x^{n-2}]B(x) \\
 &= 2^n + 2^2 \cdot 2^{n-1} + 2^2 \cdot 2^{n-2} \\
 &= 2^n + 2^{n+1} + 2^n \\
 &= 2^{n+2}.
 \end{aligned}$$

1.5.2 Problem-Solving: Recurrence Relation

Example. What is the result for $\frac{1-x}{1-2x-3x^2}$? Is it a formal power series?

Solution. Suppose this is equal to $A(x) = \sum_{n \geq 0} a_n x^n$. Cross-multiply, we get

$$\begin{aligned} 1 - x &= (1 - 2x - 3x^2)A(x) \\ &= A(x) - 2xA(x) - 3x^2A(x) \\ &= (a_0 + a_1x + a_2x^2 + \cdots) - 2x(a_0 + a_1x + a_2x^2 + \cdots) - 3x^2(a_0 + a_1x + a_2x^2 + \cdots) \\ &= (a_0 + a_1x + a_2x^2 + \cdots) - (2a_0x + 2a_1x^2 + 2a_2x^3 + \cdots) - (3a_0x^2 + 3a_1x^3 + 3a_2x^4 + \cdots) \\ &= a_0 + (a_1 - 2a_0)x + (a_2 - 2a_1 - 3a_0)x^2 + (a_3 - 2a_2 - 3a_1)x^3 + \cdots \\ &= a_0 + (a_1 - 2a_0)x + \sum_{n \geq 2} (a_n - 2a_{n-1} - 3a_{n-2})x^n \end{aligned}$$

By equating the coefficients on either side of the equation, we have equality if and only if

- $a_0 = 1$
- $a_1 - 2a_0 = -1 \implies a_1 - 2 = -1 \implies a_1 = 1$
- $(a_n - 2a_{n-1} - 3a_{n-2}) = 0$ or $a_n = 2a_{n-1} + 3a_{n-2}$ for all $n \geq 2$
 - $a_2 = 2a_1 + 3a_0 = 2 + 3 = 5$
 - $a_3 = 2a_2 + 3a_1 = 2 \cdot 5 + 3 \cdot 1 = 13$
 - \dots

Hence, the result is a formal power series: $1 + x + 5x^2 + 13x^3 + \cdots$ where the coefficients are defined recursively. \square

2 Generating Series

2.1 Introduction

A **generating function** is a way of encoding an *infinite sequence of numbers* (a_n) by treating them as the *coefficients* of a *power series*. This formal power series is the generating function. Unlike an ordinary series, this formal series is allowed to diverge, meaning that the generating function is not always a true function and the "variable" is actually an indeterminate.

2.2 Definition

2.2.1 Definition: Generating Series

Let S be a set of configurations with a weight function

$$\begin{aligned} w : S &\rightarrow \mathbb{N}_0 \\ \sigma &\mapsto w(\sigma), \end{aligned}$$

The **generating series** for S with respect to w is defined by

$$\Phi_S(s) = \sum_{\sigma \in S} x^{w(\sigma)}.$$

2.2.2 Remarks: Weight Function

One way to look at the weight function is that,

$$\left\{ \begin{array}{c} S \end{array} \right\} \xrightarrow{w} \left\{ \begin{array}{c} \mathbb{N}_0 \end{array} \right\},$$

the weight function w assigns each element $\sigma \in S$ to a non-negative integer $n \in \mathbb{N}_0$. The coefficients a_k of $\Phi_S(x)$ counts how many times an $n \in \mathbb{N}_0$ appears in the image set.

2.2.3 Connections: Generating Series vs. Formal Power Series

By collecting like-powers of x in $\Phi_S(x)$, we get

$$\Phi_S(x) = \sum_{\sigma \in S} x^{w(\sigma)} = \sum_{k \geq 0} \left(\sum_{\sigma \in S, w(\sigma)=k} 1 \right) x^k = \sum_{k \geq 0} a_k x^k,$$

where a_k denotes the number of elements in S with weight k . In other words, the coefficient of x^k in $\Phi_S(x)$ counts the *number of elements of weight k in S .

2.3 Trivial Examples

2.3.1 k -Subsets

Let S be the set of all subsets of $\{1, \dots, n\}$ and $w(\sigma) = |\sigma|$. Then the number of elements S of weight k is $\binom{n}{k}$, the number of k element subsets of $\{1, \dots, n\}$:

$$\Phi_S(x) = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots = \sum_{k \geq 0} \binom{n}{k} x^k$$

2.3.2 Binary Strings with k 1's

Let S be the set of length- n binary strings, i.e., $S = \{0, 1\}^n$, and $w(b) = \sum_i b_i$. Then the number of elements S with weight k is $\binom{n}{k}$, the number of binary strings with k 1's.

$$\Phi_S(x) = \binom{n}{0} + \binom{n}{1}x + \binom{n}{2}x^2 + \dots = \sum_{k \geq 0} \binom{n}{k} x^k$$

2.3.3 Two Dices

Let $S = \{1, \dots, 6\}^2$, $w : (a, b) \mapsto a + b$. Then

$$\begin{aligned} \Phi_S(x) &= \sum_{a,b=1}^6 x^{a+b} \\ &= x^2 + 2x^3 + 3x^4 + 4x^5 + 5x^6 + 6x^7 + 5x^8 + \dots + x^{12} \\ &= \left(\sum_{i=1}^6 x^i \right)^2 \end{aligned}$$

We can interpret the result as, for example, there are three ways for us to get a weight of 4.

Now, if we restrict $S = \{(a, b) : a \in \{1, 3, 5\}, b \in \{2, 4, 6\}\}$, with the same weight function, then

$$\begin{aligned} \Phi_S(x) &= \sum_{a,b=1}^6 x^{a+b} \\ &= x^3 + 2x^5 + 3x^7 + 2x^9 + x^{11} \\ &= (x^1 + x^3 + x^5)(x^2 + x^4 + x^6) \end{aligned}$$

This factorization will come back in the future sections.

2.3.4 General Binary Strings

Let $S = \bigcup_{n \geq 0} \{0, 1\}^n$ denote the set of all binary strings (of all lengths) and for each binary string $b \in S$, $w(b)$ returns the length of b . We then have

$$\Phi_S(x) = 1 + 2x + 4x^2 + \dots = \sum_{n \geq 0} (2x)^n = \frac{1}{1 - 2x}$$

That is, there is 1 way to build a string of length 0 (empty string), 2 ways to build a binary string of length 1 (0 and 1), 4 ways to build a binary string of length 2, etc.

2.4 Sum Lemma, Product Lemmas, Geometric Series

2.4.1 The Sum Lemma

If $A \cup B = S$ and $A \cap B = \emptyset$, then $\Phi_S(x) = \Phi_A(x) + \Phi_B(x)$.

2.4.2 The Product Lemma

Let A and B be sets of configurations with weight functions α and β respectively. If $w(\sigma) = \alpha(a) + \beta(b)$ for each $\sigma = (a, b) \in A \times B$, then $\Phi_{A \times B}(x) = \Phi_A(x)\Phi_B(x)$.

More generally, if A_1, \dots, A_k are sets, then $A_1 \times \dots \times A_k$ denotes the set of all k -tuples (a_1, a_2, \dots, a_k) where $a_i \in A_i$ for all i . Now suppose that α_i is a weight function for A_i and that w is a weight function for $A_1 \times \dots \times A_k$. If $w(\sigma) = \alpha_1(a_1) + \dots + \alpha_k(a_k)$ for each k -tuples $\sigma = (a_1, \dots, a_k)$, then $\Phi_{A_1 \times \dots \times A_k}(x) = \Phi_{A_1}(x) \dots \Phi_{A_k}(x)$.

2.4.3 Geometric Series

Recall the formula for (infinite) geometric series:

$$1 + x + x^2 + x^3 + \dots = \frac{1}{1 - x}.$$

Plugging in $x = X$, we get

$$1 + X + X^2 + X^3 + \dots = \frac{1}{1 - X}.$$

For example,

$$1 + x^3 + x^6 + x^9 + \dots = \frac{1}{1 - x^3}.$$

2.5 Product Lemma Applications

2.5.1 Binary String Sums

Exercise. Let k, n be fixed non-negative integers. How many solutions are there to $t_1 + \dots + t_n = k$, where $t_1, \dots, t_n = 0$ or 1 ?

Solution. The solutions are n -tuples (t_1, \dots, t_n) where $t_i \in \{0, 1\}$ for each $i = 1, \dots, n$. We know the answer is $\binom{n}{k}$, but we can also use the product lemma. Since we are dealing with the set $\{0, 1\}^n$, and $w(t_i) = t_i$, we get

$$\Phi_{\{0,1\}}(x) = 1x^0 + 1x^1 + 0x^2 + \dots = 1 + x$$

because there is one way to get 0 and one way to get 1. Extending this to length n , we get

$$\Phi_{\{0,1\}^n}(x) = (\Phi_{\{0,1\}}(x))^n = (1+x)^n$$

and the number of solutions is

$$[x^k](1+x)^n = \binom{n}{k}. \quad \square$$

2.5.2 Generalized Strings

Exercise. Let S be the set of all k -tuples (a_1, \dots, a_k) where each $a_i \in \mathbb{N}_0$. The weight of k -tuple $\sigma = (a_1, \dots, a_k)$ is defined as $w(\sigma) = a_1 + \dots + a_k$. Show that $\Phi_S(x) = (1-x)^{-k}$.

Solution. Let $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$. Now $S = \mathbb{N}_0 \times \dots \times \mathbb{N}_0 = \mathbb{N}_0^k$. Then, by the product lemma,

$$\Phi_S(x) = (\Phi_{\mathbb{N}_0}(x))^k.$$

Now, $\mathbb{N}_0 = \{0, 1, 2, 3, \dots\}$, so there exists 1 way to get each $n \in \mathbb{N}_0$:

$$\Phi_{\mathbb{N}_0}(x) = 1 + x + x^2 + \dots = \frac{1}{1-x}.$$

Therefore,

$$\Phi_S(x) = (\Phi_{\mathbb{N}_0}(x))^k = \left(\frac{1}{1-x}\right)^k. \quad \square$$

2.5.3 Negative Binomial Theorem

Exercise. For any positive integer k , show that

$$(1-x)^{-k} = \sum_{n \geq 0} \binom{n+k-1}{k-1} x^n$$

Solution. Let S_n be the set of all k -tuples $(a_1, \dots, a_k) \in \mathbb{N}_0^k$ with $a_1 + \dots + a_k = n$. By **2.5.2**, there exists $[x^n](1-x)^{-k}$ number of k -tuples satisfying this property.

Let T_n denote the set of all binary strings of length $n+k-1$ with $k-1$ 1's. Then we know

$$|T_n| = \binom{n+k-1}{k-1}.$$

We will establish a bijection between S_n and T_n , and hence prove that

$$[x^n](1-x)^{-k} = |S_n| = |T_n| = \binom{n+k-1}{k-1}.$$

Define $f: S_n \rightarrow T_n$ as follows: For a k -tuple $A = (a_1, \dots, a_k) \in S_n$,

$$f(A) = 0^{a_1} 1 0^{a_2} 1 \dots 1 0^{a_k}$$

where 0^{a_i} represents a_i 0's in a row. Notice that $f(A)$ is a string with exactly $k - 1$ 1's, and the length of the string $a_1 + \cdots + a_k + (k - 1) = n + k - 1$. Therefore $f(A) \in T_n$. It is trivial to show there exists an inverse of f which maps a binary string of length n with exactly $k - 1$ 1's to a k -tuple. This shows that f is a bijection between S_n to T_n . \square

2.5.4 Coin Value

Exercise. Let $n \in \mathbb{N}_0$. Suppose we have an unlimited supply of Canadian nickels, dimes, quarters, loonies, and toonies. How many ways can we make n cents using these coins?

Solution. Let $C_k = \{0, k, 2k, 3k, \dots\}$. Each collection of coins can be represented by a 5-tuple $(n, d, q, l, t) \in C_5 \times C_{10} \times C_{25} \times C_{100} \times C_{200}$. Let $S = C_5 \times C_{10} \times C_{25} \times C_{100} \times C_{200}$. We define the weight of any 5-tuple in S to be $w(n, d, q, l, t) = n + d + q + l + t$, which represents the value of this collection of coins in cents.

Using the weight function $\alpha_k(a) = a$ for the set C_k , we see that the generating series for C_k is

$$\Phi_{C_k}(x) = 1 + x^k + x^{2k} + x^{3k} + \cdots = \frac{1}{1 - x^k}.$$

For our set S , the set function $w(n, d, q, l, t) = \alpha_5(n) + \alpha_{10}(d) + \alpha_{25}(q) + \alpha_{100}(l) + \alpha_{200}(t)$. Applying the product lemma, we get the generating series for S :

$$\begin{aligned} \Phi_S(x) &= \Phi_{C_5}(x) \cdot \Phi_{C_{10}}(x) \cdot \Phi_{C_{25}}(x) \cdot \Phi_{C_{100}}(x) \cdot \Phi_{C_{200}}(x) \\ &= \frac{1}{(1 - x^5)(1 - x^{10})(1 - x^{25})(1 - x^{100})(1 - x^{200})}. \end{aligned}$$

The number of collections that are valued at n cents is then $[x^n]\Phi_S(x)$. \square

1. *Any particular reason for the use of rational number?* 