

# Chapter 1

Logan | Nick

## 1.1 Populations, Samples, and Processes

Statistics provides methods for organizing and summarizing data and for drawing conclusions from that data

### Def

Data : a collection of facts

### Def

Population : A well defined collection of objects for which we wish to obtain info

### Def

Census : When desired info is obtained from every member of the population

- problems : Time, money, practical

### Def

Sample : A subset of the population

### 1.

You want the home price in Edwardsville

- Fewer well trained appraisers gives better results than many poorly trained

## 2. Tree Age Study

Testing is destructive, so a sample is better

## Def

variable : any characteristic whose value may differ from one subject to another.

- denote with low letters

## Note

- Don't say \$McDonald's = 10\$
- Do say \$x = \$ the length of the tibia bone in 10 year old boys.

## Def

univariate data : result from making observations of 1 variable

- these variable can be qualitative / quantitative

## Def

Bivariate data : when observations are made on each of 2 variables for each individual

- (weight.mpg) of cars

## Def

Multivariate data : observations made on many variables

- patient data

## Ex

Labor force, sample 60,000, find population + sample

- population = labor force, sample size = 60,000 households

## Branches of Stats

1. Descriptive Stats : data are collected and you wish to summarize and describe features of the data (graphs, numerical summaries)
2. Inferential stats : data is collected from a sample and used to draw a conclusion about the population
  - confidence intervals, hypothesis test, prediction, etc...

## Types of sampling

- Simple random sampling : random choice / draw of the hat sampling
- Systematic sampling : selecting every  $k^{th}$  member of the population

- Cluster sampling : divide population into groups, then select some of these groups @ random
- Stratified sampling : divide population into groups. Find subgroups of groups (strata) and then draw random sample in strata
- Convenience sampling : sampling in the most convenient way
  - best to avoid , but a good starter

## Notate

sample size :  $n$

- For a dataset with  $n$  observations on some variable  $x$ , the individual observations will be denoted as  $x_1, x_2, \dots, x_n$ .

## 1.2

### Stem and leaf plots

#### Ex

(54, 59, 35, 41, 46, 25, 47, 60, 54, 46, 49, 46, 41, 34, 22)


During these problems it helps to first organize the numebrs in the list first

```

2 | 2, 5
3 | 4, 5
4 | 1, 1, 6, 6, 6, 7, 9
5 | 4, 4, 9
6 | 0

```

## Dot plots



## Histograms

left blank for nick to fill in

## Skewed (Right and left)

add a dataset to show?

## Bell

add a dataset to show?

## Flat uniform

add a dataset to show?

## nonsymmetric

add a dataset to show?

## bimodal symmetric

add a dataset to show?

## 1.3

### Def

mean : numerical value of average

Notate

**Sample mean** :  $\bar{x}$

$$\bar{x} = \frac{x_1 + x_2 + \dots + x_n}{n} = \frac{\sum_{i=1}^n x_i}{n}$$

Notate

**Population mean** :  $\mu$

- avg of all values in the entire pop.

### Ex

2, 2, 5, 3, 8, 9, 2, 3, 1

$$\bar{x} = \frac{\sum_{i=1}^{10} x_i}{10} = 3.6$$

The mean is inappropriate in some cases b/c of outliers.

- this makes the mean a **nonresistant measure**

### Def

Median : middle value /avg of 2 middle values when sorted

Notate

Median :  $\tilde{x}$

- if  $n = \text{odd}$ , median is at  $\frac{n+1}{2}$
- if  $n = \text{even}$  median are b/n  $\frac{n}{2}$  &  $\frac{n+1}{2}$

Notate

Population Mean :  $\tilde{\mu}$

## 1.4 Measures of Variability

One way to describe a distribution is by using the standard deviation

## Quartiles

- $Q_1$  - lower quartile separates bottom 25%
- $Q_2$  - median middle 50%
- $Q_3$  - upper quartile separates upper 25%

## Ex

2, 2, 5, 1, 3, 8, 9, 2, 31

SORT

1, 1, 2, 2, 2

3, 3, 5, 8, 9

$$\tilde{x} = 2 + 3 = 2.5$$

## Five number summary

- Find min,  $Q_1$ , median,  $Q_3$ , max

Note : If median is found in list, use it in both top half and lower half.

Ex : 2 2 5 1 3 8 9 2 3 1 100

$$\bar{x} = \frac{36+100}{11} \approx 12.36$$

Sort to find median.  $\tilde{x} = 3$ .

## Mean vs. Median

- median is the equal parts point
- mean is the balance point

Notate

## Trimmed mean : $\bar{x}_{tr}$

- compromise b/n the mean & median
- to find it, remove top & bottom 10%, then calculate the mean