Chapter 1

Logan | Nick

1.1 Populations, Samples, and Processes

Statistics provides methods for organizing and summarizing data and for drawing conclusions from that data

• Def

Data: a collection of facts

• Def

Population: A well defined collection of objects for which we wish to obtain info

• Def

Census: When desired info is obtained from every member of the population

- problems: Time, money, practical
- Def

Sample: A subset of the population

1.

You want the home price in Edwardsville

- Fewer well trained appraisers gives better results than many poorly trained
- 2. Tree Age Study

Testing is destructive, so a sample is better

• Def

variable: any characteristic whose clue may differ from one subject to another.

• denote with low letters

Note

- Don't say \$McDonald's = 10\$
- Do say x = the length of the tibia bone in 10 year old boys.

• Def

univariate data: result from making observations of 1 variable

- these variable can be qualitative / quantitative
- Def

Bivariate data: when observations are made on each of 2 variables for each individual

- (weight.mpg) of cars
- Def

Multivariate data: observations made on many variables

- patient data
- Ex

Labor force, sample 60,000, find population + sample

• population = labor force, sample size = 60,000 households

Branches of Stats

- 1. Descriptive Stats: data are collected and you wish to summarize and describe features of the data (graphs, numerical summaries)
- 2. Inferential stats: data is collected from a sample and used to draw a conclusion about the population
 - confidence intervals, hypothesis test, prediction, etc...

Types of sampling

- Simple random sampling: random choice / draw of the hat sampling
- Systematic sampling: selecting every k^{th} member of the population
- Cluster sampling : divide population into groups, then select some of these groups @ random
- Stratified sampling: divide population into groups. Find subgroups of groups (strata) and then draw random sample in strata
- Convenience sampling: sampling in the most convenient way
 - best to avoid, but a good starter

Notate

sample size : n

• For a dataset with n observations on some variable x, the individual observations will be denoted as x_1, x_2, \ldots, x_n .

1.2

Stem and leaf plots


• Ex

(54, 59, 35, 41, 46, 25, 47, 60, 54, 46, 49, 46, 41, 34, 22)

During these problems it helps to first organize the numebrs in the list first

$$\begin{array}{c|c} 2 \mid 2,5 \\ 3 \mid 4,5 \\ 4 \mid 1,1,6,6,6,7,9 \\ 5 \mid 4,4,9 \\ 6 \mid 0 \end{array}$$

Dot plots



Histograms

left blank for nick to fill in

Skewed (Right and left)

add a dataset to show?

Bell

add a dataset to show?

Flat uniform

add a dataset to show?

nonsymmetric

add a dataset to show?

bimodal symmetric

add a dataset to show?

1.3

• Def

mean: numerical value of average

Notate

Sample mean : \overline{x}

$$\overline{x} = \frac{x_1 + x_2 + \ldots + x_n}{n} = \frac{\sum_{i=1}^{n} x_i}{n}$$

Notate

Population mean : μ

- avg of all values in the entire pop.
- Ex

2, 2, 5, 3, 8, 9, 2, 3, 1

$$\bar{x} = \frac{\sum_{i=1}^{10} x_i}{10} = 3.6$$

The mean is inappropriate in some cases b/c of outliers.

- this makes the mean a nonresistant measure
- Def

Median: middle value /avg of 2 middle values when sorted

Notate

 $\mathrm{Median}:\, \widetilde{x}$

- if n = odd, median is at $\frac{n+1}{2}$
- if n = even median are b/n $\frac{n}{2}$ & $\frac{n+1}{2}$

Notate

Population Mean : $\tilde{\mu}$

1.4 Measures of Variability

One way to describe a distribution is by using the standard deviation

Quartiles

- Q_1 lower quartile separates bottom 25%
- Q_2 median middle 50%
- Q_3 upper quartile separates upper 25%
- Ex

 $2,\,2,\,5,\,1,\,3,\,8,\,9,\,2,\,31$

SORT

$$1, 1, 2, 2, 2$$

 $3, 3, 5, 8, 9$

$$\tilde{x} = 2 + 3 = 2.5$$

Five number summary

• Find min, Q_1 , median, Q_3 , max

Note: If median is found in list, use it in both top half and lower half.

 $Ex: 2\ 2\ 5\ 1\ 3\ 8\ 9\ 2\ 3\ 1\ 100$

$$\overline{x} = \frac{36+100}{11} \approx 12.36$$

Sort to find median. $\tilde{x} = 3$.

Mean vs. Median

- \bullet median is the equal parts point
- mean is the balance point

Notate

Trimmed mean : \overline{x}_{tr}

- compromise b/n the mean & median
- to find it, remove top & bottom 10%, then calculate the mean