

# Gauge Theories: Electroweak

Lecturer: Richard Ball

January 22, 2019

## 1 Introduction

Imagine a world without electroweak:

- Still have electromagnetism (EM), massive hadrons, atoms, gravity etc.
- Parity (P) and charge conjugation (C) are still good symmetries
- Flavour is always conserved: everything lasts forever (and always existed)
- No neutrinos (would be non-interacting)

Neutrinos were first hypothesised by Pauli (1930) to explain the missing energy in  $\beta$ -decay. They were first observed in 1956 (Cowan-Reines). Parity violation was first directly observed in 1956/7 (Lee-Yang-Wu).

### 1.1 Chirality

Define the projection operators  $P_R \equiv \frac{1}{2}(1 + \gamma^5)$  and  $P_L \equiv \frac{1}{2}(1 - \gamma^5)$ . Recalling that  $(\gamma^5)^2 = 1$  and  $(\gamma^5)^\dagger = \gamma^5$ , we can deduce the following properties:

- $P_R^2 = P_R$
- $P_L^2 = P_L$
- $P_L P_R = P_R P_L = 0$
- $P_R + P_L = 1$
- $P_R - P_L = \gamma^5$ .

Any Dirac spinor can be split up into a right-handed and a left-handed component,  $\psi = \psi_R + \psi_L$ , using the projection operators to define  $\psi_R = P_R \psi$  and  $\psi_L = P_L \psi$ . Since  $\gamma^\mu P_L = P_R \gamma^\mu$ , it follows that  $\bar{\psi} P_R \equiv (\bar{\psi})_R = \bar{\psi}_L$  and  $\bar{\psi} P_L \equiv (\bar{\psi})_L = \bar{\psi}_R$ . So

$$\begin{aligned}\bar{\psi} \psi &= \bar{\psi}(\psi_R + \psi_L) = \bar{\psi}_L \psi_R + \bar{\psi}_R \psi_L \\ &\quad \text{and} \\ \bar{\psi} \gamma^\mu \psi &= \bar{\psi} \gamma^\mu (\psi_R + \psi_L) = \bar{\psi}_R \gamma^\mu \psi_R + \bar{\psi}_L \gamma^\mu \psi_L.\end{aligned}\tag{1}$$

The Dirac Lagrangian splits up like

$$\begin{aligned}\mathcal{L}_D &= \bar{\psi}(i\not{\partial} - m)\psi \\ &= \bar{\psi}_R i\not{\partial} \psi_R + \bar{\psi}_L i\not{\partial} \psi_L + m(\bar{\psi}_L \psi_R + \bar{\psi}_R \psi_L)\end{aligned}\tag{2}$$

so the mass term mixes  $\psi_R$  and  $\psi_L$ : if  $m \rightarrow 0$ ,  $\psi_R$  and  $\psi_L$  are independent. In this scenario they are both 2-component spinors obeying the Weyl equation  $i\not{\partial} \psi_{R/L} = 0$ .

## 1.2 Helicity

The spin operator can be expressed as  $\Sigma^i = \frac{i}{2}\epsilon^{ijk}[\gamma^j, \gamma^k] = \gamma^5 \gamma^0 \gamma^i$ . Then  $[P_L, \Sigma^i] = [P_R, \Sigma^i]$ : spin and chirality commute. Now consider *helicity*, defined

$$h \equiv \frac{2\underline{\Sigma} \cdot \underline{p}}{|\underline{p}|}.\tag{3}$$

$h$  has eigenvalues  $\pm 1$ , which follows from  $(\not{p} - m)u^\pm = 0 \implies hu^\pm = \pm u^\pm$ . But  $\not{p} = E\gamma^0 - \underline{\gamma} \cdot \underline{p}$  and  $h = \gamma^5 \gamma^0 (\underline{\gamma} \cdot \underline{p})/|\underline{p}| = \gamma^5 \gamma^0 (E\gamma^0 - \not{p})$ . So

$$\begin{aligned}\gamma^5(E - \gamma^0 \not{p})u^\pm &= \pm u^\pm \\ \implies (P_R - P_L)(E - \gamma^0 m)u^\pm &= \pm p(P_R + P_L)u^\pm \\ \text{and } (E \mp p)u_R^\pm &= m\gamma^0 u_L^\pm \\ (E \pm p)u_L^\pm &= m\gamma^0 u_R^\pm.\end{aligned}\tag{4}$$

Again, the mass term mixes R and L, but if  $m \rightarrow 0$ ,  $p = E + \mathcal{O}(\frac{m^2}{E})$  and  $2Eu_R^- = 2Eu_L^+ = 0$  so  $u_R^- = u_L^+ = 0$ , i.e.  $u_R$  has helicity +1 and  $u_L$  has helicity -1.

Note that when  $m = 0$ , helicity is Lorentz invariant (no rest frame). For  $m \neq 0$ ,

$$\begin{aligned}u_R^- &= \frac{m\gamma^0}{E+p}u_L^- \approx \frac{m}{2E}\gamma^0 u_L^- \\ \text{and } u_L^+ &\approx \frac{m}{2E}\gamma^0 u_R^+.\end{aligned}\tag{5}$$

You get similar expressions for negative energy solutions:

$$\begin{aligned}\text{for } m = 0: v_R^- &= v_L^+ = 0 \\ \text{for } m \neq 0: v_R^- &= -\frac{m\gamma^0}{2E}v_L^- \text{ and } v_L^+ = -\frac{m\gamma^0}{2E}v_R^-.\end{aligned}\tag{6}$$

## 1.3 The Chiral Representation

In the chiral representation the gamma matrices can be expressed as:

$$\gamma^0 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \quad \gamma^i = \begin{pmatrix} 0 & -\sigma^i \\ \sigma^i & 0 \end{pmatrix} \quad \gamma^5 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

so

$$P_R = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \quad P_L = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \quad \implies \psi = \begin{pmatrix} \psi_R \\ \psi_L \end{pmatrix}.$$

Furthermore, the spin operator can be written

$$\Sigma^i = \gamma^5 \gamma^0 \gamma^i = \begin{pmatrix} \sigma^i & 0 \\ 0 & \sigma^i \end{pmatrix}$$

so the helicity eigenstates are  $\begin{pmatrix} \psi_R^\pm \\ 0 \end{pmatrix}$  and  $\begin{pmatrix} 0 \\ \psi_L^\pm \end{pmatrix}$ . Now consider the positive and negative energy solutions.  $(\not{p} - m)u = 0 \implies$

$$\begin{pmatrix} -m & E + \underline{\sigma} \cdot \underline{p} \\ E - \underline{\sigma} \cdot \underline{p} & -m \end{pmatrix} \begin{pmatrix} u_R \\ u_L \end{pmatrix} = 0.$$

But  $\underline{\sigma} \cdot \underline{p} u_{L/R} = \pm p u_{L/R}^\pm$  so  $(E \pm p)u_L^\pm = m u_R^\pm$ . Using  $E^2 = p^2 + m^2$ :

$$\begin{aligned} (E \pm p)u_L^\pm &= \sqrt{(E+p)(E-p)}u_R^\pm \\ \implies \sqrt{E \pm p} u_L^\pm &= \sqrt{E \mp p} u_R^\pm \end{aligned} \tag{7}$$

and we can write

$$u^\pm = \begin{pmatrix} \sqrt{E \pm p} \xi^\pm \\ \sqrt{E \mp p} \xi^\pm \end{pmatrix}$$

where  $\xi^+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  and  $\xi^- = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$ .

*Exercise: check that the normalisations  $\bar{u}u = 2m$ ,  $u^\dagger u = 2E$ .*

Similarly,  $(\not{p} + m)v = 0$  and  $(\underline{\sigma} \cdot \underline{p})v^\pm = \mp p v^\pm$ , so

$$\begin{aligned} (E \mp p)v_L^\pm &= -m v_R^\pm \\ \implies \sqrt{E \mp p} v_L^\pm &= -\sqrt{E \pm p} v_R^\pm \end{aligned} \tag{8}$$

and

$$v^\pm = \begin{pmatrix} \sqrt{E \mp p} \xi^\pm \\ -\sqrt{E \pm p} \xi^\pm \end{pmatrix}.$$

You can relate  $\xi^\pm$  to  $\chi^\pm$  by charge conjugation:  $v^\pm = i\gamma^2 u^{\pm*}$ , where

$$i\gamma^2 = \begin{pmatrix} 0 & -i\sigma^2 \\ i\sigma^2 & 0 \end{pmatrix}; \quad -i\sigma^2 = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$$

so  $\chi^\pm = -i\sigma^2 \xi^\pm$  and if  $\xi^\pm = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix}$  then  $\chi^\pm = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \begin{pmatrix} -1 \\ 0 \end{pmatrix}$  as before.

## 1.4 Parity

Under P,  $\psi \rightarrow \psi_p = \gamma^0 \psi$ . We know that  $P_L \gamma^0 = \gamma^0 P_R$ , so  $\psi_L \rightarrow \gamma^0 \psi_L = P_R \gamma^0 \psi_L = (\psi_p)_R$ . In other words,  $(\psi_L)_p = (\psi_p)_R$ , so parity switches L and R.

Note that  $[\gamma^0, \Sigma^i] = 0$  but under P  $\underline{p} \rightarrow -\underline{p}$  so  $h \rightarrow -h$ , as expected. This means  $u_R^\pm \rightarrow u_L^\mp$  etc.: parity flips helicity but not spin.

## 1.5 Charge conjugation

Under C,  $\psi \rightarrow \psi_c = C\bar{\psi}^T$  where  $C = i\gamma^2\gamma^0$ . Then  $P_L C = C P_L$  so  $\psi_L \rightarrow C(\bar{\psi})_L^T$ . This means that charge conjugation leaves the chirality unchanged. Helicity is also unchanged: C just takes particles  $\leftrightarrow$  antiparticles.

## 1.6 Time reversal

Under T,  $\psi \rightarrow \psi_T = B\psi$  where  $B = i\gamma^1\gamma^3 = -i\gamma^5 C$  and  $B^\dagger = B = B^{-1}$ . Again,  $P_L B = B P_L$  so  $\psi_L \rightarrow B\psi_L$  and time reversal leaves chirality and helicity unchanged (it reverses both spin and momentum).

# 2 Charged Current Electroweak

## 2.1 Fermi Theory (1934)

Fermi theory is based on a point-like 4-fermion interaction

$$G_F(\bar{n}\gamma_\mu p)(\bar{\nu}\gamma^\mu e) \quad (9)$$

representing the process  $n \rightarrow p e^- \bar{\nu}$ , where  $n$  is a neutron,  $p$  is a proton,  $e^-$  is an electron and  $\bar{\nu}$  is an antineutrino. More generally, the interaction can be written as  $G_F J_\mu^\dagger J^\mu$  where  $J_\mu$  is the "weak current" and is composed of leptonic and hadronic contributions:  $J_\mu = \bar{\nu}\gamma_\mu e + \bar{p}\gamma_\mu n + \dots$ . Note that  $J_\mu$  has  $\Delta Q = +1$  so  $J_\mu^\dagger$  has  $\Delta Q = -1$  and electric charge is conserved.

Considering mass dimensions  $[\cdot]$ :  $[m\psi\bar{\psi}] = +\psi$ ,  $[\psi] = 3/2$  and  $[J_\mu] = 3$  so  $[G_F] = -2$ . In other words, the coupling scales as  $1/\text{mass}^2$ . The mass scale is  $\approx 300$  GeV, so the interaction is weak. The original Fermi interaction conserves parity (and C and CP), just like QED. This turned out to be wrong, according to theory developed by Lee and Yang (1956) and the Wu experiment in 1957. In weak interactions, P and C are violated by CP is conserved. More on this later.

## 2.2 V-A Theory

Developed by Masshart, Sudarshan, Feynman, Gell-Man ... in 1958. V-A theory is based on a vector current  $V_\mu$  and an axial current  $A_\mu$ ,

$$\begin{aligned} V_\mu &= \bar{\nu}\gamma_\mu e + \bar{p}\gamma_\mu n + \dots \\ A_\mu &= \bar{\nu}\gamma_\mu\gamma^5 e + \bar{p}\gamma_\mu\gamma^5 n + \dots \end{aligned} \quad (10)$$

whose difference gives the overall current

$$\begin{aligned} \frac{1}{2}J_\mu &= \frac{1}{2}(V_\mu - A_\mu) = \bar{\nu}\gamma_\mu\frac{1}{2}(1 - \gamma^5)e + \bar{p}\gamma_\mu\frac{1}{2}(1 - \gamma^5)n + \dots \\ &= \bar{\nu}_L\gamma_\mu e_L + \bar{p}_L\gamma_\mu n_L + \dots \end{aligned} \quad (11)$$

So the weak interactions involve **only left-handed fields**, and maximally violate P and C. In fact, under P, C and CP the currents transform in the following ways:

$$\begin{array}{lll}
P & V^\mu \rightarrow V_\mu, & A^\mu \rightarrow -A_\mu, & (V - A)^\mu \rightarrow (V + A)_\mu \\
C & V^\mu \rightarrow -V^\mu, & A^\mu \rightarrow A^\mu, & (V - A)^\mu \rightarrow (-V - A)^\mu \\
CP & V^\mu \rightarrow -V_\mu, & A^\mu \rightarrow -A_\mu, & (V - A)^\mu \rightarrow -(V - A)_\mu
\end{array} \tag{12}$$

so  $(V - A)^{\mu\dagger}(V - A)_\mu$  is invariant under CP (and T).

Note that that neutrinos **only** interact weakly, so  $\nu_R$  does not interact at all! If neutrinos are massless, there is no need for  $\nu_R$ . From 1930-1998 neutrinos were always assumed to be massless: here we will assume  $m_\nu = 0$  so  $\nu$  are always left handed and  $\bar{\nu}$  are always right handed. In reality  $m_\nu \leq 0.3$  eV, which is very small.

So we have

$$\mathcal{L}_{4F} = -\frac{G_F}{\sqrt{2}} J_\mu^\dagger J^\mu \tag{13}$$

where the  $\sqrt{2}$  is historical.  $J_\mu = J_\mu^l + J_\mu^h$  can be split up into a leptonic and a hadronic current, each of which comprises three generations. For example, the leptonic current

$$\frac{1}{2} J_\mu^l = \bar{\nu}_e \gamma_\mu e_L + \bar{n} u_{(\mu)} \gamma_\mu \mu_L + \bar{\nu}_\tau \gamma_\mu \tau_L. \tag{14}$$

Lepton number  $L_e = N_{e^-} - N_{e^+} - N_{\nu_e} + N_{\bar{\nu}_e}$  and the corresponding  $L_\mu$  and  $L_\tau$  are conserved, according to Noether's Theorem.

The hadronic current

$$\frac{1}{2} J_\mu^h = \bar{u}_L \gamma_\mu d'_L + \bar{c}_L \gamma_\mu s'_L + \bar{t}_L \gamma_\mu b'_L \tag{15}$$

is simplest to express in terms of the quarks. The baryon number  $B = \sum_{gen} (N_u - N_{\bar{u}} - N_d + N_{\bar{d}})$  is conserved. Things are complicated due to the effects of quark mixing - see later.

(V-A) Theory is quite complicated, but there is only one coupling,  $G_F$ : we call this universality. Explicitly,

$$\mathcal{L}_{4F} = -\frac{G_F}{\sqrt{2}} (J_\mu^{l\dagger} J^{l\mu} + (J_\mu^{h\dagger} J^{l\mu} + J_\mu^{l\dagger} J^{h\mu}) + J_\mu^{h\dagger} J^{h\mu}) \tag{16}$$

so there are three types of (charged current) weak interaction

- **leptonic:** only leptons (and neutrinos) e.g.  $\mu \rightarrow e \nu_\mu \bar{\nu}_e$
- **semi-leptonic:** both leptons and quarks e.g.  $\pi^- = d\bar{u} \rightarrow \mu^- \bar{\nu}_\mu$
- **hadronic:** only quarks e.g.  $\Lambda \rightarrow p \pi$

But all of them violate P and C and conserve CP, only involve left-handed particles and only involve one coupling  $G_F$ .

**Example:**  $\pi^+ \approx u\bar{d} \rightarrow \mu^+ \nu_\mu$ .

