Measure Theory

Kasper Rosenkrands

MATØK7 Fall 2020

Contents

1	Betingede forventningsværdier	2
2	Processer med uafhængige of stationære tilvækst, specielt standard brownske bevægelser	3
3	Martingaler og kvadratisk variation	4
4	Itô-formlen	5
5	Girsanov-transformationen	6

Notation

Measure with densities

Theorem 0.1 (Measures with densities). Let $(\Omega, \mathcal{A}, \mu)$ be a measure space and let $f: \Omega \to [0, \infty]$ be measureable. Then

$$(f \odot \mu)(A) := \int_A f \, d\mu = \int_\Omega \chi_A f \, d\mu, \quad A \in \mathcal{A},$$

defines a new measure $f \odot \mu : (A) \rightarrow [0, \infty]$, called the measure with density f with respect to μ . For every $N \in \mathcal{A}$, the following implication holds,

$$\mu(N) = 0 \implies (f \odot \mu)(N) = 0.$$

Product σ -algrebras

Definition 0.1. Let $n \in \mathbb{N}$, $n \ge 2$, and suppose that, for every $i \in \{1, ..., n\}$, we are given a measurable space $(\Omega_i, \mathcal{A}_i)$.

• The smallest σ -algebra on $\times_{i=1}^n \Omega_i$ containing

$$\mathcal{A}_1 * \cdots * \mathcal{A}_n := \{A_1 \times \cdots \times A_n \mid A_1 \in \mathcal{A}_1, \dots, A_n \in \mathcal{A}_n\} \subset \mathcal{P}\left(\underset{i=1}{\overset{n}{\times}} \Omega_i \right)$$

is called the product σ -algebra defined by measns of A_1, \ldots, A_n . It is denoted by

$$\bigotimes_{i=1}^n \mathcal{A}_i := \sigma \left(\mathcal{A}_1 * \cdots * \mathcal{A}_n \right).$$

• Let Γ be a set and let $f_i: \Gamma \to \Omega_i$ be an arbitrary map for every $i \in \{1, \ldots, n\}$. Then the smallest σ -algebra on Γ turning all maps f_1, \ldots, f_n into measurable maps, i.e.,

$$\sigma(f_1,\ldots,f_n) := \sigma\left(\left\{f_i^{-1}(A_i) \mid A_i \in \mathcal{A}_i, i \in 1,\ldots,n\right\}\right),\,$$

is called the initial σ -algebra generated by f_1, \ldots, f_n .

1 Betingede forventningsværdier

2 Processer med uafhængige of stationære tilvækst, specielt standard brownske bevægelser

3 Martingaler og kvadratisk variation

relevante dele til forelæsning 9

- afsnit 5.8.1
 - def 5.101
 - eks 5.103
 - def 5.104
 - $-~{\rm eks}~5.105$
 - $\ \mathrm{sæt} \ 5.110$
- \bullet afsnit 5.8.2
 - def 5.112
 - bem 5.114
 - $\ \mathrm{bem} \ 5.115$
 - sæt 5.118
 - sæt 5.120
- afsnit 5.8.3
 - def 5.122
 - $-\ \mathrm{bem}\ 5.123$
 - sæt 5.125
 - -sæt (med def) 5.126

4 Itô-formlen

5 Girsanov-transformationen