## P 8 Product Stability

### P.8.1 Stability Summary and Conclusion

The stability studies for Solifenacin Succinate Tablets BP 5 mg were carried out at following conditions.

| **Protocol** | **Description** |
|---|---|
| **Name of product** | Solifenacin Succinate Tablets BP 5 mg |
| **Container closure system** | ALU/ALU Blister pack of 10 tablets. |
| **Mother Batch Size** | 1,20,000 Tablets |
| **Stability conditions** | Accelerated conditions (40°C ± 2°C / 75% RH ± 5% RH) |
| **Frequency** | Initial, 1,2,3 and 6 Months |
| **Stability conditions** | Long term (30°C ± 2°C / 75% RH ± 5% RH) |
| **Frequency** | Initial, 3 Months, 6 Months, 9 Months, 12 Months, 18 Months, 24 months, 36 Months |
| **Storage** | Store below 30° C, in dark & dry place. |

Following are the stability indicating tests and specifications that are carried out on the samples of Solifenacin Succinate Tablets BP 5 mg kept for long term and accelerated stability studies.

| **Sr. No.** | **Ingredients** | **Specifications** |
|---|---|---|
| 1. | Description | Light yellow coloured, round shaped, film coated tablet, plain on both sides. Such 10 tablets packed in ALU/ALU Blister. |
| 2. | Dissolution | NLT 75% (Q) |
| 3. | Disintegration Time | NMT 30 min. |
| 4. | Hardness | NLT 3 kp |
| 5. | Assay | 95.0% to 105.0% of the stated amount. |
| 6. | Related Substances By HPLC | |
| | Impurity I | NMT 1.5% |
| | Impurity 1 | NMT 0.5% |
| | Unknown Impurity | NMT 0.2% |
| | Total Unknown Impurities | NMT 1.0% |
| 7. | Microbial Limit Test | |
| | Total Aerobic Microbial Count (TAMC) | NMT 1000 cfu/gm |
| | Total Yeasts and Moulds Count (TYMC) | NMT 100 cfu/gm |
| | Pathogens:<br>a) Escherichia coli<br>b) Salmonella<br>c) S. aureus<br>d) P. aeruginosa | absent/gm<br>absent/10gm<br>absent/gm<br>absent/gm |