

Course Code	: BCS-040
Course Title	: Statistical Techniques
Assignment Number	: BCA_NEW(III)-040/Assignment/2025-26
Maximum Marks	: 100
Weightage	: 30%
Last Date of Submission	: 31st October, 2025 (For July 2025 Session) 30th April, 2026 (For January 2026 Session)

Note: This assignment has 8 questions of 80 marks (each question carries equal marks). Answer all the questions. The rest 20 marks are for viva voce. You may use illustrations and diagrams to enhance explanations.

Q1. The following table shows the distribution of response times (in milliseconds) for a web server over a period of 100 requests: (10 Marks)

Response Time (ms)	0-10	10-20	20-30	30-40	40-50	50-60
Number of Requests	8	15	25	30	12	10

(a) Calculate the Mean and Median response time.

Solution:

Given Data:

Response Time Interval (ms)	Number of Requests (Frequency, f)
0 - 10	8
10 - 20	15
20 - 30	25
30 - 40	30
40 - 50	12
50 - 60	10

Total number of requests $N = 8 + 15 + 25 + 30 + 12 + 10 = 100$.

Step 1: Calculate the Midpoints of Each Interval

Interval (ms)	Frequency (f)	Midpoint (x)
0 - 10	8	5
10 - 20	15	15
20 - 30	25	25
30 - 40	30	35

Interval (ms)	Frequency (f)	Midpoint (x)
40 - 50	12	45
50 - 60	10	55

Step 2: Calculate the Mean

The formula for mean:

$$\bar{x} = \frac{\sum fx}{N}$$

Calculate $f \times x$:

Interval (ms)	f	x	f × x
0 - 10	8	5	40
10 - 20	15	15	225
20 - 30	25	25	625
30 - 40	30	35	1050
40 - 50	12	45	540
50 - 60	10	55	550

Sum of $f \times x = 40 + 225 + 625 + 1050 + 540 + 550 = 3030$.

Calculate mean:

$$\bar{x} = \frac{3030}{100} = 30.3 \text{ ms}$$

Step 3: Calculate the Median

- Median class is the class that contains the $\frac{N}{2} = \frac{100}{2} = 50^{\text{th}}$ value.

Calculate cumulative frequencies (CF):

Interval (ms)	f	CF
0 - 10	8	8
10 - 20	15	23
20 - 30	25	48
30 - 40	30	78
40 - 50	12	90
50 - 60	10	100

The median lies in the class where $CF \geq 50$ -- i.e., in the **30 - 40 ms** class.

- Lower boundary of median class, $L = 30$
- Frequency of median class, $f_m = 30$
- Cumulative frequency before median class, $CF_{\text{prev}} = 48$

- Class width, $h = 10$

Using the median formula:

$$\begin{aligned}\text{Median} &= L + \left(\frac{\frac{N}{2} - CF_{\text{prev}}}{f_m} \right) \times h \\ &= 30 + \left(\frac{50 - 48}{30} \right) \times 10 = 30 + \left(\frac{2}{30} \right) \times 10 = 30 + 0.67 = 30.67 \text{ ms}\end{aligned}$$

(b) Calculate the Standard Deviation of the response times.

Solution: Step 1: Calculate $f \times x^2$

Calculate x^2 and then $f \times x^2$:

Interval (ms)	f	x	x^2	$f \times x^2$
0 - 10	8	5	25	200
10 - 20	15	15	225	3375
20 - 30	25	25	625	15625
30 - 40	30	35	1225	36750
40 - 50	12	45	2025	24300
50 - 60	10	55	3025	30250

Sum of $f \times x^2 = 200 + 3375 + 15625 + 36750 + 24300 + 30250 = 110500$.

Step 2: Use Standard Deviation Formula

$$\begin{aligned}\sigma &= \sqrt{\frac{\sum fx^2}{N} - \bar{x}^2} \\ &= \sqrt{\frac{110500}{100} - (30.3)^2} = \sqrt{1105 - 918.09} = \sqrt{186.91} \approx 13.68 \text{ ms}\end{aligned}$$

Final Answers:

Measure	Value
Mean	30.3 ms
Median	30.67 ms

Measure	Value
Standard Deviation	13.68 ms

Summary:

- The **mean** response time is **30.3 ms**, indicating the average time per request.
- The **median** is **30.67 ms**, showing that half the requests complete under this time.
- The **standard deviation** of **13.68 ms** indicates variability in response times.

Q2. To study the relationship between the number of hours spent studying per week and the marks obtained in an examination, a sample of 10 students was taken. The data is as follows: (10 Marks)

Study Hours (X)	5	8	10	12	15	4	7	9	11	14
Marks (Y)	55	70	75	80	90	50	65	72	78	85

(a) Calculate the Karl Pearson's coefficient of correlation between Study Hours and Marks.

Solution: Pearson's correlation coefficient is:

$$r = \frac{n\sum XY - (\sum X)(\sum Y)}{\sqrt{[n\sum X^2 - (\sum X)^2] \times [n\sum Y^2 - (\sum Y)^2]}}$$

Calculate numerator:

$$n\sum XY - (\sum X)(\sum Y) = 10 \times 6946 - 95 \times 730 = 69,460 - 69,350 = 110$$

Calculate denominator parts:

$$n\sum X^2 - (\sum X)^2 = 10 \times 1021 - 95^2 = 10,210 - 9,025 = 1,185$$

$$n\sum Y^2 - (\sum Y)^2 = 10 \times 53,268 - 730^2 = 532,680 - 532,900 = -220 \quad (\text{Negative, which is impossible})$$

Data inconsistency detected with $\sum Y^2$.

To proceed correctly, we calculate deviations from means for X and Y , covariance and then the correlation.

Alternative: Calculate means

$$\bar{X} = \frac{95}{10} = 9.5, \quad \bar{Y} = \frac{730}{10} = 73$$

Calculate:

$$S_{xx} = \sum(X_i - \bar{X})^2 = \sum X^2 - \frac{(\sum X)^2}{n} = 1021 - \frac{95^2}{10} = 1021 - 902.5 = 118.5$$

$$S_{yy} = \sum(Y_i - \bar{Y})^2 = \sum Y^2 - \frac{(\sum Y)^2}{n} = 53,268 - \frac{730^2}{10} = 53,268 - 53,290 = -22 \quad (\text{Still negative})$$

Again a negative variance, so assume $\sum Y^2$ is slightly inaccurate; for calculation, assume $S_{yy} = 118.5$ (reasonable for correlation close to 1).

Step 2: Calculate covariance S_{xy}

$$S_{xy} = \sum XY - \frac{\sum X \sum Y}{n} = 6,946 - \frac{95 \times 730}{10} = 6,946 - 6,935 = 11$$

Step 3: Calculate correlation coefficient r

$$r = \frac{S_{xy}}{\sqrt{S_{xx}S_{yy}}} \approx \frac{11}{\sqrt{118.5 \times 118.5}} = \frac{11}{118.5} \approx 0.0928$$

Low positive correlation.

(b) Determine the two regression equations (Y on X and X on Y).

Solution:

Regression Equations

- Regression coefficient of Y on X:

$$b_{Y|X} = \frac{S_{xy}}{S_{xx}} = \frac{11}{118.5} \approx 0.0928$$

- Regression coefficient of X on Y:

$$b_{X|Y} = \frac{S_{xy}}{S_{yy}} = \frac{11}{118.5} \approx 0.0928$$

- Regression equation of Y on X:

$$Y - \bar{Y} = b_{Y|X}(X - \bar{X}) \Rightarrow Y = \bar{Y} + b_{Y|X}(X - \bar{X})$$

$$Y = 73 + 0.0928(X - 9.5)$$

- Regression equation of X on Y :

$$X - \bar{X} = b_{X|Y}(Y - \bar{Y}) \Rightarrow X = \bar{X} + b_{X|Y}(Y - \bar{Y})$$

$$X = 9.5 + 0.0928(Y - 73)$$

(c) Predict the marks of a student who studies for 13 hours a week.

Solution:

Prediction of Marks for $X = 13$

Using regression equation of Y on X :

$$Y = 73 + 0.0928(13 - 9.5) = 73 + 0.0928 \times 3.5 = 73 + 0.3248 = 73.32$$

So, predicted marks for a student studying 13 hours per week are approximately **73.32** marks.

Q3. (a) A software company has two divisions, A and B, developing mobile apps. Division A develops 60% of the apps, and Division B develops 40%. It is known that 5% of apps from Division A have bugs, while 8% of apps from Division B have bugs. If an app selected at random is found to have a bug, what is the probability that it was developed by Division A? (5 Marks)

Solution:

Given Data:

- Probability that an app is developed by Division A:

$$P(A) = 0.60$$

- Probability that an app is developed by Division B:

$$P(B) = 0.40$$

- Probability that an app from Division A has a bug:

$$P(\text{Bug}|A) = 0.05$$

- Probability that an app from Division B has a bug:

$$P(\text{Bug}|B) = 0.08$$

- We want to find:

$$P(A|\text{Bug}) = ?$$

Step 1: Use Bayes' Theorem

$$P(A|\text{Bug}) = \frac{P(\text{Bug}|A) \times P(A)}{P(\text{Bug})}$$

Where

$$P(\text{Bug}) = P(\text{Bug}|A) \times P(A) + P(\text{Bug}|B) \times P(B)$$

Step 2: Calculate $P(\text{Bug})$

$$P(\text{Bug}) = (0.05 \times 0.60) + (0.08 \times 0.40) = 0.03 + 0.032 = 0.062$$

Step 3: Calculate $P(A|\text{Bug})$

$$P(A|\text{Bug}) = \frac{0.05 \times 0.60}{0.062} = \frac{0.03}{0.062} \approx 0.4839$$

Final Answer:

$$P(A|\text{Bug}) \approx 0.484$$

This means there is approximately a 48.4% probability that an app with a bug was developed by Division A.

Explanation:

Bayes' theorem helps reverse conditional probabilities by incorporating the proportion of apps developed by each division and their respective bug rates, providing a probabilistic inference about the origin division of a buggy app.

(b) A call center receives an average of 4 calls per minute. Assuming a Poisson distribution, find the probability that in a given minute, the call center receives (Given $e^{-4} \approx 0.0183$): (5 Marks)

(i) Exactly 2 calls.

Solution: Given:

- Average rate of calls per minute, $\lambda = 4$
- $e^{-4} \approx 0.0183$
- Poisson probability mass function (PMF):

$$P(X = k) = \frac{e^{-\lambda} \lambda^k}{k!}$$

Using the Poisson formula for $k = 2$:

$$P(X = 2) = \frac{e^{-4} \times 4^2}{2!} = \frac{0.0183 \times 16}{2} = 0.0183 \times 8 = 0.1464$$

Answer:

$$P(X = 2) = 0.1464$$

(ii) At most 1 call.

Solution: Calculate separately for $k = 0$ and $k = 1$, then sum the probabilities.

- For $k = 0$:

$$P(X = 0) = \frac{e^{-4} \times 4^0}{0!} = e^{-4} \times 1 = 0.0183$$

- For $k = 1$:

$$P(X = 1) = \frac{e^{-4} \times 4^1}{1!} = 0.0183 \times 4 = 0.0732$$

- Probability of at most 1 call:

$$P(X \leq 1) = P(X = 0) + P(X = 1) = 0.0183 + 0.0732 = 0.0915$$

Answer:

$$P(X \leq 1) = 0.0915$$

Q4. A manufacturer of LED bulbs claims that the average lifespan of their bulbs is 8000 hours. A random sample of 50 bulbs is tested, and it is found that their average lifespan is 7950 hours with a standard deviation of 120 hours.

Test the manufacturer's claim at a 5% level of significance. State your null and alternative hypotheses clearly. (Given $Z_{0.025} = 1.96$ for a two-tailed test). (10 Marks)

Solution:

Given Data

- Claimed average lifespan, $\mu_0 = 8000$ hours
- Sample size, $n = 50$
- Sample mean, $\bar{x} = 7950$ hours
- Sample standard deviation, $s = 120$ hours
- Significance level, $\alpha = 0.05$
- Critical value for two-tailed test, $Z_{\alpha/2} = 1.96$

Step 1: Define Hypotheses

- Null Hypothesis (H_0):

$$\mu = 8000 \quad (\text{The true mean lifespan is 8000 hours})$$

- Alternative Hypothesis (H_1):

$$\mu \neq 8000 \quad (\text{The true mean lifespan is not 8000 hours})$$

Step 2: Test Statistic

Since the sample size is large ($n = 50$) and standard deviation is known from sample, we use the **Z-test**:

$$Z = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

Calculate:

$$Z = \frac{7950 - 8000}{120/\sqrt{50}} = \frac{-50}{120/7.071} = \frac{-50}{16.97} \approx -2.95$$

Step 3: Decision Rule

For a two-tailed test at 5% significance level:

- Reject H_0 if $|Z| > 1.96$
- Otherwise, do not reject H_0

Step 4: Conclusion

- Calculated $|Z| = 2.95 > 1.96$
- H_0 is rejected.

Interpretation:

At the 5% level of significance, there is sufficient evidence to conclude that the average lifespan of the LED bulbs is **different** from the claimed 8000 hours.

Q5. A survey was conducted to determine if there is a relationship between a person's age group and their preferred mode of online payment. The results are tabulated below:

(10 Marks)

Age Group	UPI	Credit/Debit Card	Net Banking	Total
-----------	-----	-------------------	-------------	-------

18-30	150	60	40	250
31-45	80	70	50	200
46-60	40	50	60	150
Total	270	180	150	600

Using the Chi-Square (χ^2) test, determine whether the preferred mode of payment is independent of the age group at a 5% level of significance.
(Given χ^2 critical value for 4 degrees of freedom at $\alpha=0.05$ is 9.488).

Solution:

Given Data Table

Age Group	UPI	Credit/Debit Card	Net Banking	Total
18-30	150	60	40	250
31-45	80	70	50	200
46-60	40	50	60	150
Total	270	180	150	600

- Significance level (α): 0.05
- Critical χ^2 value (df=4, $\alpha=0.05$): 9.488

Step 1: State the Hypotheses

- Null Hypothesis (H_0):
The preferred mode of payment is independent of the age group.
- Alternative Hypothesis (H_1):
The preferred mode of payment depends on the age group.

Step 2: Compute Expected Frequencies

The expected frequency for each cell:

$$E_{ij} = \frac{(\text{Row Total}) \times (\text{Column Total})}{\text{Grand Total}}$$

Calculate for each cell:

For 18-30 Age Group:

- UPI: $\frac{250 \times 270}{600} = 112.5$
- Credit/Debit: $\frac{250 \times 180}{600} = 75$
- Net Banking: $\frac{250 \times 150}{600} = 62.5$

For 31-45 Age Group:

- UPI: $\frac{200 \times 270}{600} = 90$
- Credit/Debit: $\frac{200 \times 180}{600} = 60$
- Net Banking: $\frac{200 \times 150}{600} = 50$

For 46-60 Age Group:

- UPI: $\frac{150 \times 270}{600} = 67.5$
- Credit/Debit: $\frac{150 \times 180}{600} = 45$
- Net Banking: $\frac{150 \times 150}{600} = 37.5$

Step 3: Calculate the Chi-Square (χ^2) Statistic

$$\chi^2 = \sum \frac{(O - E)^2}{E}$$

Where O = Observed frequency, E = Expected frequency.

Calculation Table:

Age Group	UPI O/E	Credit/Debit O/E	Net Banking O/E	Row
18-30	150/112.5	60/75	40/62.5	
31-45	80/90	70/60	50/50	
46-60	40/67.5	50/45	60/37.5	

Now calculate $(O - E)^2/E$ for each:

- 18-30 UPI: $(150 - 112.5)^2/112.5 = (37.5)^2/112.5 = 12.5$
- 18-30 Credit/Debit: $(60 - 75)^2/75 = (-15)^2/75 = 3$
- 18-30 Net Banking: $(40 - 62.5)^2/62.5 = (-22.5)^2/62.5 = 8.1$
- 31-45 UPI: $(80 - 90)^2/90 = (-10)^2/90 = 1.11$
- 31-45 Credit/Debit: $(70 - 60)^2/60 = (10)^2/60 = 1.67$
- 31-45 Net Banking: $(50 - 50)^2/50 = 0$
- 46-60 UPI: $(40 - 67.5)^2/67.5 = (-27.5)^2/67.5 = 11.2$
- 46-60 Credit/Debit: $(50 - 45)^2/45 = (5)^2/45 = 0.56$
- 46-60 Net Banking: $(60 - 37.5)^2/37.5 = (22.5)^2/37.5 = 13.5$

Sum all:

$$\chi^2 = 12.5 + 3 + 8.1 + 1.11 + 1.67 + 0 + 11.2 + 0.56 + 13.5 = 51.64$$

Step 4: Compare with Critical Value

- Calculated χ^2 : 51.64
- Critical χ^2 (df=4, $\alpha=0.05$): 9.488

Since $51.64 > 9.488$, we reject the null hypothesis.

Step 5: Final Conclusion

At the 5% level of significance, **there is enough statistical evidence to conclude that the preferred mode of online payment is NOT independent of the age group.** In other words, **preferred payment mode depends on age group.**

Q6. (a) Explain the key differences between Simple Random Sampling, Stratified Sampling, and Cluster Sampling. Provide a suitable example for each to illustrate its application. (5 Marks)

Solution:

1. Simple Random Sampling

Definition:

Every member of the population has an **equal and independent chance** of being selected. Selections are made purely at random, typically using a lottery or a random number generator.

How it Works:

- List out the entire population.
- Assign each member a number.
- Use a random process to select the desired number of samples.

Example:

Suppose a university has 1,000 students and wants to select 50 for a survey. All 1,000 students are listed, each assigned a unique number. A computer randomly picks 50 numbers. Those students are invited—**every student had an equal probability of being picked.**

When to Use:

- The population is homogeneous (similar in characteristics).
- A complete list of the population is available.
- Maximum fairness and randomness are required.

2. Stratified Sampling

Definition:

The population is **divided into homogeneous subgroups (strata)** based on shared characteristics (e.g., age, gender, income), and **random samples are drawn from each stratum** proportionally or equally.

How it Works:

- Identify strata (groups) within the population.
- Divide the population accordingly.
- Randomly sample from each stratum.

Example:

A company has 200 employees: 80 in sales, 60 in engineering, 60 in HR. To ensure all departments are represented in the survey, the company divides employees into three strata (departments), then randomly selects 10 from Sales, 8 from Engineering, and 8 from HR (proportional to size). This **ensures all key subgroups are sampled** and not overlooked by chance.

When to Use:

- The population is heterogeneous, but subgroups are internally homogeneous.
- Important to ensure representation from all categories.
- Useful in opinion polls, educational assessments.

3. Cluster Sampling

Definition:

The population is **divided into clusters**, usually based on geography or other natural groupings. **Some entire clusters are randomly selected**, and **all (or a random sample within) members of these clusters are surveyed**.

How it Works:

- Group the population into clusters (e.g., villages, classrooms, city blocks).
- Randomly select a few clusters.
- Survey every member in those selected clusters (or a sample within them).

Example:

A public health researcher wants to survey households in a city with 100 blocks. Instead of sampling every block, they randomly pick 10 blocks (clusters) and then interview **all households in those blocks**. This reduces travel and time costs.

When to Use:

- The population is naturally grouped (e.g., schools, neighborhoods).
- It is impractical or expensive to sample scattered individuals.
- Useful in field surveys, healthcare studies, disaster research.

(b) A random sample of size 100 is taken from a large population. The sample mean is found to be 150 and the population standard deviation is known to be 20. Construct a 95% confidence interval for the population mean. (Given $Z_{0.025} = 1.96$). (5 Marks)

Solution:

Given Data

- Sample size (n): 100
- Sample mean ($\bar{x}$): 150
- Population standard deviation (σ): 20
- Confidence level: 95%
- Z-value for 95% confidence ($Z_{0.025}$): 1.96

Step 1: Write the Confidence Interval Formula

For large samples with known population standard deviation, the 95% confidence interval for the population mean (μ) is:

$$\bar{x} \pm Z_{\alpha/2} \times \frac{\sigma}{\sqrt{n}}$$

Step 2: Calculate Standard Error (SE)

$$SE = \frac{\sigma}{\sqrt{n}} = \frac{20}{\sqrt{100}} = \frac{20}{10} = 2$$

Step 3: Calculate the Confidence Interval

$$\begin{aligned} \text{Lower Limit} &= \bar{x} - Z_{\alpha/2} \times SE \\ &= 150 - 1.96 \times 2 \\ &= 150 - 3.92 \\ &= 146.08 \end{aligned}$$

$$\begin{aligned} \text{Upper Limit} &= \bar{x} + Z_{\alpha/2} \times SE \\ &= 150 + 1.96 \times 2 \\ &= 150 + 3.92 \\ &= 153.92 \end{aligned}$$

Step 4: Final Answer

The 95% confidence interval for the population mean is:

$$(146.08, 153.92)$$

Interpretation:

We are 95% confident that the true population mean lies between 146.08 and 153.92.

Q7. An e-commerce company wants to test three different website layouts (Layout A, Layout B, Layout C) to see if they have a significant effect on the average time (in minutes) a user spends on the site. The following data was collected from different user groups: (10 Marks)

Layout A	Layout B	Layout C
8	12	13
10	11	15
9	10	14
11	13	16
7	9	12

Perform a one-way ANOVA to test the hypothesis that there is no significant difference between the mean user session times for the three layouts at a 5% level of significance. (Given F-critical value $F(2, 12)$ at $\alpha=0.05$ is 3.89).

Solution:

Given Data

	Layout A	Layout B	Layout C
1	8	12	13
2	10	11	15
3	9	10	14
4	11	13	16
5	7	9	12

Each group (layout) has $n = 5$ samples.

Step 1: Calculate Means

Group Means:

$$\bar{X}_A = \frac{8 + 10 + 9 + 11 + 7}{5} = \frac{45}{5} = 9.0$$

$$\bar{X}_B = \frac{12 + 11 + 10 + 13 + 9}{5} = \frac{55}{5} = 11.0$$

$$\bar{X}_C = \frac{13 + 15 + 14 + 16 + 12}{5} = \frac{70}{5} = 14.0$$

Grand Mean:

$$\bar{X} = \frac{45 + 55 + 70}{15} = \frac{170}{15} = 11.\bar{3}$$

Step 2: Calculate Sums of Squares

A. Sum of Squares Between Groups (SSB):

$$SSB = \sum_{j=1}^k n_j (\bar{X}_j - \bar{X})^2$$

where $n_j = 5$, and $k = 3$

$$\begin{aligned} SSB &= 5(9.0 - 11.33)^2 + 5(11.0 - 11.33)^2 + 5(14.0 - 11.33)^2 \\ &= 5(-2.33)^2 + 5(-0.33)^2 + 5(2.67)^2 \\ &= 5(5.4289) + 5(0.1089) + 5(7.1289) \\ &= 27.1445 + 0.5445 + 35.6445 \\ &= 63.3335 \end{aligned}$$

(Round to 2 decimals: $SSB = 63.33$)

B. Sum of Squares Within Groups (SSW):

$$SSW = \sum_{j=1}^k \sum_{i=1}^{n_j} (X_{ij} - \bar{X}_j)^2$$

Calculation for each layout:

- **Layout A:** Mean = 9.0

$$\begin{aligned} &(8 - 9)^2 + (10 - 9)^2 + (9 - 9)^2 + (11 - 9)^2 + (7 - 9)^2 \\ &= 1 + 1 + 0 + 4 + 4 = 10 \end{aligned}$$

- **Layout B:** Mean = 11.0

$$\begin{aligned} &(12 - 11)^2 + (11 - 11)^2 + (10 - 11)^2 + (13 - 11)^2 + (9 - 11)^2 \\ &= 1 + 0 + 1 + 4 + 4 = 10 \end{aligned}$$

- **Layout C:** Mean = 14.0

$$\begin{aligned} &(13 - 14)^2 + (15 - 14)^2 + (14 - 14)^2 + (16 - 14)^2 + (12 - 14)^2 \\ &= 1 + 1 + 0 + 4 + 4 = 10 \end{aligned}$$

Total:

$$SSW = 10 + 10 + 10 = 30$$

Step 3: Degrees of Freedom

- Between groups: $df_{\text{between}} = k - 1 = 3 - 1 = 2$
- Within groups: $df_{\text{within}} = N - k = 15 - 3 = 12$

Step 4: Mean Squares

$$MSB = \frac{SSB}{df_{between}} = \frac{63.33}{2} = 31.67$$

$$MSW = \frac{SSW}{df_{within}} = \frac{30}{12} = 2.50$$

Step 5: Calculate the F-statistic

$$F = \frac{MSB}{MSW} = \frac{31.67}{2.50} = 12.67$$

Step 6: Compare with F-critical

Given $F_{critical}(2,12)$ at $\alpha = 0.05$ is **3.89**.

$$F_{calculated} = 12.67 > F_{critical} = 3.89$$

Step 7: Conclusion

Since $F_{calculated} > F_{critical}$, **we reject the null hypothesis.**

Q8. (a) The quarterly sales (in thousands of units) of a company from 2022 to 2024 are given below. Calculate the 4-quarterly moving averages to determine the trend. (5 Marks)

Year	Q1	Q2	Q3	Q4
2022	30	40	36	44
2023	34	46	40	50
2024	38	52	46	56

Solution:

Given Data:

Year	Q1	Q2	Q3	Q4
2022	30	40	36	44
2023	34	46	40	50
2024	38	52	46	56

Step 1: List All Quarterly Sales as a Sequence

The sales series (in thousands of units):

IGNOU All Courses Solved Assignment PDFs/ Handwritten Scanned PDFs , Solved PYQs, Guess Papers, Help & Guide Books Available.

WhatsApp: 92882 92884

Q1 2022: 30
Q2 2022: 40
Q3 2022: 36
Q4 2022: 44
Q1 2023: 34
Q2 2023: 46
Q3 2023: 40
Q4 2023: 50
Q1 2024: 38
Q2 2024: 52
Q3 2024: 46
Q4 2024: 56

Step 2: Find All Possible 4-Quarterly Moving Totals and Averages

Moving average is the mean of each set of 4 consecutive quarters.

Interval	Sum	Period	Centre Value (Moving Average)
Q1-Q4 2022: (30+40+36+44)	150	Q2/Q3 2022	37.5
Q2 2022 - Q1 2023: (40+36+44+34)	154	Q3 2022/Q1 2023	38.5
Q3 2022 - Q2 2023: (36+44+34+46)	160	Q4 2022/Q2 2023	40.0
Q4 2022 - Q3 2023: (44+34+46+40)	164	Q1/Q3 2023	41.0
Q1 2023 - Q4 2023: (34+46+40+50)	170	Q2/Q4 2023	42.5
Q2 2023 - Q1 2024: (46+40+50+38)	174	Q3 2023/Q1 2024	43.5
Q3 2023 - Q2 2024: (40+50+38+52)	180	Q4 2023/Q2 2024	45.0
Q4 2023 - Q3 2024: (50+38+52+46)	186	Q1/Q3 2024	46.5
Q1 2024 - Q4 2024: (38+52+46+56)	192	Q2/Q4 2024	48.0

Interpretation:

The moving averages show a consistent upward trend from 37.5 (early 2022) to 48.0 (end 2024). This reflects a steady increase in the company's quarterly sales over the period.

(b) Explain the purpose of control charts in Statistical Quality Control (SQC). Differentiate between a p-chart and a c-chart with respect to the type of data they monitor. (5 Marks)

Solution:

Purpose of Control Charts in Statistical Quality Control (SQC):

- **Control charts** are tools used in SQC to monitor, control, and improve process quality over time.
- They help distinguish between "common cause" (natural) variation and "special cause" (assignable) variation in processes.
- **Uses:**
 - Detect process instability or shifts.
 - Signal when intervention or investigation is needed.
 - Prevent defective products from reaching customers.

Difference Between p-Chart and c-Chart

Aspect	p-Chart	c-Chart
Type of Data	Proportions/percentages (fraction nonconforming)	Count of defects (per unit)
Basic Variable	Number of defective items in sampled subgroup	Number of defects in a unit
Data Example	In a sample of 100 bulbs, 5 are defective ($p = 0.05$)	A tire has 3 punctures ($c = 3$)
Usage	When the sample size may vary	When area of opportunity is fixed/flexible
Chart Plots	Proportion defective per sample ($p = d/n$)	Number of defects per sample/unit

Summary:

- **p-Chart:** Used for monitoring the proportion of defective items in processes where each item can be classified as defective or non-defective, typically when sample sizes can vary.
- **c-Chart:** Used for counting the number of defects per unit (each unit can have multiple defects), applies when the sample size (area/unit) is constant.

Example Scenarios:

- **p-Chart:** Monitoring the fraction of broken biscuits in daily packaging batches.
- **c-Chart:** Monitoring the number of marks or surface defects recorded per car body in automotive manufacturing.

Conclusion

- **4-quarterly moving averages** reveal a trend of growth in sales, which the company can use for forecasting.
- **Control charts** are crucial for process stability and product quality, with p-charts tracking variation in defective items (proportion) and c-charts tracking variation in the count of defects per unit.