P03

The neighbor-joining method: a new method for reconstructing phylogenetic trees

Authors	Naruya Saitou, M Nei

Published	Molecular Biology and Evolution (1987)
Paper Link	https://academic.oup.com/mbe/article-pdf/4/4/406/11167444/7sait.pdf
Domain	Biology & biomedical science

Abstract

A new method called the neighbor-joining method is proposed for reconstructing phylogenetic trees from evolutionary distance data. The principle of this method is to find pairs of operational taxonomic units (OTUs [= neighbors]) that minimize the total branch length at each stage of clustering of OTUs starting with a starlike tree. The branch lengths as well as the topology of a parsimonious tree can quickly be obtained by using this method. Using computer simulation, we studied the efficiency of this method in obtaining the correct unrooted tree in comparison with that of five other tree-making methods: the unweighted pair group method of analysis, Farris's method, Sattath and Tversky's method, Li's method, and Tateno et al.'s modified Farris method. The new, neighbor-joining method and Sattath and Tversky's method are shown to be generally better than the other methods.

Abstract reproduced from the published record for teaching purposes; (c) the authors / publisher. Source: https://doi.org/10.1093/oxfordjournals.molbev.a040454

Notes: record the page or section for every claim you make about this paper.

Reading prompts

1. What problem does the paper try to solve? Write it in one sentence, in your own words.
2. What is unknown or hidden, and what can actually be observed or measured?
3. What method do the authors use? Is it new, borrowed from another paper, or a modification of one?
4. What evidence do they give (proofs, simulations, experiments, real data)?
5. What assumptions or limitations can you find?
6. Which other papers in the corpus does it cite, use, extend or resemble? Note the page or section.