


Data Representation vs. Data Classification

- PCA finds the most accurate *data representation* in a lower dimensional space
- Project data in the directions of maximum variance
- However the directions of maximum variance may be useless for classification




- Fisher Linear Discriminant project to a line which preserves direction useful for *data classification*


Fisher Linear Discriminant

- **Main idea:** find projection to a line s.t. samples from different classes are well separated

Example in 2D




*bad line to project to,
classes are mixed up*



*good line to project to,
classes are well separated*

Fisher Linear Discriminant

- Suppose we have 2 classes and d -dimensional samples $\mathbf{x}_1, \dots, \mathbf{x}_n$ where
 - n_1 samples come from the first class
 - n_2 samples come from the second class
- consider projection on a line
- Let the line direction be given by unit vector \mathbf{v}



- Scalar $\mathbf{v}^t \mathbf{x}_i$ is the distance of projection of \mathbf{x}_i from the origin
- Thus it $\mathbf{v}^t \mathbf{x}_i$ is the projection of \mathbf{x}_i into a one dimensional subspace

Fisher Linear Discriminant


- Thus the projection of sample \mathbf{x}_i onto a line in direction \mathbf{v} is given by $\mathbf{v}^t \mathbf{x}_i$
- How to measure separation between projections of different classes?
- Let $\tilde{\mu}_1$ and $\tilde{\mu}_2$ be the means of projections of classes 1 and 2
- Let μ_1 and μ_2 be the means of classes 1 and 2
- $|\tilde{\mu}_1 - \tilde{\mu}_2|$ seems like a good measure

$$\tilde{\mu}_1 = \frac{1}{n_1} \sum_{\mathbf{x}_i \in C_1}^{n_1} \mathbf{v}^t \mathbf{x}_i = \mathbf{v}^t \left(\frac{1}{n_1} \sum_{\mathbf{x}_i \in C_1}^{n_1} \mathbf{x}_i \right) = \mathbf{v}^t \mu_1$$

similarly, $\tilde{\mu}_2 = \mathbf{v}^t \mu_2$

Fisher Linear Discriminant


- How good is $|\tilde{\mu}_1 - \tilde{\mu}_2|$ as a measure of separation?
 - The larger $|\tilde{\mu}_1 - \tilde{\mu}_2|$, the better is the expected separation



- the vertical axes is a better line than the horizontal axes to project to for class separability
- however $|\hat{\mu}_1 - \hat{\mu}_2| > |\tilde{\mu}_1 - \tilde{\mu}_2|$

Fisher Linear Discriminant


- The problem with $|\tilde{\mu}_1 - \tilde{\mu}_2|$ is that it does not consider the variance of the classes




Fisher Linear Discriminant

- We need to normalize $|\tilde{\mu}_1 - \tilde{\mu}_2|$ by a factor which is proportional to variance
- Have samples $\mathbf{z}_1, \dots, \mathbf{z}_n$. Sample mean is $\mu_z = \frac{1}{n} \sum_{i=1}^n \mathbf{z}_i$
- Define their **scatter** as
$$\mathbf{s} = \sum_{i=1}^n (\mathbf{z}_i - \mu_z)^2$$
- Thus scatter is just sample variance multiplied by n
 - scatter measures the same thing as variance, the spread of data around the mean
 - scatter is just on different scale than variance

larger scatter:



smaller scatter:



Fisher Linear Discriminant

- Fisher Solution: normalize $|\tilde{\mu}_1 - \tilde{\mu}_2|$ by scatter
- Let $\mathbf{y}_i = \mathbf{v}^t \mathbf{x}_i$, i.e. \mathbf{y}_i 's are the projected samples
- Scatter for projected samples of class 1 is

$$\tilde{\mathbf{s}}_1^2 = \sum_{\mathbf{y}_i \in \text{Class 1}} (\mathbf{y}_i - \tilde{\mu}_1)^2$$


- Scatter for projected samples of class 2 is

$$\tilde{\mathbf{s}}_2^2 = \sum_{\mathbf{y}_i \in \text{Class 2}} (\mathbf{y}_i - \tilde{\mu}_2)^2$$

Fisher Linear Discriminant

- We need to normalize by both scatter of class 1 and scatter of class 2
- Thus Fisher linear discriminant is to project on line in the direction \mathbf{v} which maximizes

want projected means are far from each other

$$J(\mathbf{v}) = \frac{\overbrace{(\tilde{\mu}_1 - \tilde{\mu}_2)^2}}{\tilde{\mathbf{s}}_1^2 + \tilde{\mathbf{s}}_2^2}$$


want scatter in class 1 is as small as possible, i.e. samples of class 1 cluster around the projected mean $\tilde{\mu}_1$


want scatter in class 2 is as small as possible, i.e. samples of class 2 cluster around the projected mean $\tilde{\mu}_2$

Fisher Linear Discriminant

$$J(\mathbf{v}) = \frac{(\tilde{\mu}_1 - \tilde{\mu}_2)^2}{\tilde{\mathbf{s}}_1^2 + \tilde{\mathbf{s}}_2^2}$$

- If we find \mathbf{v} which makes $J(\mathbf{v})$ large, we are guaranteed that the classes are well separated

projected means are far from each other



small $\tilde{\mathbf{s}}_1$ implies that projected samples of class 1 are clustered around projected mean

small $\tilde{\mathbf{s}}_2$ implies that projected samples of class 2 are clustered around projected mean

Fisher Linear Discriminant Derivation

$$\mathbf{J}(\mathbf{v}) = \frac{(\tilde{\mu}_1 - \tilde{\mu}_2)^2}{\tilde{\mathbf{S}}_1^2 + \tilde{\mathbf{S}}_2^2}$$

- All we need to do now is to express \mathbf{J} explicitly as a function of \mathbf{v} and maximize it
 - straightforward but need linear algebra and Calculus
- Define the separate class scatter matrices \mathbf{S}_1 and \mathbf{S}_2 for classes 1 and 2. These measure the scatter of original samples \mathbf{x}_i (before projection)

$$\mathbf{S}_1 = \sum_{\mathbf{x}_i \in \text{Class 1}} (\mathbf{x}_i - \mu_1)(\mathbf{x}_i - \mu_1)^t$$

$$\mathbf{S}_2 = \sum_{\mathbf{x}_i \in \text{Class 2}} (\mathbf{x}_i - \mu_2)(\mathbf{x}_i - \mu_2)^t$$

Fisher Linear Discriminant Derivation

- Now define the **within** the class scatter matrix

$$\mathbf{S}_W = \mathbf{S}_1 + \mathbf{S}_2$$

- Recall that $\tilde{\mathbf{s}}_1^2 = \sum_{y_i \in \text{Class } 1} (\mathbf{y}_i - \tilde{\mu}_1)^2$

- Using $\mathbf{y}_i = \mathbf{v}^t \mathbf{x}_i$ and $\tilde{\mu}_1 = \mathbf{v}^t \mu_1$

$$\begin{aligned}\tilde{\mathbf{s}}_1^2 &= \sum_{y_i \in \text{Class } 1} (\mathbf{v}^t \mathbf{x}_i - \mathbf{v}^t \mu_1)^2 \\&= \sum_{y_i \in \text{Class } 1} (\mathbf{v}^t (\mathbf{x}_i - \mu_1))^t (\mathbf{v}^t (\mathbf{x}_i - \mu_1)) \\&= \sum_{y_i \in \text{Class } 1} ((\mathbf{x}_i - \mu_1)^t \mathbf{v})^t ((\mathbf{x}_i - \mu_1)^t \mathbf{v}) \\&= \sum_{y_i \in \text{Class } 1} \mathbf{v}^t (\mathbf{x}_i - \mu_1) (\mathbf{x}_i - \mu_1)^t \mathbf{v} = \mathbf{v}^t \mathbf{S}_1 \mathbf{v}\end{aligned}$$

Fisher Linear Discriminant Derivation

- Similarly $\tilde{\mathbf{s}}_2^2 = \mathbf{v}^t \mathbf{S}_2 \mathbf{v}$
 - Therefore $\tilde{\mathbf{s}}_1^2 + \tilde{\mathbf{s}}_2^2 = \mathbf{v}^t \mathbf{S}_1 \mathbf{v} + \mathbf{v}^t \mathbf{S}_2 \mathbf{v} = \mathbf{v}^t \mathbf{S}_w \mathbf{v}$
 - Define between the class scatter matrix
- \mathbf{S}_B measures separation between the means of two classes (before projection)
 - Let's rewrite the separations of the projected means

$$\begin{aligned}(\tilde{\mu}_1 - \tilde{\mu}_2)^2 &= (\mathbf{v}^t \mu_1 - \mathbf{v}^t \mu_2)^2 \\&= \mathbf{v}^t (\mu_1 - \mu_2)(\mu_1 - \mu_2)^t \mathbf{v} \\&= \mathbf{v}^t \mathbf{S}_B \mathbf{v}\end{aligned}$$

Fisher Linear Discriminant Derivation

- Thus our objective function can be written:

$$\mathbf{J}(\mathbf{v}) = \frac{(\tilde{\mu}_1 - \tilde{\mu}_2)^2}{\tilde{\mathbf{s}}_1^2 + \tilde{\mathbf{s}}_2^2} = \frac{\mathbf{v}^t \mathbf{S}_B \mathbf{v}}{\mathbf{v}^t \mathbf{S}_W \mathbf{v}}$$

- Minimize $\mathbf{J}(\mathbf{v})$ by taking the derivative w.r.t. \mathbf{v} and setting it to 0

$$\begin{aligned} \frac{d}{d\mathbf{v}} \mathbf{J}(\mathbf{v}) &= \frac{\left(\frac{d}{d\mathbf{v}} \mathbf{v}^t \mathbf{S}_B \mathbf{v} \right) \mathbf{v}^t \mathbf{S}_W \mathbf{v} - \left(\frac{d}{d\mathbf{v}} \mathbf{v}^t \mathbf{S}_W \mathbf{v} \right) \mathbf{v}^t \mathbf{S}_B \mathbf{v}}{(\mathbf{v}^t \mathbf{S}_W \mathbf{v})^2} \\ &= \frac{(2\mathbf{S}_B \mathbf{v}) \mathbf{v}^t \mathbf{S}_W \mathbf{v} - (2\mathbf{S}_W \mathbf{v}) \mathbf{v}^t \mathbf{S}_B \mathbf{v}}{(\mathbf{v}^t \mathbf{S}_W \mathbf{v})^2} = 0 \end{aligned}$$

Fisher Linear Discriminant Derivation

- Need to solve $\mathbf{v}^t \mathbf{S}_W \mathbf{v} (\mathbf{S}_B \mathbf{v}) - \mathbf{v}^t \mathbf{S}_B \mathbf{v} (\mathbf{S}_W \mathbf{v}) = 0$

$$\Rightarrow \frac{\mathbf{v}^t \mathbf{S}_W \mathbf{v} (\mathbf{S}_B \mathbf{v})}{\mathbf{v}^t \mathbf{S}_W \mathbf{v}} - \frac{\mathbf{v}^t \mathbf{S}_B \mathbf{v} (\mathbf{S}_W \mathbf{v})}{\mathbf{v}^t \mathbf{S}_W \mathbf{v}} = 0$$

$$\Rightarrow \mathbf{S}_B \mathbf{v} - \frac{\mathbf{v}^t \mathbf{S}_B \mathbf{v} (\mathbf{S}_W \mathbf{v})}{\mathbf{v}^t \mathbf{S}_W \mathbf{v}} = 0$$

$$\Rightarrow \underbrace{\mathbf{S}_B \mathbf{v}} = \lambda \mathbf{S}_W \mathbf{v}$$

generalized eigenvalue problem

Fisher Linear Discriminant Derivation

$$\mathbf{S}_B \mathbf{v} = \lambda \mathbf{S}_W \mathbf{v}$$

- If \mathbf{S}_W has full rank (the inverse exists), can convert this to a standard eigenvalue problem

$$\mathbf{S}_W^{-1} \mathbf{S}_B \mathbf{v} = \lambda \mathbf{v}$$

- But $\mathbf{S}_B \mathbf{x}$ for any vector \mathbf{x} , points in the same direction as $\mu_1 - \mu_2$

$$\mathbf{S}_B \mathbf{x} = (\mu_1 - \mu_2)(\mu_1 - \mu_2)^t \mathbf{x} = (\mu_1 - \mu_2) \underbrace{((\mu_1 - \mu_2)^t \mathbf{x})}_{\alpha} = \alpha(\mu_1 - \mu_2)$$

- Thus can solve the eigenvalue problem immediately


$$\mathbf{v} = \mathbf{S}_W^{-1}(\mu_1 - \mu_2)$$

$$\mathbf{S}_W^{-1} \mathbf{S}_B \underbrace{[\mathbf{S}_W^{-1}(\mu_1 - \mu_2)]}_{\mathbf{v}} = \mathbf{S}_W^{-1} [\alpha(\mu_1 - \mu_2)] = \underbrace{\alpha}_{\lambda} \underbrace{[\mathbf{S}_W^{-1}(\mu_1 - \mu_2)]}_{\mathbf{v}}$$


Fisher Linear Discriminant Example

- Data
 - Class 1 has 5 samples $\mathbf{c}_1 = [(1,2), (2,3), (3,3), (4,5), (5,5)]$
 - Class 2 has 6 samples $\mathbf{c}_2 = [(1,0), (2,1), (3,1), (3,2), (5,3), (6,5)]$
- Arrange data in 2 separate matrices

$$\mathbf{c}_1 = \begin{bmatrix} 1 & 2 \\ \vdots & \vdots \\ 5 & 5 \end{bmatrix} \quad \mathbf{c}_2 = \begin{bmatrix} 1 & 0 \\ \vdots & \vdots \\ 6 & 5 \end{bmatrix}$$



- Notice that PCA performs very poorly on this data because the direction of largest variance is not helpful for classification



Fisher Linear Discriminant Example

- First compute the mean for each class

$$\mu_1 = \text{mean}(\mathbf{c}_1) = [3 \quad 3.6] \qquad \mu_2 = \text{mean}(\mathbf{c}_2) = [3.3 \quad 2]$$

- Compute scatter matrices \mathbf{S}_1 and \mathbf{S}_2 for each class

$$\mathbf{S}_1 = 4 * \text{cov}(\mathbf{c}_1) = \begin{bmatrix} 10 & 8.0 \\ 8.0 & 7.2 \end{bmatrix} \qquad \mathbf{S}_2 = 5 * \text{cov}(\mathbf{c}_2) = \begin{bmatrix} 17.3 & 16 \\ 16 & 16 \end{bmatrix}$$

- Within the class scatter:

$$\mathbf{S}_W = \mathbf{S}_1 + \mathbf{S}_2 = \begin{bmatrix} 27.3 & 24 \\ 24 & 23.2 \end{bmatrix}$$

- it has full rank, don't have to solve for eigenvalues


- The inverse of \mathbf{S}_W is $\mathbf{S}_W^{-1} = \text{inv}(\mathbf{S}_W) = \begin{bmatrix} 0.39 & -0.41 \\ -0.41 & 0.47 \end{bmatrix}$

- Finally, the optimal line direction \mathbf{v}

$$\mathbf{v} = \mathbf{S}_W^{-1}(\mu_1 - \mu_2) = \begin{bmatrix} -0.79 \\ 0.89 \end{bmatrix}$$

Fisher Linear Discriminant Example

- Notice, as long as the line has the right direction, its exact position does not matter
- Last step is to compute the actual **1D** vector **y**. Let's do it separately for each class




$$Y_1 = \mathbf{v}^t \mathbf{c}_1^t = \begin{bmatrix} -0.65 & 0.73 \end{bmatrix} \begin{bmatrix} 1 \cdots 5 \\ 2 \cdots 5 \end{bmatrix} = \begin{bmatrix} 0.81 \cdots 0.4 \end{bmatrix}$$

$$Y_2 = \mathbf{v}^t \mathbf{c}_2^t = \begin{bmatrix} -0.65 & 0.73 \end{bmatrix} \begin{bmatrix} 1 \cdots 6 \\ 0 \cdots 5 \end{bmatrix} = \begin{bmatrix} -0.65 \cdots -0.25 \end{bmatrix}$$

Multiple Discriminant Analysis (MDA)

- Can generalize FLD to multiple classes
- In case of c classes, can reduce dimensionality to 1, 2, 3, ..., $c-1$ dimensions
- Project sample \mathbf{x}_i to a linear subspace $\mathbf{y}_i = \mathbf{V}^t \mathbf{x}_i$
 - \mathbf{V} is called projection matrix



Multiple Discriminant Analysis (MDA)

- Let
 - n_i by the number of samples of class i
 - and μ_i be the sample mean of class i
 - μ be the total mean of all samples

$$\mu_i = \frac{1}{n_i} \sum_{x \in \text{class } i} x \quad \mu = \frac{1}{n} \sum_{x_i} x_i$$

- Objective function: $J(V) = \frac{\det(V^t S_B V)}{\det(V^t S_W V)}$

- within the class scatter matrix S_W is

$$S_W = \sum_{i=1}^c S_i = \sum_{i=1}^c \sum_{x_k \in \text{class } i} (x_k - \mu_i)(x_k - \mu_i)^t$$

- between the class scatter matrix S_B is

$$S_B = \sum_{i=1}^c n_i (\mu_i - \mu)(\mu_i - \mu)^t$$

maximum rank is $c - 1$

Multiple Discriminant Analysis (MDA)

$$J(V) = \frac{\det(V^t S_B V)}{\det(V^t S_W V)}$$


- First solve the **generalized eigenvalue** problem:
$$S_B \mathbf{v} = \lambda S_W \mathbf{v}$$
- At most **$c-1$** distinct solution eigenvalues
- Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{c-1}$ be the corresponding eigenvectors
- The optimal projection matrix V to a subspace of dimension k is given by the eigenvectors corresponding to the largest k eigenvalues
- Thus can project to a subspace of dimension at most **$c-1$**


FDA and MDA Drawbacks


- Reduces dimension only to $k = c - 1$ (unlike PCA)
 - For complex data, projection to even the best line may result in unseparable projected samples


- Will fail:

1. $J(\mathbf{v})$ is always 0: happens if $\mu_1 = \mu_2$



PCA performs
reasonably well
here: 



PCA also
fails: 

2. If $J(\mathbf{v})$ is always large: classes have large overlap when projected to any line (PCA will also fail)

