



# 背景知识：医学文献的结构

## 正文常包含以下模块：

- 1、引言 (Introduction)
- 2、结果 (Results)
- 3、讨论 (Discussion)
- 4、方法 (Method)

## 参考文献

### Review

## Studying CaMKII: Tools and standards

Carolyn Nicole Brown<sup>1</sup> and Karl Ulrich Beyer<sup>1,2</sup>

<sup>1</sup>Department of Pharmacology, University of Colorado Anschutz Medical Campus, Aurora, CO 80045, USA

<sup>2</sup>Correspondence: [karl.beyer@ucdenver.edu](mailto:karl.beyer@ucdenver.edu)

<https://doi.org/10.1016/j.celrep.2024.113982>

### SUMMARY

The  $\text{Ca}^{2+}$ /calmodulin (CaM)-dependent protein kinase II (CaMKII) is a ubiquitous mediator of cellular  $\text{Ca}^{2+}$  signals with both enzymatic and structural functions. Here, we briefly introduce the complex regulation of CaMKII and then provide a comprehensive overview of the expanding toolbox to study CaMKII. Beyond a variety of distinct mutants, these tools now include optical methods for measurement and manipulation, with the latter including light-induced inhibition, stimulation, and sequestration. Perhaps most importantly, there are now three mechanistically distinct classes of specific CaMKII inhibitors, and their combined use enables the interrogation of CaMKII functions in a manner that is powerful and sophisticated yet also accessible. This review aims to provide guidelines for the interpretation of the results obtained with these tools, with careful consideration of their direct and indirect effects.

### INTRODUCTION

The  $\text{Ca}^{2+}$ /calmodulin (CaM)-dependent protein kinase II (CaMKII) is most famous for its prominent roles in neurons<sup>1,2</sup> and in the heart.<sup>3,4</sup> However, at least one CaMKII isozyme is expressed in any cell type examined. Additionally, CaMKII is a multifunctional protein kinase that can phosphorylate a large variety of substrate proteins at Ser/Thr residues and thereby contribute to a variety of  $\text{Ca}^{2+}$ -induced functions. This review aims to help you, the reader, to study the role of CaMKII in your favorite cell type and/or your favorite cellular function. Thus, while we will provide an overview of some sophisticated methods such as light-induced manipulations, we will focus on approaches that are readily accessible: pharmacological inhibitors. Such CaMKII inhibitors now include three different classes with distinct mechanisms of inhibition, making these pharmacological approaches more powerful than ever, especially with some guidance regarding the interpretation of the results. We will emphasize both primary and secondary effects, both for pharmacological inhibitors and for tool mutants. First, however, we will provide a brief overview of the complex regulation of CaMKII.

### THE MOLECULAR BASIS OF CaMKII REGULATION

CaMKII is a family of four isozymes ( $\alpha$ ,  $\beta$ ,  $\gamma$ , and  $\delta$ ) encoded by different genes, with alternative splicing giving rise to more diversity.<sup>1,2,5</sup> Unless noted otherwise, any amino acid residues mentioned in this review will refer to the brain-specific CaMKII $\alpha$  isozyme, which was also the first described member of the larger family of CaM kinases; the numbers of the homologous residue in the other three isozymes are typically one numerical value higher.

### CaMKII structure and stimulation by $\text{Ca}^{2+}$ /CaM

Every CaMKII isozyme contains an N-terminal kinase domain, followed by a short regulatory domain that contains the binding

site for  $\text{Ca}^{2+}$ /CaM, a variable linker region that is subject to alternative splicing, and a C-terminal association or hub domain (Figure 1A) that mediates the formation of predominantly 12meric holoenzymes (Figure 1B). The holoenzymes are largely in an extended conformation, with the kinase domains radiating outward from the central hub and with the  $\text{Ca}^{2+}$ /CaM-binding site of the regulatory domain accessible (Figures 1B and 1C). Notably, while the central hub is rigid, the positioning of the kinase domain is much more flexible.<sup>6,7</sup>

The activity of each individual CaMKII subunit is stimulated by the direct binding of  $\text{Ca}^{2+}$ /CaM to its regulatory domain (Figures 1C and 1D). This  $\text{Ca}^{2+}$ /CaM binding likely enhances ATP affinity,<sup>8</sup> but the vast majority of the dramatic ~1,000-fold increase in enzymatic kinase activity is mediated by enabling substrate access. In the basal state (Figure 1C) the regulatory domain is bound to the kinase domain, in part via interactions of the region around T286 on the regulatory domain with the T286-binding site on the kinase domain (T site), which results also in the block of substrate access to the neighboring substrate-binding site (S site).  $\text{Ca}^{2+}$ /CaM binding to the regulatory domain then displaces it, now allowing substrate access to the S site on the kinase domain, thereby allowing enzymatic kinase activity (i.e., the phosphorylation of the hydroxyl group of a Ser or Thr residue on a substrate protein).

### $\text{Ca}^{2+}$ /CaM-induced autophosphorylation at T286

Displacing the regulatory domain allows access to several features of CaMKII: (1) to the S site, which enables substrate binding and thereby enzymatic activity (as described above); (2) to the T site, which enables binding to other proteins (as described in the next paragraph); and (3) to T286 on the regulatory domain, thereby making T286 accessible for phosphorylation by a neighboring subunit within the holoenzyme (Figure 1E). Once phosphorylated, pT286 prevents complete reassociation of the regulatory domain with the T site, thereby keeping the CaMKII in a





# 在文献库中检索文献

Pubmed, ScienceDirect, Web of Science, Embase, Cochrane...



ScienceDirect

Web of Science™




Embase



# 在文献库中检索文献

## 常规搜索

以Pubmed为例，通过关键词进行文献检索



EasyPubMed 浏览器插件

# 在文献库中检索文献：高级检索

举例：如何检索某个特定的老师？

浙江大学张岩与北京大学张岩课题组在《细胞》发表 ...  
2024年3月3日 · 该研究基于结构解析设计了完全偏向于Gi通路的APLNR激动剂，可降低心肌肥厚和心衰的风险，为心脏疾病的治疗提供新策略。浙江大学张 ...



PubMed®  
User Guide

Add terms to the query box

Affiliation



Enter a search term



AND



Show Index

Query box

(Zhang Yan[Author]) AND (Zhejiang University[Affiliation])



Search



# 查阅文献的两大痛

如何查得全面？

如何查得高效？



# 全面查找：从文献到文献

一篇好综述是一个方向的说明手册

进一步明确方向，进而细化关键词

丰富的参考文献

# 全面查找：从文献到文献



研究论著类 (article) 同样可以作为文献导航

查作者

查推荐

查参考文献



# 全面查找：从文献到文献

研究论著类 (article) 同样可以作为文献导航

查作者

查推荐

查参考文献


# 全面查找：从文献到文献

全收集：文献→文献→文献

整理：

按照发表时间排序

文献整理软件，如Zotero、Endnote等



# 高效查找：降低阅读门槛

学位论文：中国知网、北京大学图书馆等

高质量的中文综述

高质量的中文研究方法

了解国内该领域的主要研究者



(知网终于上桌了)

# 高效查找：降低阅读门槛



科普公众号：BioArt、iNature、ScienceAAAS等




定期分享最新发表的工作，中文撰写、高度凝练



BioArt ★  
科技自媒体  
上海 已实名



iNature ★  
科普自媒体  
浙江 已实名



ScienceAAAS  
北京



Nature | 成纤维细胞——脑损伤修复的“  
双面卫士”



Nat Genet | 端粒损耗成为衰老造血和白  
血病发生中克隆选择的工具



STM | 秦士杰等合作完成刚毛棉鼠基因组  
测序并揭示其与人类保守易感呼...



Trends Mol Biol | 临床基因治疗恢复听力：  
一场范式转变






# 高效查找: 降低阅读门槛

## AI工具: 秘塔、玻尔等

引用真实文献, 避免“瞎编文献”的AI幻觉



请为我介绍深度学习辅助的蛋白质设计技术的应用...

学术 简读

10

新讲解

深度学习辅助的蛋白质设计技术是一种利用深度学习算法来设计和优化蛋白质结构与功能的方法。该技术通过深度学习模型(如卷积神经网络、生成对抗网络、变分自编码器)来处理蛋白质序列、结构和功能数据,从而实现蛋白质的高效设计和优化。

深度学习在蛋白质设计中的应用主要包括以下几个方面:

1. 蛋白质结构预测与设计: 深度学习模型(如 AlphaFold, AlphaFold2)在蛋白质结构预测方面取得了显著进展,为蛋白质设计提供了基础。深度学习模型可以用于预测蛋白质的三维结构、折叠方式和功能。
2. 蛋白质序列设计: 深度学习模型可以生成新的蛋白质序列,以实现特定的功能或结构。

International Journal of Molecular Sciences

MDPI

Review

### Protein Design with Deep Learning

Marianne Defresne <sup>1,2</sup>, Sophie Barbe <sup>1</sup> and Thomas Schiex <sup>2,\*</sup>

<sup>1</sup> Toulouse Biotechnology Institute, Université de Toulouse, CNRS, INRAE, INSA, ANITI, 31077 Toulouse, France; marianne.defresne@insa-toulouse.fr (M.D.); sophie.barbe@insa-toulouse.fr (S.B.)  
<sup>2</sup> Université Fédérale de Toulouse, ANITI, INRAE, UR 675, 31326 Toulouse, France  
\* Correspondence: thomas.schiex@insa-toulouse.fr; Tel.: +33-561-28-5428

**Abstract:** Computational Protein Design (CPD) has produced impressive results for engineering new proteins, resulting in a wide variety of applications. In the past few years, various efforts have aimed at replacing or improving existing design methods using Deep Learning technology to leverage the amount of publicly available protein data. Deep Learning (DL) is a very powerful tool to extract patterns from raw data, provided that data are formatted as mathematical objects and the architecture processing them is well suited to the targeted problem. In the case of protein data, specific representations are needed for both the amino acid sequence and the protein structure in order to capture respectively 1D and 3D information. As no consensus has been reached about the most suitable representations, this review describes the representations used so far, discusses their strengths and weaknesses, and details their associated DL architecture for design and related tasks.