Neuromechanics of Human Motion

Perception, Illusions, & Bayes' Theorem

Joshua Cashaback, PhD



Recap — Action Potentials

- 1. Basic nerve anatomy
- 2. Know what contributes to resting membrane potential
 - a. ions, electrochemical gradients, leak channels and Na-K pumps
- 3. Understand how action potentials occur
 - a. voltage gates, depolarization, repolarization, hyperpolarization (refractory period)
- 4. Hodgkin-Huxley Model



Recap — Action Potentials

- 1. Basic nerve anatomy
- 2. Know what contributes to resting membrane potential
 - a. ions, electrochemical gradients, leak channels and Na-K pumps
- 3. Understand how action potentials occur
 - a. voltage gates, depolarization, repolarization, hyperpolarization (refractory period)
- 4. Hodgkin-Huxley Model

Any questions on the assignment?



Lecture Objectives — Perception, Illusions, Bayes

- 1. Define Perception and Sensory Illusion.
- 2. Discuss (multi)sensory illusions
- 3. Discuss how the brain uses prior experience
- 4. Examples of how the brain integrates multiple senses
- 5. Learn about noise in the nervous system
- 6. Probabilities Primer (Bayes Theorem)




Perception

the organization, identification, and interpretation of sensory information to represent and understand the environment.



Perception

the organization, identification, and interpretation of sensory information to represent and understand the environment.



Motor Learning (elite, new skill)



Perception

the organization, identification, and interpretation of sensory information to represent and understand the environment.



autism, dyslexia



Perception

the organization, identification, and interpretation of sensory information to represent and understand the environment.



Neuroscience: understand the brain before you can fix it.

Large portions of our brain are involved with sensory information,
percept and integrating both together.

Perception

the organization, identification, and interpretation of sensory information to represent and understand the environment.



Technology creation

- Robotics
- · Face recognition
- · Immersive virtual reality

human-brain interfaces (HBI), neurolink

how does the brain transform sensory information?



What is a Sensory Illusion?

Sensory Illusion

Where perception deviates from reality.





What is a Sensory Illusion?

Sensory Illusion

Where perception deviates from reality.



A way to probe perception



What is a Sensory Illusion?

Sensory Illusion

Where perception deviates from reality.



A way to probe perception

Visual, Auditory, Haptic, Propriception, Multisensory



Illusions



Illusions

Before we begin...



Illusions




Warning!

Some of the following images may cause nausea

Checkerboard patterns may cause seizures in certain forms of epilepsy




Visual Illusions — Circles






Visual Illusions — Luminence



Pupil size is actually influenced by this illusion (Laeng and Endestad, 2012)




Visual Illusions — Diamonds in the Air






Visual Illusions — Moving Snakes






Visual Illusions — Tilt and Movement





Proprioception Illusion — Position Sense


Goodwin, McCloskey, Matthews (1972) Science




Biceps vibration causes arm flexion and a mismatch between actual and felt movement

Triceps vibration causes arm extension and a mismatch between actual and felt movement

Tracking passive movement is very accurate








Insight into Sensory Physiology




Probe Perception



1. Insight into Sensory Physiology





2. Insight into how the BRAIN integrates sensory information

Hollow Face Illusion

http://www.youtube.com/watch?v=sKa0eaKsdA0




Hollow Face Illusion

http://www.youtube.com/watch?v=sKa0eaKsdA0


Your Brain fills in blanks based on prior experience






Different size boxes, same weight





Transducers to record force, measure motor output, and probe expectations





Big box feels light





small box feels heavier that big box!






Our prior experience and expectation modifies our perception






How does this influence our motor output?





Mismatch, despite same weight






Adaptation




Grip/lift errors are rapidly corrected





Size Weight Illusion



Despite motor output changes, illusion remains (e.g., thick, dashed line = participants keep reporting '7')



Size Weight Illusion — What can it tell us?





- 1. We don't always know the physical properties of our world
- 2. Our brain fills in gaps (based on prior experience)
- 3. Sensorimotor system can operate independent of conscious perception
 - a. perception influenced early behaviour but stayed constant while control changed.
 - b. (partially) separate systems for judging weight and lifting objects




Multisensory Integration





McGurk Effect — Visual and Auditory




The McGurk effect is a perceptual phenomenon that demonstrates an interaction between hearing and vision in speech perception.

The illusion occurs when the auditory component of one sound (ba) is paired with the visual component of another sound (ga), leading to the perception of a third sound (da).

https://www.youtube.com/watch?v=G-IN8vWm3m0 (note, video using different combination of syllables)



McGurk Effect — Visual and Auditory




The McGurk effect is a perceptual phenomenon that demonstrates an interaction between hearing and vision in speech perception.

The illusion occurs when the auditory component of one sound (ba) is paired with the visual component of another sound (ga), leading to the perception of a third sound (da).

Our Brain combines our sensory inputs together (even when they are in conflict!)




Rubber Hand Illusion — Visual and Somatosensory



Jared Madina (UD psych): https://www.youtube.com/watch?v=RhHzcVdyvWg




Rubber Hand Illusion — Visual and Somatosensory



Funny Extension: https://www.youtube.com/watch?v=MG22iFL-VgE



Rubber Hand Illusion — Visual and Somatosensory



- 1. Combining vision and somatosensory
- 2. Adaptation
- 3. Phantom limb pain



Noise in the Nervous System





Figure 1) Overview of the behavioural loop and the stages at which noise is present in the nervous system, and sources of extension proise induct the transduction of signals. This is exemplified here by a photocorpora and is signal, amplification cascade is 15 sources of cellular noise include the init channels of excitable membranes, spragic transmission and interest in the section of the second of the s



Optimally Combining Noisy Sensors



Ernst and Banks (2002) Nature

- 1. Nervous system is statistically optimal
- 2. Noise typically modelled with Gaussian or Normal probability distributions



Bayesian Integration

Bayes' Theorem can account for:

- 1. The role of prior experience
- 2. Adaptation
- 3. Sensory noise (statistically optimal)
 - a. Multisensory integration (explain illusions)
 - b. cost functions
 - c. state estimation (e.g., Kalman filters)



Probability Primer

- 1. Mutually Exclusive (Disjoint)
- 2. Joint Probabilitites
- 3. Complement Probability
- 4. Conditional Probability
- 5. Marginal Probability



Probability Primer

- 1. Mutually Exclusive (Disjoint)
- 2. Joint Probabilitites
- 3. Complement Probability
- 4. Conditional Probability
- 5. Marginal Probability

With these defined, we can derive Bayes' theorem!



Notation

S =sample space (all possible outcomes)

p(A) = probability of event A


 $A \cup B =$ union of events A and B

 $A \cap B$ = intersection of events A and B

p(B|A) = probability of B given A

p(A') or $p(A^C)$ or $p(\bar{A}) = \text{complement probability of } p(A)$

Mutually Exclusive (Disjoint Probability)




$$p(A \cup B) = p(A) + p(B)$$

0.7 = 0.4 + 0.3




Joint Probability



$$p(A \cup B) = p(A) + p(B) - p(A \cap B)$$
$$0.5 = 0.4 + 0.3 - 0.2$$




Complement Probability




H	Red	Yellow	Green	Marginal probability P(H)
Not Hit	0.198	0.09	0.14	0.428
Hit	0.002	0.01	0.56	0.572
Total	0.2	0.1	0.7	1

Probabilities of getting in an accident at an intersection given different lights















$$p(A \cap B') = p(A) - p(A \cap B)$$



$$p(A \cap B') = p(A) - p(A \cap B)$$

The marginal p(A) or p(B) is found by summating their disjoint parts.

$$p(A) = p(A \cap B) + p(A \cap B')$$
, and similarly $p(B) = p(A \cap B) + p(A' \cap B)$



Conditional Probability Examples


```
p(accepted) = 0.3

p(funding|accepted) = 0.43

p(funding \cap accepted) = p(funding|accepted) \cdot p(accepted)

p(funding \cap accepted) = 0.43 \cdot 0.3 = 0.13
```


Conditional Probability



$$p(A \cap B) = p(B|A) \cdot p(A)$$
$$p(B|A) = \frac{p(A \cap B)}{p(A)}$$



Conditional Probability



$$p(A \cap B) = p(B|A) \cdot p(A)$$


$$p(B|A) = \frac{p(A \cap B)}{p(A)}$$

$$p(A \cap B) = p(A|B) \cdot p(B) \text{ (in terms of B)}$$

$$p(A|B) = \frac{p(A \cap B)}{p(B)}$$



Conditional Probability



$$p(A \cap B) = p(B|A) \cdot p(A)$$


$$p(B|A) = \frac{p(A \cap B)}{p(A)}$$

$$p(A \cap B) = p(A|B) \cdot p(B) \text{ (in terms of B)}$$

$$p(A|B) = \frac{p(A \cap B)}{p(B)}$$




Conditional Probability Complements






Conditional Probability Complements



$$p(A \cap B') = p(B'|A) \cdot p(A)$$

Conditional Probability Complements



$$p(A \cap B') = p(B'|A) \cdot p(A)$$

Other friendly complements:
 $p(A' \cap B) = p(B|A') \cdot p(A')$
 $p(A' \cap B') = p(B'|A') \cdot p(A')$



Bayes' Theorem



Reverand Thomas Bayes (1701-1761)

Good News! Bayes' Theorem is simply a conditional probability!



Deriving Bayes' Theorem

Remember:

$$p(A|B) = \frac{p(A \cap B)}{p(B)}, (eq.1) \text{(slide 45)}$$

$$p(A \cap B) = p(B|A) \cdot p(A), (eq.2) \text{(slide 44)}$$

Substitute (eq.2) into (eq.1):

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$

Thats it!



$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$



$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$

Calculate p(B) by using its marginal probability

$$p(B) = p(A \cap B) + p(A' \cap B), (eq.4) \text{(slide 42)}$$



$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$

Calculate p(B) by using its marginal probability

$$p(B) = p(A \cap B) + p(A' \cap B), (eq.4) \text{(slide 42)}$$

Substitute (eq.4) into (eq.3)

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(A \cap B) + p(A' \cap B)}, (eq.5)$$



$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}$$
, (eq.3)
Calculate p(B) by using its marginal probability
 $p(B) = p(A \cap B) + p(A' \cap B)$, (eq.4)(slide 42)
Substitute (eq.4) into (eq.3)
 $p(A|B) = \frac{p(B|A) \cdot p(A)}{p(A \cap B) + p(A' \cap B)}$, (eq.5)
Since,
 $p(A \cap B) = p(B|A) \cdot p(A)$, (eq.6)(slide 44)
 $p(A' \cap B) = p(B|A') \cdot p(A')$, (eq.7)(slide 49)

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$

Calculate p(B) by using its marginal probability

$$p(B) = p(A \cap B) + p(A' \cap B), (eq.4) \text{ (slide 42)}$$

Substitute (eq.4) into (eq.3)

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(A \cap B) + p(A' \cap B)}, (eq.5)$$

Since.

$$p(A \cap B) = p(B|A) \cdot p(A), (eq.6) \text{ (slide 44)}$$

$$p(A' \cap B) = p(B|A') \cdot p(A'), (eq.7) \text{(slide 49)}$$

Substitute (eq.6) and (eq.7) into (eq.5)

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B|A) \cdot p(A) + p(B|A') \cdot p(A')}, (eq.8)$$



Alternative Definition

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}, (eq.3)$$
$$p(B|A) = \frac{p(A|B) \cdot p(B)}{p(A)}$$



Why Bayesian?

Powerful way to continually account for new evidence given prior beliefs

$$p(A|B) = \frac{p(B|A) \cdot p(A)}{p(B)}$$



Next Class

- 1. point estimate examples
- 2. continuous probability
- 3. combining multiple senses
- 4. cost functions



Homework

1. Plot a Normal probability distribution

$$p(x|\mu,\sigma^2) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

- . manipulate the mean (μ) and standard deviation (σ)
- . discretize along x (make the number of data points used a variable)

2. Plot a Beta probability distribution

$$p(x|\alpha,\beta) = \frac{1}{B(\alpha,\beta)} x^{\alpha-1} (1-x)^{\beta-1}$$

- . manipulate α and β
- . from scipy import special
- . B = scipy.special.beta(alpha, beta)
- . discretize along x (make the number of data points used a variable)



Acknowledgements

Gavin Buckingham Michael Barnett-Cowan

