## Years in which Broadbalk soil was sampled 1843-1944, and chemical properties re-analysed in 2000.

Year	Depth	%N	%SOC	Olsen P	рН	Exchangeable cations \$	CaCO3	No. plots
1865	Topsoil	٧ *	٧ *	٧ *	٧ *	٧*	٧*	6
	Subsoil	٧ *	٧ *	٧ *		٧*	٧ *	6
1881	Topsoil	٧ *	٧*	٧ *	٧ *	٧*	٧*	72
	Subsoil	٧ *	٧ *	٧ *		٧*	٧ *	16
1893	Topsoil	٧ *	٧*	٧ *	٧ *	٧*	٧*	52
	Subsoil	٧ *	٧ *	٧ *		٧ *	٧ *	36
1914	Topsoil	٧ *	٧ *	٧ *	٧ *	٧*	٧ *	56
	Subsoil	٧ *	٧ *	٧ *		٧ *	٧ *	40
1936	Topsoil	٧ *	٧ *	٧ *	٧ *	٧ *	٧ *	23
1944	Topsoil	٧ *	٧*	٧ *	٧ *	٧*	٧*	70
	Subsoil	٧ *	√ *	٧ *	٧ *	√ *	٧*	50

 $<sup>\</sup>vee$  \* Selected plots and sections. This table lists the major soil samplings; other samples will have been taken from selected plots and sections in other years, to investigate specific questions.

<sup>\$</sup> Exchangeable cations – Ca, K, Mg and Na (no Na in 1881 or 1944)