

## Lecture 1 Farkas's Lemma, Strong Duality and Criss-Cross Algorithm

Instructor: Tamás Terlaky

Scribe: Yutong Dai

Last Modified: 2019-10-14

**Disclaimer:** *This note has not been subjected to the usual scrutiny reserved for formal publications.*

Consider the following primal-dual problem

$$\left\{ \begin{array}{ll} \min & c^T x \\ \text{s.t.} & Ax = b \\ & x \geq 0, \end{array} \right. (P) \quad \left\{ \begin{array}{ll} \max & b^T y \\ \text{s.t.} & A^T y \leq c \end{array} \right. (D).$$

Correspondingly, we have following version of Farkas' Lemma

**Lemma 1.1: Farkas' Lemma (Primal Form)**

One and only one of the following two systems of linear inequalities has a feasible solution.

$$(I_p) \left\{ \begin{array}{l} Ax = b \\ x \geq 0 \end{array} \right. \quad (II_p) \left\{ \begin{array}{l} A^T y \leq 0 \\ b^T y > 0 \end{array} \right. .$$

**Theorem 1.2: Weak Duality**

For any primal-dual feasible solution pair  $(x, y)$ , we have  $c^T x \geq b^T y$ . Moreover,  $=$  holds iff

$$x^T (A^T y - c) = 0,$$

which is known as the complementary slackness condition.

**Theorem 1.3: Strong Duality**

Exactly one of the following could happen for a  $(P) - (D)$  problem pair.

1. If both the primal and dual problems have a feasible solution, then for any optimal solutions  $\bar{x}$  and  $\bar{y}$ , we have  $c^T \bar{x} = b^T \bar{y}$ .
2. If the primal infeasible and the dual is feasible, then the dual is unbounded.
3. If the dual infeasible and the primal is feasible, then the primal is unbounded.
4. Both the primal and dual can be infeasible.

## The equivalence of Farkas' Lemma and Strong Duality

**First, let's consider when strong duality holds.** Then consider the following case,  $(P)$  is infeasible with  $c = 0$  and  $(D)$  is feasible. Then, we know that  $(D)$  is unbounded. Therefore, we know that  $\exists y^*$  such that  $A^T y^* \leq 0$  and  $b^T y^* > 0$ . This implies, when  $(II_p)$  is feasible,  $(I_p)$  is infeasible.

**Then we consider using Farkas' Lemma to prove Strong duality.**

1. By weak duality, we have  $c^T x \geq b^T y$ , for any feasible solution pair  $(x, y)$ . Now, we want to prove  $b^T y \geq c^T x$ . It's equivalent to prove the system (1.1) is feasible. So if (1.1) is feasible, then we have (1) in strong duality proved.

$$\left\{ \begin{array}{l} -b^T y + c^T x + \rho = 0, \rho \geq 0 \\ Ax = b, x \geq 0 \\ A^T y + s = b, s \geq 0, y \text{ is free.} \end{array} \right. \implies \underbrace{\begin{bmatrix} 0 & 0 & A & 0 & 0 \\ -A^T & A^T & 0 & I & 0 \\ b^T & -b^T & c^T & 0 & 1 \end{bmatrix}}_{\bar{A}} \underbrace{\begin{bmatrix} y^- \\ y^+ \\ x \\ s \\ \rho \end{bmatrix}}_{\bar{x}} = \underbrace{\begin{bmatrix} b \\ c \\ 0 \end{bmatrix}}_{\bar{b}} \quad (1.1)$$

2. Assume the system (1.1) is infeasible, then by Farkas' lemma, we know that the system (1.2) must be feasible.

$$\left\{ \begin{array}{l} \bar{A}^T \bar{y} \leq 0 \\ \bar{b}^T \bar{y} > 0 \end{array} \right. \quad \text{where } \bar{y} = \begin{bmatrix} u \in \mathbb{R}^m \\ v \in \mathbb{R}^n \\ l \in \mathbb{R}^1 \end{bmatrix} \implies \left\{ \begin{array}{l} Av = bl \\ A^T u + cl \leq 0 \\ b^T u + c^T v > 0 \\ v \leq 0, l \leq 0 \end{array} \right. \quad (1.2)$$

Set  $\bar{l} = -l \geq 0, \bar{v} = -v \geq 0$ . If  $\bar{l} > 0$ . Then set  $x = \frac{\bar{v}}{\bar{l}}, y = \frac{u}{\bar{l}}$ . It's easy to verify that  $(x, y)$  is a feasible primal-dual solution pair. But  $b^T y = \frac{1}{\bar{l}} b^T u > \frac{1}{\bar{l}} c^T \bar{v} = c^T x$ , which contradicts with the weak duality. This implies  $\bar{l} = 0$ .

Provided  $\bar{l} = 0$ , then (1.3) always holds.

$$\mathbb{P} \left\{ \begin{array}{l} A\bar{v} = 0 \\ \bar{v} \geq 0 \end{array} \right. \quad \mathbb{Q} \left\{ \begin{array}{l} A^T u \leq 0 \\ b^T u > c^T \bar{v} \end{array} \right. \quad (1.3)$$

As  $\mathbb{P}$  and  $\mathbb{Q}$  both hold, by Farkas' lemma, we know  $c^T \bar{v} \neq 0$ .

- $c^T \bar{v} < 0$ . Assume the primal is feasible and dual is infeasible. Consider a primal feasible solution  $x^*$ , then  $x(\alpha) = x^* + \alpha \bar{v}$ , where  $\alpha > 0$ , is also a solution. Then  $c^T x(\alpha) = c^T x^* + \alpha c^T \bar{v} \rightarrow -\infty$  as  $\alpha \rightarrow +\infty$ . Therefore, the primal is unbounded.
- $c^T \bar{v} > 0$ . This implies  $b^T u > 0$ . Assume the dual is feasible and primal is infeasible. Consider a dual feasible solution  $y^*$ , then  $y(\alpha) = y^* + \alpha u$ , where  $\alpha > 0$ , is also a solution. Then  $b^T y(\alpha) = b^T y^* + \alpha b^T u \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$ . Therefore, the dual is unbounded.

So (2) and (3) in the strong duality theorem are proved. It remains to prove (4). Consider  $A = 0, c = -1, b = 1$ . In this case both (P) and (D) are infeasible.

## Criss-cross Algorithm and the Strong Duality

From Theorem 1.4, we know that the Criss-cross algorithm terminates in finite steps.

1. If the Criss-cross algorithm terminates with feasible a primal-dual solution pair  $(x, y)$ , then the complementary slack conditions hold. By the weak duality, we have  $c^T x = b^T y$ .
2. Suppose the Criss-cross algorithm terminates with following case.

p-th row

$[A_B^{-1}]_p$	-	$\oplus$	$\cdots$	$\oplus$

Then we claim the primal is infeasible. This can be easily seen from fact that the sum of non-negative elements cannot be strictly negative. That is, the no  $x \geq 0$  satisfies the p-th constraint.

Further, if we assume the dual is feasible. Let  $(\bar{y}, \bar{s})$  be a dual feasible solution such that  $A^T \bar{y} + \bar{s} = c$ ,  $\bar{s} \geq 0$ . We need to find a  $\hat{y}$  such that  $b^T \hat{y} > 0$ . Then  $b^T y(\alpha) = b^T \bar{y} + \alpha b^T \hat{y} \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$ .

Let  $\tilde{y}^T =$  p-th row of  $A_B^{-1}$ , then  $\tilde{y}^T b < 0$ ,  $A^T \tilde{y} \geq 0$ . Set  $\hat{y} = -\tilde{y}$ , then  $b^T \hat{y} > 0$ ,  $A^T \hat{y} \geq 0$ <sup>1</sup>. Hence, the dual is unbounded.

3. Suppose the Criss-cross algorithm terminates with following case.

q-th row

	+	
	$\ominus$	
	$\vdots$	
	$\ominus$	
	$A_B^{-1} A[:, q]$	

Then we claim the dual is infeasible. We prove this by using Farkas' lemma. Define

$$t_q = \begin{cases} \tau_{iq}, & i \text{ in the basis} \\ -1, & i=q \\ 0, & i \text{ not in the basis} \end{cases},$$

we can see in the proof of Theorem 1.4 that  $t_q \in \text{Null}(A)$ , i.e.,  $At_q = 0$ . From the table, we know  $t_{iq} = A_B^{-1} A[i, q] \leq 0, i$  in the basis.. Set  $\hat{x} = -t_q$ . Then  $A\hat{x} = 0, \hat{x} \geq 0$ .

$$\begin{aligned} c^T \hat{x} &= c_B^T \hat{x}_B + C_N^T \hat{x}_N \\ &\stackrel{(1)}{=} c_q - c_B^T (A_B^{-1} A[:, q]) \\ &\stackrel{(2)}{=} s_q < 0, \end{aligned}$$

where (1) holds as  $q$  not in the basis and (2) holds as  $-s_q > 0$  (from the tableau). Therefore, we have  $c^T \hat{x} < 0, A\hat{x} = 0, \hat{x} \geq 0$ , the by Farkas's lemma (dual form), we know  $A^T y \leq c$  is infeasible.

<sup>1</sup>This again implies the primal is infeasible from the Farkas' lemma.

Further, suppose the primal is feasible. Let  $\bar{x}$  be a primal feasible solution such that  $A^T \bar{x} = b, \bar{x} \geq 0$ . Since  $c^T \hat{x} < 0$ , then  $c^T x(\alpha) = c^T \bar{x} + \alpha c^T \hat{x} \rightarrow -\infty$  as  $\alpha \rightarrow +\infty$ .

Hence, the primal is unbounded.

## Appendix 1: Farkas' Lemma and its variants

Recall the primal form defined in the Lemma 1.1

$$(I_p) \begin{cases} Ax = b \\ x \geq 0 \end{cases} \quad (II_p) \begin{cases} A^T y \leq 0 \\ b^T y > 0 \end{cases}.$$

It's essentially can be derived from the primal-dual problem pair

$$\begin{cases} \min & c^T x \\ \text{s.t.} & Ax = b \\ & x \geq 0, \end{cases} \quad (P) \quad \begin{cases} \max & b^T y \\ \text{s.t.} & A^T y \leq c \end{cases} \quad (D).$$

$(I_p)$  are just constraints from  $(P)$ . The first constraint in  $(II_p)$  are derived by setting  $c = 0$ , then plug it into the constraint from  $(D)$ . As  $c = 0$ , then if  $(P)$  is feasible, the optimal is 0. In order to construct contradictions, we require  $b^T y > 0$ . As by weak duality we always have  $b^T y \leq 0$ . Then, we recover the second constraint.

As for the proof, if  $(I_p)$  is feasible it is easy to see  $(II_p)$  is infeasible. If  $(I_p)$  is infeasible, then  $b$  not in the cone(columns of  $A$ ). By separation theorem, we can set  $y$  to the norm vector of the separating hyperplane, we have  $(II_p)$  feasible.

An common variant is represented in the dual form,

$$(I_d) \begin{cases} Ax = 0 \\ x \geq 0 \\ c^T x < 0 \end{cases} \quad (II_d) \begin{cases} A^T y \leq c \end{cases}.$$

As for the proof, we can rewrite them to the equivalent form of  $(I_p)$  and  $(II_p)$ . For example, denote  $y = y^+ - y^-$ , and change  $II_d$  to

$$(II'_d) \begin{cases} [A, -A, I] \begin{pmatrix} y^+ \\ y^- \\ s \end{pmatrix} = c \\ y^+, y^-, s \geq 0. \end{cases}.$$

Then  $(II'_d)$  is equivalent to  $I_p$ . Similarly, we can re-write  $I_d$  to the form of  $II_p$ .

## Appendix 2: Criss-Cross Algorithm

- Basic Tableau Setup

For a given coefficient matrix  $A \in \mathbb{R}^{m \times n}$ ,  $m < n$  and  $\text{Rank}(A) = m$ , we can partition  $A$  as  $A = [A_B, A_N]$ , where  $A_B$  is invertible. Denote  $I_B = \{i | i\text{th column of } A \text{ in the } A_B\}$ ,  $I_N = \{1, \dots, n\} \setminus I_B$ . We can rewrite the constraint in (D) as  $A^T y + s = c$ . Then, can partition  $c, s$  according to  $I_B, I_N, x, y$  respectively. Now, the solution pair  $(x, y)$  can be set to

$$x_B = A_B^{-1}b, \quad x_N = \mathbf{0}, \quad s_B = 0, \quad s_N = c_N^T - c_B^T A_B^{-1} A_N.$$

$c_B^T A_B^{-1} b$	$-s_B^T = 0$	$-s_N^T = -c_N^T + c_B^T A_B^{-1} A_N$
$x_B = A_B^{-1} b$	$I$	$A_B^{-1} A_N = (\tau_{ij})$

- Pivot

The pivot step can be described as

$$\begin{aligned}
 I'_B &\leftarrow I_B \cup \{l\} \setminus \{k\} \\
 \tau'_{ij} &= \tau_{ij} - \frac{\tau_{il}\tau_{kj}}{\tau_{kl}} & \forall i \in I'_B \setminus \{l\}; j \in I'_N \setminus \{k\} \\
 \tau'_{ik} &= -\frac{\tau_{il}}{\tau_{kl}}, & \forall i \in I'_B \setminus \{l\} \\
 \tau'_{lj} &= \frac{\tau_{kj}}{\tau_{kl}}, & \forall j \in I'_N \setminus \{l\} \\
 \tau'_{lk} &= \frac{1}{\tau_{kl}}
 \end{aligned}$$

		j	l			j	k
i		$\tau_{ij}$	$\tau_{il}$		i	$\tau'_{ij}$	$\tau'_{ik}$
k		$\tau_{kj}$	$\tau_{kl}$		l	$\tau'_{lj}$	$\tau'_{lk}$

Pivot on  $\tau_{kl} \implies$

- Criss-cross Algorithm Procedure

**Initialization**

let  $A_B$  be an arbitrary initial basis;  
 $I_B$  resp.  $I_N$  is the index set of  
the basis and nonbasis variables;

**while** true **do**

**if**  $x_B \geq 0$  and  $s_N \geq 0$  **then**

(I) **stop**: the current solution solves the LP problem;

**else**

$p := \min\{i \in I_B : x_i < 0\};$

$q := \min\{j \in I_N : \underline{s_j} < 0\};$

$r := \min\{p, q\};$

**if**  $r = p$  **then**

**if** the  $p$ -row of the tableau is nonnegative **then**

(II) **stop**: (LP) is inconsistent;

**else**

let  $q := \min\{j \in I_N : \tau_{pj} < 0\};$

**endif**

**else** (i.e.  $r = q$ )

**if** the  $q$ -column of the tableau is nonpositive **then**

(III) **stop**: (LD) is inconsistent;

**else**

let  $p := \min\{i \in I_B : \tau_{iq} > 0\};$


**endif**

**endif**

perform a pivot:  $I_B := I_B \cup \{q\} \setminus \{p\};$


**endwhile**

**end.**



**Theorem 1.4**

Criss-cross Algorithm terminates in finite steps.



**Proof:** Assume we will visit a basis twice, meaning we are trapped in a cycle. Denote  $q = \max\{i \in \{1, 2, \dots, n\} | i \text{ enters the basis during the cycle.}\} = \max\{i \in \{1, 2, \dots, n\} | i \text{ leaves the basis during the cycle.}\}$ . From the figure, we know that  $x_q$  can only enter the basis via either pattern A or B. Similarly,  $x_q$  can only leave the basis via either pattern C or D.

It remains to prove that are of the following four situations are impossible.

1.  $B \Rightarrow D$
2.  $B \Rightarrow C$
3.  $A \Rightarrow D$
4.  $A \Rightarrow C$

**$B \Rightarrow D$  case:**

The matrix  $A$  can be partitioned as  $A = [A_B \mid A_N]$ , then we can get the coefficient matrix in the tableau  $[I \mid A_B^{-1} A_N]$ . So we know that  $t_p \in \mathbb{R}^{1 \times n}$ , where  $p < q$ , must be

$$[t_p]_j = \begin{cases} 0 & x_j \text{ in the basis and } j \neq p \\ 1 & j = p \\ \tau_{pj} & j \text{ not in the basis} \end{cases}$$


Similarly we can construct a  $t_k \in \mathbb{R}^{n \times 1}$ , where  $k < q$ , such that

$$[t_k]_i = \begin{cases} 0 & x_i \text{ not in the basis and } i \neq k \text{ (1)} \\ -1 & i = k \text{ (2)} \\ \tau_{ik} & i \text{ in the basis (3)}. \end{cases}$$

(3) is directly from the tableau; (1) and (2) are constructed based the following fact

$$\begin{bmatrix} I & A_B^{-1}A_N \end{bmatrix} \begin{bmatrix} A_B^{-1}A_N \\ -I \end{bmatrix} = \mathbf{0}.$$

Therefore, we know that  $\langle t_k, t_p \rangle = 0$ . From the figure below, and the facts that



- 1) the way we define  $t_p$  and  $t_k$ ;
- 2) If  $j > p$ , both sets  $S_1 = \{j | x_j \text{ in the basis}\}$ ,  $S_2 = \{j | x_j \text{ not in the basis}\}$  remain unchanged.

We know that  $\langle t_k, t_p \rangle < 0$ . Contradiction! So  $B \Rightarrow D$  is impossible.

**A  $\Rightarrow$  D case:**

From above analysis, we already construct  $t_k$  such that  $t_k \in \text{Null}(A)$ . So we want to find a vector in  $\text{row}(A)$ .

Note that

$$\begin{cases} A^T y_A + (-S_A) = c \\ A^T y_D + (-S_D) = c, \end{cases}$$

we know  $S \triangleq S_A - S_D = A^T(y_A - y_D)$ . Therefore,  $\langle t_k, t_p \rangle = 0$ .



	$j < q$	$j = q$	$j > q$	
$S_A$	$\ominus$ .... $\ominus$ +		?	
	$k$			
$S_D$	+	0	?	0 if the indices are in the basis
	$k$			
$t_k$	$\ominus$ .... -1 $\ominus$ +		?	0 if the indices are NOT in the basis

Similarly as the previous case, for  $j > q$ ,  $x_j$  will be always [in/not in] basis when  $A \Rightarrow D$ .

- $\langle S_A, t_k \rangle = \underbrace{\text{non-negative}}_{j < q} + \underbrace{\text{positive}}_{j = q} + \underbrace{0}_{j > q} = \text{positive}$
- $\langle S_D, t_k \rangle = \underbrace{0}_{j < q, j \neq k} + \underbrace{\text{negative}}_{j = k} + \underbrace{0}_{j \geq q} = \text{negative}$

Therefore,  $\langle S_A - S_D, t_k \rangle = \text{positive}$ , Hence, contradiction!

**B  $\Rightarrow$  C case:**

	$j < q$	$j = q$	$j > q$	
$X_C$	$\oplus$ .... $\oplus$ -		?	0 if the indices are NOT in the basis
	$p$			
$X_B$	-	0	?	0 if the indices are NOT in the basis
	$p$			
$t_p$	$\oplus$ .... 1 $\oplus$ -		?	

Consider the solution from tableau  $C$  and  $B$  and denote them as  $X_C$  and  $X_B$ . Then we know that  $AX_C = b$ ,  $AX_B = b$ , (here we abuse the notation  $X_B$ , i.e., not the basic solution part.) So  $X_C - X_B \in \text{Null}(A)$ . As  $t_p \in \text{Row}(A)$ , we know that  $\langle X_C - X_B, t_p \rangle = 0$ .

- $\langle X_C, t_p \rangle = \underbrace{\text{non-negative}}_{j < q} + \underbrace{\text{positive}}_{j = q} + \underbrace{0}_{j > q} = \text{positive}$

- Based on the fact that  $X_B, t_p$  are from the same tableau,

$$\langle S_D, t_k \rangle = \underbrace{0}_{j < q, j \neq p} + \underbrace{\text{negative}}_{j=p} + \underbrace{0}_{j=q} + \underbrace{0}_{j \geq q} = \text{negative}$$

Therefore,  $\langle X_C - X_B, t_p \rangle = \text{positive}$ , Hence, contradiction!

**A  $\Rightarrow$  C**

	$j < q$	$j = q$	$j > q$		
$X_A$			0	?	0 if the indices are NOT in the basis
$S_A$			+	?	0 if the indices are in the basis
$X_C$			-	?	0 if the indices are NOT in the basis
$S_C$			0	?	0 if the indices are in the basis

Based on the same reasons used before, we know that

$X_A - X_C \in \text{Null}(A), S_A - S_C \in \text{Row}(A)$ , hence  $\langle X_A - X_C, S_A - S_C \rangle = 0$ .

- As  $S_A, X_C$  from the same tableau, we know that  $\langle X_A, S_A \rangle = 0$ . Similarly,  $\langle X_C, S_C \rangle = 0$
- $\langle X_A, S_C \rangle = \text{non-positive}$ , and  $\langle X_C, S_A \rangle = \text{negative}$

So we know that  $\langle X_A - X_C, S_A - S_C \rangle > 0$ . Hence, contradiction!

To sum up, cycling is impossible during the criss-cross algorithm, hence terminating in finite steps.