

Assignment 2

Last date: February 19, 2025

- Log sum inequality: for any non negatives a_1, \dots, a_n and non negatives b_1, \dots, b_n we have $\sum_{i=1}^n a_i \ln \frac{a_i}{b_i} \geq (\sum_i a_i) \ln \frac{\sum_i a_i}{\sum_i b_i}$.

- In calculation of KL divergence, $0 \ln 0$ is considered as 0.

(1 marks each):

1. Using log sum inequality, show that $KL(P, Q) \geq 0$.
2. Consider an algorithm that always pulls arm 1 (throughout T rounds). Give an instance on which this algorithm achieves an expected regret of 0. Also give an instance on which it achieves an expected regret of T .
3. Which of the following statement(s) is/are correct (Justify briefly - no marks for just writing the answer).
 - (a) The expected regret of Successive Elimination algorithm is $O(\sqrt{KT \log T})$ on all instances.
 - (b) The expected regret of Successive Elimination algorithm can be $\leq (KT \log T)^{1/4}$ on some instances.
 - (c) The expected regret of Successive Elimination algorithm can not be $\leq (KT \log T)^{1/4}$ on all instances
4. Assuming *Good* (as defined in UCB algorithm), show that $\mu_a \leq UCB_a(t)$ for UCB for any t .
5. Assuming *Good* (as defined in Successive Elimination algorithm), prove that the optimal arm will never be eliminated in the Successive Elimination Algorithm.
6. In the class we show a lower bound of $\Omega(\sqrt{KT})$ on the expected regret of any algorithm. But we also see that the UCB can achieve an instance-dependent upper bound of $O(\log T) \sum_{a: \Delta_a > 0} \frac{1}{\Delta_a}$. Explain why they do not contradict each other.
7. Let $P = (p, 1-p)$ and $Q = (q, 1-q)$ be two distributions on two elements. Show

$$d_{tv}(P^{\otimes 2}, Q^{\otimes 2}) \leq 2 d_{tv}(P, Q)$$

8. Show for any n ,

$$KL(P^{\otimes n}, Q^{\otimes n}) = n \cdot KL(P, Q)$$

(2-mark)

Prove for distributions P and Q defined on $[n]$,

$$d_{tv}(P, Q) = \max_{S \subseteq [n]} |P(S) - Q(S)|$$

(1-mark)

In the first assignment, you showed that Testing Coin($\epsilon, 1/5$) problem can be solved in $O(1/\epsilon^2)$ coin tosses. Show that Testing Coin(ϵ, δ) can be solved in $O(\frac{\log 1/\delta}{\epsilon^2})$ coin tosses for any $\delta > 0$.

(3-mark)

In the class, we saw a lower bound of $\frac{(1-2\delta)^2}{\epsilon^2}$ on coin tosses for deterministic algorithms for TestingCoin(ϵ, δ) problem. The above question shows, the upper bound is $O(\frac{\log 1/\delta}{\epsilon^2})$. Clearly, the upper and lower bounds are tight in terms of ϵ but not on δ . Show that the lower bound is also $\Omega(\frac{\log 1/\delta}{\epsilon^2})$ for deterministic algorithms. You can assume $\epsilon \leq 1/3$ (in fact, if you want, you can assume both $\epsilon, \delta \leq c$ for any constant $c < 1$ of your choice).

(Hint: Instead of Pinsker Inequality, use the following stronger bound

$$d_{tv}(P, Q) \leq \sqrt{1 - e^{-KL(P, Q)}}$$

)

(3+3 = 6 marks)

Consider bandit instances in which the mean reward of all arms lie in $[1/2, 1/2 + \gamma]$ for some $\gamma > 0$. Modify Successive Elimination and UCB algorithm to achieve better regret bounds (both instance independent and instance dependent bounds) for these instances. Just to clarify, the algorithm knows the value of γ . Your answer should have both description of the modified algorithms as well as their regret analysis.

(3 marks)

Consider a following problem : input consists of K coins and $\epsilon > 0$. It is promised that either

- all coins are fair or
- there is exactly one coin which is biased and have $\Pr(H) = 1/2 + \epsilon$ and rest other coins are fair.

The goal is to correctly determine the one of the above possibility with at least $4/5$ probability. That is, if the input coins satisfy the first item, algorithm

should return ‘Fair’ with at least $4/5$ probability. On the other hand, if the input coins satisfy the second item, algorithm should return ‘Biased’ with at least $4/5$ probability.

Modify the lower bound proof for Biased Coin Identification problem to show the same lower bound (of $\Omega(K/\epsilon^2)$) for the above problem.

Prove the Pinsker Inequality:

- (2-mark)

Let $P = (p, 1 - p)$ and $Q = (q, 1 - q)$ be two binary distributions (distributions on two elements). Show

$$d_{tv}(P, Q) \leq \sqrt{\frac{1}{2}KL(P, Q)}$$

(Hint: use elementary calculus)

- (1-mark)

Consider two distributions $P = (p_1, \dots, p_n)$ and $Q = (q_1, \dots, q_n)$ on $[n]$. Consider any set $S \subseteq [n]$. Let $P' = (P(S), 1 - P(S))$ and $Q' = (Q(S), 1 - Q(S))$ be binary distributions where recall that $P(S) = \sum_{i \in S} p_i$ and $Q(S) = \sum_{i \in S} q_i$. Show

$$KL(P', Q') \leq KL(P, Q)$$

(Hint: use log sum inequality)

- (2-mark)

Consider two distributions $P = (p_1, \dots, p_n)$ and $Q = (q_1, \dots, q_n)$ on $[n]$. Show that

$$d_{tv}(P, Q) \leq \sqrt{\frac{1}{2}KL(P, Q)}$$

(Proof of $E[\delta] \leq O(\sqrt{K/T})$ for the MOSS algorithm)

Let r_1, \dots, r_T be T independent samples from a distribution D with mean μ (D is supported on $[0, 1]$ so all r_i and μ is in $[0, 1]$). For any $1 \leq x \leq T$, let $\hat{\mu}_x = \frac{\sum_{i=1}^x r_i}{x}$ and

$$I_x = \hat{\mu}_x + \sqrt{\frac{\log^+(\frac{T}{Kx})}{x}}$$

where $\log^+(z) = \max(z, 0)$. Let $\delta = \max(0, \mu - \min_{1 \leq x \leq T} I_x)$

- (3-marks)

For any y , prove that

$$\Pr(\delta \geq y) \leq \frac{K}{T} \cdot O\left(\frac{1}{y^2}\right)$$

(Hint: use Hoeffding's maximal inequality - Suppose X_1, X_2, \dots are iid such that each X_i is in $[0, 1]$ and $E[X_i] = \mu$ then for any $t > 0$ and $m \geq 1$, we have

$$\Pr(\exists 1 \leq r \leq m : \sum_{i=1}^r (\mu - X_i) \geq t) \leq \exp(-\frac{2t^2}{m}).$$

Note that the above inequality is stronger than the one we used throughout the class. To apply the above inequality, t should not depend on r . However, you might come up with an expression where t may depend on r . In such case, use the following trick:

$$\Pr(\exists 1 \leq r \leq m : \sum_{i=1}^r (\mu - X_i) \geq t(r)) \leq \sum_j \Pr(\exists 2^j \leq r \leq 2^{j+1} : \sum_{i=1}^r (\mu - X_i) \geq f(j))$$

where $t(r) \geq f(j)$ for $r \in [2^j, 2^{j+1}]$. Now with some manipulation, you should be able to apply the Hoeffding's maximal inequality.

Additionally, following inequality might be useful: $\sum_{j=1}^{\infty} 2^j \exp(-2^j y^2) = O(\frac{1}{y^2})$

- (2-marks)

Prove that $E[\delta] = O(\sqrt{K/T})$.

(Hint: note that δ is a continuous r.v.. Define a new r.v. β such that β is discrete and $\delta \leq \beta$ and so $E[\delta] \leq E[\beta]$. Obviously, you need to use the result of the first part, i.e., $\Pr(\delta \geq y) \leq \frac{K}{T} \cdot O(\frac{1}{y^2})$ which should be hint on how to define β .)