MATH 623 Homework 2

Yuan Yin

October 17, 2018

Problem 3

5.

The main code:

```
T = 1; K = 1; r = 0.02; delta = 0.3; N = 1000; M = 10^5;
s_bar = K * exp(3 * delta * T^(0.5) / K^(0.5) + abs(r - delta^2 * 0.5 / K) * T);
dt = T / M; ds = s_bar / N; p = zeros(N+1, M+1);
for i = 1 : (N + 1)
   p(i, M + 1) = \max(ds * (i - 1) - K, 0);
end
for i = 1 : (M + 1)
   p(N + 1, i) = s_{bar} - exp(-r * (T - dt * (i - 1))) * K;
   p(1, i) = 0;
end
for i = (M + 1) : -1 : 2
   for j = 2 : N
       alpha1 = 1 - delta^2 * (j - 1) * dt / ds;
       alpha2 = delta^2 * (j - 1) * dt * 0.5 / ds + r * (j - 1) * dt * 0.5;
       alpha3 = delta^2 * (j - 1) * dt * 0.5 / ds - r * (j - 1) * dt * 0.5;
       p(j, i-1) = alpha1 * p(j, i) + alpha2 * p(j+1, i) + alpha3 * p(j-1, i);
   end
end
s = 0 : ds : s_bar;
plot(s, p(:,1))
```

the plot of P(s,0) is as below:

