

The Torus Geometric Refinement Transform (TGRT)

A constant-Jacobian multiresolution transform on the solid ring torus

Abstract. We construct an equal-volume (constant-Jacobian) parameterization of the solid ring torus and build on it a multiresolution transform—the *Torus Geometric Refinement Transform* (TGRT). The TGRT combines dyadic refinement in a radial variable with Fourier analysis in two angular variables, yielding orthonormal bases, Parseval identities, and simple cell-count formulas. The associated discrete exterior calculus (DEC) inherits Kronecker structure, enabling near-diagonalization of the discrete Hodge Laplacian; leakage off the block-diagonal scales with torus thickness $\varepsilon = a/R$. The framework mirrors the SGRT on \mathbb{B}^3 but for the solid torus $\mathcal{T}_{R,a}$, and is designed for fast, stable analysis and PDE discretization.

1 Geometry of the ring torus

Let $R > a > 0$ and define the *ring torus* (solid) as the image of coordinates (ρ, θ, φ) under

$$\Phi_0(\rho, \theta, \varphi) = ((R + \rho \cos \theta) \cos \varphi, (R + \rho \cos \theta) \sin \varphi, \rho \sin \theta), \quad (1)$$

with $\rho \in [0, a]$, $\theta \in [0, 2\pi)$ (poloidal), $\varphi \in [0, 2\pi)$ (toroidal). The standard metric in these orthogonal coordinates is

$$ds^2 = d\rho^2 + \rho^2 d\theta^2 + (R + \rho \cos \theta)^2 d\varphi^2, \quad (2)$$

with volume form

$$dV = \rho (R + \rho \cos \theta) d\rho d\theta d\varphi. \quad (3)$$

The centerline (core circle) is $\mathbf{C}(\varphi) = (R \cos \varphi, R \sin \varphi, 0)$.

2 Equal-volume parameterization and constant Jacobian

Define the radial area variable $u = \frac{1}{2}\rho^2$ so $du = \rho d\rho$, and define a ρ -dependent angular warp $\psi = \psi(\theta; \rho)$ by

$$\psi := \theta + \frac{\rho}{R} \sin \theta \iff d\psi = \frac{R + \rho \cos \theta}{R} d\theta. \quad (4)$$

Theorem 1 (Constant Jacobian map). *For any $R > a > 0$, the change of variables $(\rho, \theta, \varphi) \mapsto (u, \psi, \varphi)$ given by (4) is a C^∞ diffeomorphism of the parameter rectangle, and the induced volume form is constant:*

$$dV = R du d\psi d\varphi. \quad (5)$$

Proof. From (4), $\partial_\theta \psi = 1 + (\rho/R) \cos \theta \in (1 - \rho/R, 1 + \rho/R) \subset (0, 2)$ for all $\rho \in [0, a]$ with $a < R$, hence $\theta \mapsto \psi$ is strictly increasing and 2π -periodic, yielding a diffeomorphism on the circle for each fixed ρ . The Jacobian determinant of $(\rho, \theta, \varphi) \mapsto (u, \psi, \varphi)$ equals $\det(\partial(u, \psi, \varphi)/\partial(\rho, \theta, \varphi)) = \rho \frac{R + \rho \cos \theta}{R}$, so substituting into (3) gives $dV = R du d\psi d\varphi$. \square

Corollary 2 (Equal-volume dyadic bricks). *Uniform splits $u \in [0, U]$ with $U = \frac{1}{2}a^2$, $\psi \in [0, 2\pi)$, $\varphi \in [0, 2\pi)$ into 2^k bins per axis produce $N_k = 8^k$ bricks, each of volume*

$$\Delta V_k = R \frac{U}{2^k} \frac{2\pi}{2^k} \frac{2\pi}{2^k} = \frac{2\pi^2 R a^2}{8^k}. \quad (6)$$

3 The TGRT: spaces, basis, and transform

Let $\mathcal{T}_{R,a} = \Phi_0([0, a] \times \mathbb{T} \times \mathbb{T})$ denote the solid ring torus. Via the diffeomorphism $(u, \psi, \varphi) \mapsto x = \Phi(u, \psi, \varphi) := \Phi_0(\rho(u), \theta(u, \psi), \varphi)$, pull a function $f \in L^2(\mathcal{T}_{R,a})$ back to $g(u, \psi, \varphi) = f(\Phi(u, \psi, \varphi))$. By Theorem 1, the L^2 inner product is flat:

$$\langle f, h \rangle_{L^2(\mathcal{T})} = \int_{\mathcal{T}_{R,a}} f \bar{h} dV = R \int_0^U \int_0^{2\pi} \int_0^{2\pi} g \bar{g}_h du d\psi d\varphi. \quad (7)$$

Scaling and wavelet factors in u ; Fourier in ψ, φ

Fix a dyadic level $j \in \mathbb{Z}_{\geq 0}$ and let $\{I_{j,\ell}\}_{\ell=0}^{2^j-1}$ be the partition of $[0, U]$ into intervals of length $U/2^j$. Define the normalized u -scaling and Haar wavelet functions

$$\phi_{j,\ell}(u) := \frac{1}{\sqrt{U/2^j}} \mathbf{1}_{I_{j,\ell}}(u), \quad (8)$$

$$\psi_{j,\ell}(u) := \frac{1}{\sqrt{U/2^j}} \left(\mathbf{1}_{I_{j+1,2\ell}}(u) - \mathbf{1}_{I_{j+1,2\ell+1}}(u) \right), \quad (9)$$

which are orthonormal in $L^2([0, U], du)$. For angular variables, use the standard Fourier system $\{(2\pi)^{-1/2} e^{im\psi}\}_{m \in \mathbb{Z}}$ and $\{(2\pi)^{-1/2} e^{in\varphi}\}_{n \in \mathbb{Z}}$.

Definition 3 (TGRT basis). For each j, ℓ and integers m, n , define the *TGRT scaling* and *TGRT wavelet* atoms on $\mathcal{T}_{R,a}$ by

$$\Phi_{j,\ell,m,n}^{\text{sc}}(u, \psi, \varphi) := \frac{1}{\sqrt{R}} \phi_{j,\ell}(u) \frac{e^{im\psi}}{\sqrt{2\pi}} \frac{e^{in\varphi}}{\sqrt{2\pi}}, \quad (10)$$

$$\Phi_{j,\ell,m,n}^{\text{w}}(u, \psi, \varphi) := \frac{1}{\sqrt{R}} \psi_{j,\ell}(u) \frac{e^{im\psi}}{\sqrt{2\pi}} \frac{e^{in\varphi}}{\sqrt{2\pi}}. \quad (11)$$

The $1/\sqrt{R}$ factor makes the family orthonormal in the inner product (7).

Proposition 4 (Orthonormality and Parseval). *The TGRT atoms form an orthonormal family in $L^2(\mathcal{T}_{R,a})$. For any $f \in L^2(\mathcal{T}_{R,a})$ with pulled-back g , the TGRT analysis coefficients*

$$\widehat{f}_{j,\ell}^{\text{sc}}[m, n] = \langle g, \Phi_{j,\ell,m,n}^{\text{sc}} \rangle, \quad \widehat{f}_{j,\ell}^{\text{w}}[m, n] = \langle g, \Phi_{j,\ell,m,n}^{\text{w}} \rangle \quad (12)$$

obey the Parseval identity: the sum of squared magnitudes equals $\|f\|_{L^2(\mathcal{T})}^2$.

Proof. Follows immediately from constant measure (7) and orthonormality of Haar in u and complex exponentials in ψ, φ . \square

Synthesis. With the usual multiresolution convention (fix a base scale j_0),

$$g(u, \psi, \varphi) = \sum_{\ell, m, n} \widehat{f}_{j_0, \ell}^{\text{sc}}[m, n] \Phi_{j_0, \ell, m, n}^{\text{sc}} + \sum_{j \geq j_0} \sum_{\ell, m, n} \widehat{f}_{j, \ell}^{\text{w}}[m, n] \Phi_{j, \ell, m, n}^{\text{w}}, \quad (13)$$

with convergence in L^2 . Composing with $\Phi(u, \psi, \varphi)$ yields f on $\mathcal{T}_{R,a}$.

4 Dyadic cells, counts, and boundaries

Let $N = 2^k$. The TGRT grid at level k has N^3 bricks (cells) with constant volume $\Delta V_k = 2\pi^2 R a^2 / 8^k$. The interval edges are

$$u_j = \frac{j}{N} U, \quad \psi_m = \frac{2\pi m}{N}, \quad \varphi_n = \frac{2\pi n}{N}, \quad j, m, n \in \{0, \dots, N\}. \quad (14)$$

The corresponding radii are $\rho_j = \sqrt{2u_j} = a\sqrt{j/N}$. The level-1 bricks are the eight blocks $(I/O) \times (T/B) \times (L/R)$.

5 Discrete Exterior Calculus on the TGRT grid

We outline mass-lumped Hodge stars consistent with the metric induced by (1). Incidence (coboundary) matrices factor as Kronecker products across the three axes, ensuring $d^2 = 0$ exactly on the grid.

Primal edge lengths and dual face areas

Let a brick be indexed by its cell-centered values (u_c, ψ_c, φ_c) , with $\rho_c = \sqrt{2u_c}$ and θ_c determined implicitly by $\psi_c = \theta_c + (\rho_c/R) \sin \theta_c$. Using the embedding (1) and Jacobian $J = R$ from Theorem 1, one finds the following *cell-centered* expressions (exact in the continuous metric, applied in lumped form):

$$\ell_u = \rho_{j+1} - \rho_j, \quad (15)$$

$$\ell_\psi = \frac{\rho_c R}{R + \rho_c \cos \theta_c} \Delta \psi, \quad (16)$$

$$\ell_\varphi = (R + \rho_c \cos \theta_c) \Delta \varphi, \quad (17)$$

$$A_{\psi\varphi} = \rho_c R \Delta \psi \Delta \varphi, \quad (18)$$

$$A_{u\varphi} = \frac{R + \rho_c \cos \theta_c}{\rho_c} \Delta u \Delta \varphi, \quad (19)$$

$$A_{u\psi} = \frac{R}{R + \rho_c \cos \theta_c} \Delta u \Delta \psi. \quad (20)$$

These follow from $J = |\partial_u \Phi \cdot (\partial_\psi \Phi \times \partial_\varphi \Phi)| = R$ and the identities $|\partial_u \Phi| = 1/\rho$, $|\partial_\varphi \Phi| = R + \rho \cos \theta$, $|\partial_\psi \Phi| = \rho R / (R + \rho \cos \theta)$.

Mass-lumped Hodge stars

With (15)–(18), diagonal Hodge stars are

$$(\star_0)_{\text{cell}} = \underbrace{R \Delta u \Delta \psi \Delta \varphi}_{\Delta V}, \quad (21)$$

$$(\star_1)_u = \frac{A_{\psi\varphi}}{\ell_u}, \quad (\star_1)_\psi = \frac{A_{u\varphi}}{\ell_\psi}, \quad (\star_1)_\varphi = \frac{A_{u\psi}}{\ell_\varphi}, \quad (22)$$

$$(\star_2)_{u\psi} = \frac{\ell_\varphi}{A_{u\psi}}, \quad (\star_2)_{u\varphi} = \frac{\ell_\psi}{A_{u\varphi}}, \quad (\star_2)_{\psi\varphi} = \frac{\ell_u}{A_{\psi\varphi}}, \quad (23)$$

$$(\star_3)_{\text{cell}} = \Delta V^{-1}. \quad (24)$$

Thus the DEC Hodge Laplacian on 0-forms is $\Delta_0 = \delta d = \star_0^{-1} d_0^\top \star_1 d_0$; analogous formulas hold for 1- and 2-forms.

6 Laplacian structure and thin-torus asymptotics

For scalar f , the continuous Laplacian in (ρ, θ, φ) is (cf. (2))

$$\Delta f = \partial_{\rho\rho} f + \frac{R + 2\rho \cos \theta}{\rho(R + \rho \cos \theta)} \partial_\rho f + \frac{1}{\rho^2} \partial_{\theta\theta} f - \frac{\sin \theta}{\rho(R + \rho \cos \theta)} \partial_\theta f + \frac{1}{(R + \rho \cos \theta)^2} \partial_{\varphi\varphi} f. \quad (25)$$

Under (u, ψ, φ) , angle coefficients acquire a mild ψ -dependence that vanishes as $\varepsilon = a/R \rightarrow 0$.

Proposition 5 (Mode-coupling scales with thickness). *Let $L_\psi g = \partial_\psi((1 + \varepsilon \cos \psi) \partial_\psi g)$ be the model angular operator at fixed (u, φ) . For input $g(\psi) = e^{im\psi}$, the energy leaked from the main Fourier mode m into sidebands $m \pm 1$ scales as $O(\varepsilon^2)$. Hence, in the TGRT Fourier basis, off-diagonal blocks induced by angular metric variation are $O(\varepsilon)$ in operator norm and energy leakage is $O(\varepsilon^2)$.*

Sketch. Expand $\cos \psi e^{im\psi} = \frac{1}{2}(e^{i(m+1)\psi} + e^{i(m-1)\psi})$; a first-order Born approximation shows coefficients to $m \pm 1$ are $\propto \varepsilon/2$. Parseval then yields energy $\propto (\varepsilon/2)^2$. \square

7 Algorithms and complexity

Forward TGRT (analysis). For a grid with $N_u \times N_\psi \times N_\varphi$ samples:

- 1) Pull back samples to $g(u, \psi, \varphi)$ via the diffeomorphism Φ .
- 2) Apply 1D Haar (or other compactly supported) transform along u (cost $O(N_u)$ per (ψ, φ) slice).
- 3) Apply FFTs in ψ and φ (cost $O(N_\psi \log N_\psi + N_\varphi \log N_\varphi)$ per u -coefficient).

Total cost $O(N_u N_\psi N_\varphi \log(N_\psi N_\varphi))$. Inverse TGRT reverses the steps.

Boundary conditions. Periodic in ψ, φ . In u , one may impose Dirichlet/Neumann/Robin at $u = U$ and regularity at $u = 0$ (axis). For vector fields, impose physical boundary conditions on tangential/normal components on the outer surface $\rho = a$.

8 Notes on novelty and scope

The constant-Jacobian map (4)–(5) packages classical ingredients (tubular coordinates and an equiareal poloidal warp) into a simple diffeomorphism that exactly flattens the measure on the *solid* torus. The TGRT then leverages this to build orthonormal multiresolution bases (Haar \times Fourier \times Fourier) and DEC discretizations with Kronecker structure. The approach extends to anisotropic refinements (non-dyadic) and to the surface torus $\rho = a$ by dropping the u -dimension.

Appendix A: Inverting ψ to θ

For fixed $\rho < R$, $\psi = \theta + (\rho/R) \sin \theta$ is strictly increasing, so each (ρ, ψ) determines a unique θ . Standard Newton iteration with initial guess $\theta^{(0)} = \psi$ converges quadratically:

$$\theta^{(k+1)} = \theta^{(k)} - \frac{\theta^{(k)} - \psi + (\rho/R) \sin \theta^{(k)}}{1 + (\rho/R) \cos \theta^{(k)}}. \quad (26)$$

Appendix B: Surface TGRT (T^2)

At $\rho = a$, the induced metric is $ds^2 = a^2 d\theta^2 + (R + a \cos \theta)^2 d\varphi^2$ and the area form is $dA = a(R + a \cos \theta) d\theta d\varphi$. The map $\psi = \theta + (a/R) \sin \theta$ yields $dA = aR d\psi d\varphi$, i.e. a constant area element, thus a 2D TGRT with Haar in no radial direction and Fourier in both angles (or 2D Haar on (ψ, φ)).

Acknowledgements. This writeup is part of an ongoing program developing geometric refinement transforms (GRT/SGRT/CFRT/DFRT/TGRT) for physics and computation.