Health economic value in the AI era: a framework for intertemporal choice

Richard T. Millian

Hangzhou Normal University

Overtime pay, time saved on meals and faster AI-assisted work offer readily visible gains. Changes in sleep, physical burden and recovery time may be less visible in the same decision. Drawing on health capital, time allocation, health capability and decision analysis, we propose a framework for comparing health-related consequences. Feasible alternatives are compared over a common horizon across resources and time, health and functioning, experience, and capability and agency, before differences are appraised against personal priorities. The framework separates measurement, prediction and appraisal while retaining units, timing and uncertainty. This distinction matters in the AI era: faster tasks need not create more rest, and better assisted performance need not imply improved independent ability. Cases involving single-day and repeated overtime, everyday food choices and digital leisure illustrate the approach. We develop testable propositions concerning the allocation of time gains, persistence of effects and decision quality. The purpose is to clarify the conditions and consequences of choices, rather than assign health a universal price or prescribe a common way of life.




Fig. 1 | A framework for comparing health-related consequences. Feasible alternatives and a common horizon define the comparison. Resources and time, health and functioning, experience, and capabilities and agency remain distinct. Priorities inform appraisal. Connections represent analytical steps, not estimated causal effects.




Fig. 2 | Linking measurement, prediction and appraisal. a, Three kinds of information address the same decision. b, Conditional dominance and trade-offs on qualitative axes without numerical units. Other relevant outcomes and uncertainty may change the ordering.

From economic returns to changes in daily life

Whether overtime pay improves life cannot be judged from the payment alone. Extra income may come from longer daytime work, repeated night shifts or greater efficiency with AI. These arrangements may differ in their implications for sleep, energy and family time. Comparison starts with concrete questions: how much money is gained, where time goes, and whether the person can work, study or care for family the next day. Figure 1 organizes these questions into a common framework; Figure 2 separates measurement, prediction and appraisal.

The starting point is established economics. Grossman's health-capital approach treats health as a stock relevant to both well-being and productive time; Becker's allocation-of-time framework makes non-market time part of economic choice. [1, 2] Marx's discussion of the working day directs attention to bargaining, limits on labour and the reproduction of labour power. [3] These traditions motivate different questions rather than a single interchangeable theory. In particular, health losses cannot simply be appended to Marxian surplus value as an empirically measured quantity without a separate account of exposure, counterfactual outcomes and incidence.

Health capability approaches further distinguish achieved states from the real opportunities people have to maintain health and act on their goals. [4] This matters when an apparent choice cannot be refused without losing housing or essential care. Evidence on health-dependent utility also cautions against assuming that a given monetary increment has the same welfare significance in every health state. [5] Neither point makes income irrelevant. Resources can enable care, security and participation; the question is which consequences a particular increment enables or displaces under actual constraints.

We use health economic value to mean the decision-relevant significance of health and functioning within these trade-offs. It is not synonymous with total welfare, earning capacity or a person's worth. Money, time, experience and agency enter the comparison because they interact with health and carry value of their own. The proposed contribution is an integrative accounting procedure: specify a feasible counterfactual, trace consequences over time, attach evidence and uncertainty, and make the evaluative judgement explicit. A legitimate result can be a preference, a conditional choice or an unresolved comparison.

Table 1 | Minimum comparison record. Proposed fields, not a validated scale. Health capability, ICF, WHO-5, MCDA and decision-quality research inform the distinctions. [4, 6, 7, 8, 9]

Domain	Record	Interpretation
Decision	Feasible options; ability to refuse; common horizon	Distinguish nominal options from real opportunities
Resources and time	Net income change; total work; displaced activities	Record where saved time is used
Health and functioning	Relevant symptoms, function or clinical outcomes	Interpret against baseline and clinical context
Experience	Enjoyment, relief or distress; during and after	Separate immediate responses from sustained change
Capabilities and agency	Independent performance; valued activities; discretion	Include accessibility gains from assistance
Evidence and appraisal	Population; comparator; horizon; uncertainty; goals	Retain unknowns and differences in priorities

From measurement to appraisal

A laboratory report can describe a blood glucose level, but cannot by itself establish whether a choice is worthwhile. Measurement describes the current state; intervention research estimates possible consequences of changing diet; willingness to bear costs in money, time or taste depends on personal priorities. Measurement, prediction and appraisal therefore require different justifications. Reference ranges do not rank human worth, and average study results do not directly specify an individual future.

Monetary valuation can answer defined policy questions, but compensation is not biological restoration. Conversely, irreversibility does not logically make valuation impossible: people still compare prevention, mitigation and foregone opportunities. What fails is the inference that a payment necessarily neutralizes a loss. A stated minimum payment depends on income, alternatives and understanding; accepting dangerous work under severe deprivation does not establish that the health loss is trivial. Behavioural welfare analysis provides reasons to examine the conditions under which choices reveal evaluative preferences. [10]

The framework adapts the transparency of cost–consequence analysis and the explicit criteria of multicriteria decision analysis, while retaining their limitations. [8, 11] It adds no validated universal scale. An established decision guide already elicits options, uncertainty, values and support needs. [12] Our narrower proposal makes time reallocation, outcome trajectories, assisted versus unaided functioning and the distribution of consequences explicit within such support. Any advantage over existing tools must be demonstrated, not inferred from a new label.

Standardized health-status instruments can supply comparable information; preference-based measures such as EQ-5D can support health-economic evaluation when used with the appropriate valuation framework. [13] They do not exhaust autonomy, relationships or personally valued capability. Self-determination theory offers a complementary account of autonomy, competence and relatedness, without implying that every activity has the same meaning for every person. [14] Reading, gaming, exercise and companionship therefore enter as alternatives whose consequences require assessment, rather than predetermined categories of virtue or harm.

Table 1 provides a minimum record. Its purpose is comparability of the decision process, not uniformity of preferences. When meaningful consequences diverge, the analyst should show the conflict and examine sensitivity to priorities. A single index is optional only when its scales, weights, uncertainty and double-counting risks are defensible; declining to aggregate is often the more informative result.




Figure 3 | Task savings and the allocation of a day. a, The accounting identity includes prompting, verification and rework in T1, additional tasks in A and other work-time changes in B. **b,** Hypothetical 24-hour budgets with an 8-hour baseline workday, a 4-hour task saving and B = 0. Non-work time is not assigned a health effect.

Intertemporal comparison and time constraints

Let $x_i(a;\tau)$ denote person i's consequence profile under feasible option a at time τ. Components retain their own units: disposable resources, time allocation, health and functioning, experience and capabilities. For components with a meaningful numerical difference, define:

$$\Delta x_i(a,b;\tau) = x_i(a;\tau) - x_i(b;\tau). \quad (1)$$

This is a contrast, not a causal prediction model. Categorical changes remain descriptive. Every estimated component requires a population, comparator, horizon, source and uncertainty statement. Timing may require a trajectory rather than an endpoint. Let z_j denote an appropriately oriented criterion, with larger values preferred within its applicable range. Conditional dominance requires:

$$\Delta z_j \geq 0\ (\forall j);\ \Delta z_k > 0\ (\exists k). \quad (2)$$

This ordering applies only if the criteria cover the relevant consequences and uncertainty does not overturn it. Many biomarkers are not monotonic, and omitted consequences can reverse an apparent ranking. Where trade-offs remain, personal appraisal is necessary; no universal exchange rate follows from equations (1)–(2).

For AI-assisted work, let S = T0 − T1 be time saved on a fixed task; T1 includes prompting, verification and rework. Let A be time added through additional tasks and B all other net changes in work time, defined to avoid overlap. Then:

$$\Delta W = -S + A + B. \quad (3)$$

Figure 3 illustrates the identity. Saving four task hours produces four fewer work hours if A = B = 0, one fewer if A = 3 and B = 0, and one more if A = 5 and B = 0. These are hypothetical allocations, not estimated AI effects. Even a fall in total work does not determine health: the activities replacing work, discretion over them and their subsequent consequences remain to be measured.

Table 2 | Different levels of AI outcomes. Studies illustrate task efficiency, experience and judgement; their designs and endpoints differ.

Setting and design	Main finding	Interpretive scope
Professional writing Randomized; 453 [15]	Faster task completion and better output quality	Full-day allocation and health not measured
Customer support Staggered rollout; 5,172 [16]	About 15% higher resolutions per hour; outcome availability varies	Causal assumptions; who receives saved time
Companionship Seven-day Study 3 [17]	Immediate loneliness relief; persistent pre-interaction benefit not demonstrated	Differential attrition; control not time-matched
Medical scenarios 1,298 people [18]	Control versus pooled AI: identification OR 1.76 (95% CI 1.45–2.13)	Scenario performance, not clinical morbidity

Three levels of AI benefit

AI can expand productive capacity. In a randomized study of 453 professionals, Noy and Zhang found improvements in writing-task speed and quality. [15] In a field study covering 5,172 customer-support agents, Brynjolfsson and colleagues used staggered deployment and difference-in-differences analysis; their preferred estimate implied about a 15% increase in issues resolved per hour. [16] This was not a randomized trial, and the field outcome was throughput, not health. Together these studies support productivity gains while leaving open who receives the time or income dividend. The study lacked compensation data, so productivity gains cannot directly be interpreted as increased worker income. [16]

The relevant distinction is also between performance with assistance and capability retained without it. Bastani and colleagues' educational experiment illustrates why an evaluation should include an unaided assessment rather than infer learning from assisted practice alone. [19] This does not establish that AI generally erodes cognition. It specifies a measurement requirement: immediate performance, transfer to unfamiliar tasks and independent performance answer different questions. Accessibility and supported functioning can be valuable even where unaided performance is unchanged.

AI companionship raises a temporal distinction. In the seven-day third study reported by De Freitas and colleagues, loneliness relief was most clearly immediate; a persistent improvement in pre-interaction loneliness was not demonstrated. [17] Differential attrition and a non-time-matched control limit interpretation. This is evidence for examining when benefit occurs, not grounds for dismissing companionship or diagnosing dependence. For short-form content and virtual interaction, useful records would include sleep displacement, enjoyment, relationships and the ability to disengage; content type or duration alone does not establish harm.

Information accuracy and user benefit also differ. Bean and colleagues found that people working through hypothetical medical scenarios did not reproduce the models' standalone performance; Figure 5d retains the reported control-versus-AI contrast. [18] The outcome was scenario interpretation, not clinical disease. Earlier Facebook-deactivation research likewise shows the value of measuring consequences beyond platform use, although social-media findings cannot simply be transferred to generative AI. [20] Health-literate AI already emphasizes comprehension, agency, accountability and proportionality. [21] Our proposal complements that agenda by specifying the consequences and time horizons to compare, rather than claiming these concerns as new.

a Locate each measure in the proposed pathway



b Select measures by construct and timing

Domain	Candidate measure	Interpretation
Metabolic state	Glucose or lipids	Clinical role and baseline
Physical function	Strength; activity limitations	Ability to perform daily activities
Vision and skin	Symptoms; discomfort	Functional and experiential impact
Cognition	Assisted / unaided task	Performance and task transfer
Mental well-being	WHO-5	Preceding two weeks
Experience and agency	Enjoyment; distress; discretion	During use and subsequent follow-up

Choose a small relevant set; do not sum different points on the same pathway.

Fig. 4 | Measurement levels and their decision relevance. The upper pathway separates exposure, state and consequences in daily life; arrows indicate relationships to investigate. Candidate measures below are selected for the decision and its timing. ICF links functioning to context; WHO-5 covers mental well-being over the preceding two weeks. [6, 7]

Choosing measures for the decision

Measures should follow the decision. A dinner comparison may require price, preparation time and satiety; night-work comparisons also need sleep, functioning and recovery. Select relevant measures rather than every available test, after establishing feasible alternatives, a common horizon and baseline conditions. Figure 4 organizes these measurement levels.

Separate exposures, states and consequences. Sleep opportunity, sitting time, diet and screen use describe behaviours or environments. Symptoms, laboratory measures and task performance describe selected states. Their significance depends on daily functioning, experience and valued activities. The ICF helps situate functioning in context. [6] Adding sedentary time, a correlated biomarker and a downstream disease risk into a single score can count the same pathway repeatedly. Candidate causal links should therefore be specified before aggregation or mediation analysis.

Glucose, lipids, blood pressure, strength, vision and skin symptoms answer different questions; blood quality is too vague. Changes require interpretation against baseline, measurement conditions and appropriate ranges. Appearance may affect experience but cannot establish internal organ status. Physical–psychological relationships require measurement, and differences in health must not become a ranking of human dignity or worth.

Experience and functioning should be observed at appropriate times. WHO-5 records mental well-being over the preceding two weeks; it is not a diagnostic test or an instantaneous response measure. [7] A brief attention task, an activity diary and a symptom report likewise capture different constructs. Selection should consider reliability, responsiveness, language, accessibility and burden. Repeated measurement may distinguish transient discomfort from a sustained trajectory, but collecting more data is not automatically better and can increase anxiety or exclusion.

Finally, attach uncertainty to each consequential claim. Distinguish observed findings from projections and unknowns; an unmeasured effect is not zero. Report absolute risks when defensible baseline estimates and applicable models exist, rather than translating relative effects into unsupported personal forecasts. Preserve the person's priorities alongside the evidence, and revisit the comparison if circumstances or outcomes change.




Figure 5 | Selected estimates with their original uncertainty. a, Stroke incidence from seven cohorts, follow-up 1–20 years. [22] **b,** Energy intake in a 20-person inpatient crossover diet trial. [23] **c,** Energy intake change versus control in a sleep-extension trial; 81 randomized, 80 in the reported analysis set. [24] **d,** Relevant-condition identification in hypothetical medical scenarios: control versus pooled AI assistance. [18] Bars denote 95% CIs in a, c and d; ±1 SE in b. Ratio axes are logarithmic. RR and OR are distinct; neither is an absolute probability change. No effects are pooled or converted to a personal health price.

Empirical estimates and their scope

Figure 5 preserves four different estimands. Long working hours were associated with stroke incidence in a cohort meta-analysis; this does not estimate the effect of one overtime week. [22] Hall and colleagues' 20-person inpatient crossover experiment found greater energy intake on the ultra-processed diet, whereas Tasali and colleagues' sleep-extension trial found lower estimated energy intake relative to control. [23, 24] The former interval is one standard error, not a 95% confidence interval. Bean and colleagues' odds ratio concerns identification in medical scenarios, not the probability of illness. [18]

These findings make particular consequences credible enough to investigate. They do not validate the entire framework, establish a universal lifestyle hierarchy or provide coefficients for pricing an individual's health. Transporting an effect requires attention to population, exposure, comparator, outcome and follow-up. The figure therefore keeps study designs, units and uncertainty visible instead of pooling unlike outcomes.

Trade-offs in everyday settings



Fig. 6 | Daily income and overnight schedules. Conceptual illustration. With CNY 200 base pay and additional net supplements of CNY 400 or 1,000, totals are CNY 600 or 1,200. The image does not depict study participants or measured health outcomes.

Case 1: daily pay and overnight supplements. Suppose ordinary daily pay is CNY 200, with an additional CNY 400 for working until 02:00 or CNY 1,000 for working through the night. Assuming all amounts are net of tax and ordinary pay is unchanged, daily income is CNY 200, 600 or 1,200 (Fig. 6). If CNY 400 and 1,000 instead denote total daily pay, the increments are CNY 200 and 800. Distinguishing base pay, supplements and total income prevents an initial accounting error.

The schedule must then be specified: when work starts, when the person must rise the next day, whether another shift follows, and whether commuting or care duties can change. Finishing at 02:00 with a day off is a different option from finishing at 02:00 before an early shift. An end time alone does not establish sleep duration, and comparison should extend beyond the work episode to subsequent functioning and recovery.

Whether the person gains also depends on what the money enables. Paying rent or care costs may relieve an immediate difficulty; fatigue, restricted activities and other burdens still belong in the comparison. Conversely, the absence of obvious discomfort does not establish an absence of effects. Confirmed income changes, functioning that needs observation and unknown longer-term consequences can be listed separately, allowing the person to identify losses they regard as unacceptable.

Negotiable arrangements may enlarge the choice set: fewer consecutive night shifts, protected time off or the ability to stop. Including them turns a question about the price of health into a comparison of ways to meet income needs while preserving health and valued activities. These are options to evaluate; their effects require evidence.

Case 2 Comparing a week with a month




Figure 7 | Align the overtime alternatives. Both options cover 28 days; supplements are assumed net and ordinary earnings equal. Option B adds CNY 90,000 and 21 overtime days. The timeline and arithmetic are hypothetical. Exposure, baseline state, recovery and later outcomes require additional evidence before an overall appraisal.

Consider a hypothetical 28-day window. Option A offers a net supplement of CNY 10,000 for seven overtime days; option B offers CNY 100,000 for 28. Holding ordinary earnings equal, $\Delta Y = 100{,}000 - 10{,}000 = 90{,}000$ CNY, alongside 21 additional overtime days. Figure 7 aligns the denominators. It does not assign the remaining days in option A a health value automatically. Their activities, recovery opportunities and effects on family members must be recorded; effects extending beyond day 28 require a longer outcome horizon.

A disciplined appraisal asks about shift duration and timing, baseline symptoms, the ability to stop, essential financial needs and the next-best feasible option. The same schedule may be evaluated differently under different constraints without making its health consequences unreal. The framework cannot determine that CNY 100,000 is sufficient compensation. It can identify what is known, what remains uncertain and which conflicts drive the decision. A legally or contractually specified payment is an allocation of responsibility; it is not evidence that lost functioning has been restored.

Sleep restriction also shows why subjective adaptation and objective function should be distinguished. In Van Dongen and colleagues' laboratory programme, 35 adults were randomized to 4, 6 or 8 hours in bed for 14 nights; a separate total-deprivation experiment involved 13 adults. Performance deficits accumulated in the 4- and 6-hour groups, while reported sleepiness did not track their full progression. [25] These functional findings do not establish permanent neuronal destruction after one late night. Recovery and persistence require their own evidence.

Case 3 Beyond the price of dinner



Fig. 8 | Resources and circumstances behind a meal. Conceptual illustration. Purchased and home-prepared meals require appraisal of the actual food, time, labour and availability; health outcomes are not assigned by the image.

Comparing a convenient purchased dinner with a meal prepared at home requires more than ingredients or price. Suppose the alternatives are maintained for four weeks. Food expenditure, shopping, cooking and cleaning time, taste and feasibility all enter the comparison (Fig. 8). A person with a kitchen and predictable hours faces different options from a shift worker with limited cooking facilities.

Junk food is an everyday label rather than a sufficiently defined research exposure. Food composition, processing category, intake and duration need specification: takeaway is not synonymous with ultra-processed food, nor home cooking with a balanced diet. Hall and colleagues observed an energy-intake difference, but no significant between-diet differences in short-term glucose tolerance or continuous glucose measures. This neither establishes long-term equivalence nor supports deterioration in every internal measure. [23]

The economic point is that preparing a meal also uses time and labour. Who shops, cooks and cleans within the household belongs in the account, alongside the time gained from convenience. A practical comparison can begin with feasible changes in available meal combinations or preparation burden, followed by observation of eating experience and relevant outcomes, rather than requiring an idealized menu in advance.

Case 4 Screens, companionship and an evening



Fig. 9 | Leisure activities and what follows. Conceptual illustration. Digital interaction, reading, movement and social contact are alternatives to compare in terms of displacement, persistence of experience and agency.

Suppose two hours after dinner are discretionary. Short dramas or games, AI conversation, reading, walking and meeting friends are candidate activities (Fig. 9). Appraisal first asks what each displaces. Gaming may displace intended sleep for one person and otherwise unstimulating time alone for another. Online and offline activities cannot be ranked by their labels alone, and leisure need not justify every minute through increased productivity.

Observation can extend from the evening to the next day and subsequent weeks: was the activity enjoyable, did satisfaction persist, could the person stop as planned, and could they carry out valued activities the next day? Attention needs a specified construct, such as sustained attention, switching or memory. Patience likewise needs a defined behaviour; viewing duration alone cannot diagnose cognitive decline. These questions organize observation rather than establish causal pathways.

AI companionship also calls for separating immediate support from persistent change. Research has recorded post-interaction loneliness relief, while follow-up and comparator design constrain conclusions about persistence. [17] A case comparison can examine whether use supplements existing support, accompanies later sleep, and remains under the user's control. Appraisal can recognize enjoyment and support as valuable while preserving scope to change use.

The same principle applies to AI food or medical advice. A credible-sounding answer and an appropriate user decision are different outcomes; medical-scenario experiments have demonstrated a gap between models acting alone and human–AI interactions. [18] Evaluation should examine understanding, recognition of applicability and appropriate action, rather than treating verbal fluency as effectiveness.




Fig. 10 | Two research questions and their evaluation paths. Mechanism research explains how outcomes arise; tool evaluation tests the added comparison module. An active comparator gives both groups the same facts and established guide. The proposed design is not a completed trial.

Table 3 | Testable propositions and disconfirming observations. Each comparison requires prespecified estimands and outcomes.

Proposition	Key comparison	Potentially disconfirming result
Discretion affects allocation of AI time savings	Similar task gains; credible differences in control over workload	No meaningful allocation difference, with sufficiently precise estimates
Immediate relief and sustained benefit may diverge	Matched comparator; repeated pre-use, post-use and follow-up measures	Stable pre-use improvement comparable to immediate relief
The module improves decision quality	Same guide, with or without the added module	No gain in understanding or transfer, or burden exceeds benefit

Testable propositions and evaluation

Validation involves two questions: which mechanisms change health-related outcomes, and whether the framework improves informed choice. The first provides evidence for its content; the second determines its practical value. Figure 10 separates these research questions, and Table 3 specifies three propositions and observations that could weaken them.

An AI workplace study could combine task logs with whole-day time diaries and repeated outcomes, while measuring authority over workload and schedules. A randomized organizational policy or a credible natural experiment would strengthen causal inference about who captures saved time. A simple association between AI use and well-being would remain vulnerable to job selection, workload, prior health and reverse causation. Mechanism analysis should be prespecified; controlling mechanically for post-treatment time use can obscure the total effect it seeks to explain.

For the decision tool, an active comparator is essential. Both groups should receive the same factual evidence and an established guide; the intervention adds the explicit time, consequence and uncertainty module (Fig. 10). Decision-quality research and the Cochrane review of patient decision aids inform the comparator and outcome selection. [9, 26] The primary outcome should be specified before recruitment, with independent scoring of understanding rather than agreement with the author's preferred lifestyle. Transfer to an unfamiliar decision, values–choice concordance and calibrated uncertainty can be secondary outcomes.

Evaluation must also capture completion time, attrition, anxiety, accessibility and the experience of people who stop using the tool. Repeated use may introduce learning, fatigue or changes in the decision itself. A preregistered protocol should specify the target population, effect of interest, analysis of missingness, multiplicity and sample-size justification. Interviews can explain comprehension or disagreement but cannot replace causal estimates. If the added module offers no meaningful benefit over the guide, or its burden outweighs benefit, it should be simplified or abandoned; a shift toward supposedly healthier choices alone would not demonstrate success.




Fig. 11 | Conditions of choice and distribution of consequences. Workplace, platform and household conditions shape feasible options. Income, time and care burdens may accrue to different people. Arrows organize evaluation questions rather than depict estimated causal effects.

Institutions, distribution and research priorities

A consequence profile would be incomplete if it treated constrained behaviour as a failure of individual discipline. Workplace rules, platform design, housing, food access and caring responsibilities shape both available options and the capacity to act on a preference. Figure 11 places these conditions upstream of appraisal. Recording that someone would prefer rest is insufficient if income insecurity or workload rules prevent it. Decision support can reveal such a conflict; it cannot itself redistribute bargaining power or provide the missing option.

The perspective of the account must be stated. A worker's net payment, an employer's productivity, a family's unpaid care and public expenditure are distinct entries. A payment can compensate a recipient while representing a transfer between parties rather than a net social resource gain. Time saved for one person may be shifted to another through monitoring, correction or care. Comparing individual, household and societal accounts can expose these differences without assuming that their interests can be aggregated by a single unexamined weight.

Institutions can also use the framework to ask concrete questions: does a productivity policy return time to workers; does a platform permit meaningful disengagement; and does an intervention remain feasible for people with limited money, mobility or digital access? These are evaluative questions, not established effects of particular policies. The same information should not become a surveillance score used to penalize workers, deny opportunities or blame people for constrained choices. Purpose limitation and proportionate collection are part of making the proposed support usable.

The central claim is modest in form but consequential in practice: a gain is not fully characterized until the health-relevant consequences, their timing and their distribution are visible. AI strengthens the need for this accounting because task performance, independent capability and lived benefit can move differently. A common comparison procedure can make these divergences discussable without imposing a common preference. Its scientific value will depend on whether it improves explanation, prediction or decision quality beyond the methods on which it builds.

This conceptual Perspective integrates economic theory, measurement approaches and selected empirical studies through narrative synthesis. The framework remains to be validated. Its breadth may increase burden, population effects may not transfer to individuals, and preferences may change with circumstances. The next step is context-specific feasibility research followed by comparison with established decision support, assessing whether any gains in understanding and choice persist.

References

1. Grossman, M. On the concept of health capital and the demand for health. Journal of Political Economy 80, 223–255 (1972). https://doi.org/10.1086/259880

2. Becker, G. S. A theory of the allocation of time. The Economic Journal 75, 493–517 (1965). https://doi.org/10.2307/2228949

3. Marx, K. The working-day. Capital, Vol. I, ch. 10. Marxists Internet Archive (n.d.). https://www.marxists.org/archive/marx/works/1867-c1/ch10.htm

4. Ruger, J. P. Health capability: conceptualization and operationalization. American Journal of Public Health 100, 41–49 (2010). https://doi.org/10.2105/AJPH.2008.143651

5. Finkelstein, A., Luttmer, E. F. P. & Notowidigdo, M. J. What good is wealth without health? The effect of health on the marginal utility of consumption. Journal of the European Economic Association 11 (Suppl. 1), 221–258 (2013). https://doi.org/10.1111/j.1542-4774.2012.01101.x

6. World Health Organization. International Classification of Functioning, Disability and Health. WHO (n.d.). https://www.who.int/standards/classifications/international-classification-of-functioning-disability-and-health

7. World Health Organization. The World Health Organization-Five Well-Being Index (WHO-5). WHO, WHO/UCN/MSD/MHE/2024.01 (2024). https://www.who.int/publications/m/item/who-ucn-msd-mhe-2024.01

8. Thokala, P. et al. Multiple criteria decision analysis for health care decision making—an introduction: Report 1 of the ISPOR MCDA Emerging Good Practices Task Force. Value in Health 19, 1–13 (2016). https://doi.org/10.1016/j.jval.2015.12.003

9. Sepucha, K. R. et al. Establishing the effectiveness of patient decision aids: key constructs and measurement instruments. BMC Medical Informatics and Decision Making 13 (Suppl. 2), S12 (2013). https://doi.org/10.1186/1472-6947-13-S2-S12

10. Bernheim, B. D. & Rangel, A. Beyond revealed preference: choice-theoretic foundations for behavioral welfare economics. The Quarterly Journal of Economics 124, 51–104 (2009). https://doi.org/10.1162/qjec.2009.124.1.51

11. Office for Health Improvement and Disparities. Cost consequence analysis: health economic studies. GOV.UK (2020). https://www.gov.uk/guidance/cost-consequence-analysis-health-economic-studies

12. Ottawa Hospital Research Institute. Ottawa Personal Decision Guide. Patient Decision Aids (2025). https://decisionaid.ohri.ca/decguide.html

13. EuroQol Research Foundation. What is the difference between the EQ-5D descriptive system, the EQ VAS and the EQ-5D index values? EuroQol (2023). https://euroqol.org/faq/what-is-the-difference-between-the-eq-5d-5l-descriptive-system-the-eq-vas-and-the-eq-5d-index-values/

14. Ryan, R. M. & Deci, E. L. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. American Psychologist 55, 68–78 (2000). https://doi.org/10.1037/0003-066X.55.1.68

15. Noy, S. & Zhang, W. Experimental evidence on the productivity effects of generative artificial intelligence. Science 381, 187–192 (2023). https://doi.org/10.1126/science.adh2586

16. Brynjolfsson, E., Li, D. & Raymond, L. Generative AI at work. The Quarterly Journal of Economics 140, 889–942 (2025). https://doi.org/10.1093/qje/qjae044

17. De Freitas, J., Oğuz-Uğuralp, Z., Uğuralp, A. K. & Puntoni, S. AI companions reduce loneliness. Journal of Consumer Research 52, 1126–1148 (2026). https://doi.org/10.1093/jcr/ucaf040

18. Bean, A. M. et al. Reliability of LLMs as medical assistants for the general public: a randomized preregistered study. Nature Medicine 32, 609–615 (2026). https://doi.org/10.1038/s41591-025-04074-y

19. Bastani, H. et al. Generative AI without guardrails can harm learning: Evidence from high school mathematics. Proceedings of the National Academy of Sciences 122, e2422633122 (2025). https://doi.org/10.1073/pnas.2422633122

20. Allcott, H., Braghieri, L., Eichmeyer, S. & Gentzkow, M. The welfare effects of social media. American Economic Review 110, 629–676 (2020). https://doi.org/10.1257/aer.20190658

21. Ivic, R. K., Ratzan, S. C. & Parker, R. M. Building health-literate artificial intelligence. Nature Human Behaviour, Advance online publication (2026). https://doi.org/10.1038/s41562-026-02595-1

22. Descatha, A. et al. The effect of exposure to long working hours on stroke: A systematic review and meta-analysis from the WHO/ILO Joint Estimates of the Work-related Burden of Disease and Injury. Environment International 142, 105746 (2020). https://doi.org/10.1016/j.envint.2020.105746

23. Hall, K. D. et al. Ultra-processed diets cause excess calorie intake and weight gain: An inpatient randomized controlled trial of ad libitum food intake. Cell Metabolism 30, 67–77.e3 (2019). https://doi.org/10.1016/j.cmet.2019.05.008

24. Tasali, E. et al. Effect of sleep extension on objectively assessed energy intake among adults with overweight in real-life settings: A randomized clinical trial. JAMA Internal Medicine 182, 365–374 (2022). https://doi.org/10.1001/jamainternmed.2021.8098

25. Van Dongen, H. P. A., Maislin, G., Mullington, J. M. & Dinges, D. F. The cumulative cost of additional wakefulness: dose-response effects on neurobehavioral functions and sleep physiology from chronic sleep restriction and total sleep deprivation. Sleep 26, 117–126 (2003). https://doi.org/10.1093/sleep/26.2.117

26. Stacey, D. et al. Decision aids for people facing health treatment or screening decisions. Cochrane Database of Systematic Reviews 1, CD001431 (2024). https://doi.org/10.1002/14651858.CD001431.pub6