

First Exam — Lessons 0 & 1 (Version A)

Time allowed: 60 minutes

Resources: Personal notes only. Electronic devices, calculators, and statistical tables are not permitted.

Instructions: Answer all four questions. Show algebraic steps and clearly label each part. Unless stated otherwise, provide exact values (fractions, radicals, exponentials). Partial credit is available when reasoning is clearly explained.

Question	Topic	Points
1	Probability structures and independence	25
2	Bayes' rule and conditional reasoning	25
3	Exponential model derivations	25
4	Interpreting summaries and convergence	25
Total		100

Question 1 — Probability Foundations (25 points)

A coin is tossed twice. Throughout parts (a)–(c), assume the coin is fair.

- (6 pts) Specify a probability space $(\Omega, \mathcal{F}, P)$ for this experiment. Clearly list the sample space, a natural σ -algebra, and the probability measure.
- (6 pts) Let A be the event “at least one head appears” and B the event “exactly one head appears.” Compute $P(A)$, $P(B)$, and $P(A \cap B)$.
- (5 pts) Are A and B independent? Justify your answer using the formal definition.
- (8 pts) Now suppose the coin has probability p of landing heads, where $0 < p < 1$. With the same definitions of A and B , derive expressions for $P_p(A)$, $P_p(B)$, and $P_p(A \cap B)$. For which value(s) of p are A and B independent? Provide algebraic reasoning.

Question 2 — Bayes' Rule in Component Testing (25 points)

A factory purchases circuit boards from two suppliers. Boards from S_A account for $\frac{3}{5}$ of the total, and boards from S_B account for the remaining $\frac{2}{5}$. Historical quality data show that $\frac{1}{10}$ of boards from S_A are defective, while $\frac{1}{20}$ of boards from S_B are defective. Each board is screened by a test that returns “positive” for a defective board with probability $\frac{9}{10}$, and returns “positive” for a non-defective board with probability $\frac{1}{20}$.

Recall: sensitivity = $P(\text{Positive} \mid \text{Defective})$ and specificity = $P(\text{Negative} \mid \text{Non-defective})$. The false positive rate is $1 - \text{specificity} = P(\text{Positive} \mid \text{Non-defective})$.

- (7 pts) Compute the probability that a randomly selected board produces a positive test result.

2. (8 pts) Given a positive test result, determine the probability that the board is defective. Express your answer as a reduced fraction.
 3. (6 pts) Given a negative test result, determine the probability that the board is defective. Express your answer as a reduced fraction.
 4. (4 pts) Let $\pi = P(\text{Defective})$, $s = P(\text{Positive} \mid \text{Defective})$, and $f = P(\text{Positive} \mid \text{Non-defective})$. Derive a formula for $P(\text{Defective} \mid \text{Positive})$ in terms of π , s , and f , and briefly state how decreasing f affects this probability.
-

Question 3 — Exponential Waiting-Time Model (25 points)

Let T be the waiting time (in hours) until the next service request arrives, and suppose T follows an exponential distribution with rate $\lambda = \frac{1}{4}$.

Recall: The probability density function of an exponential random variable with rate λ is given by

$$f_T(t) = \begin{cases} \lambda e^{-\lambda t}, & t \geq 0, \\ 0, & t < 0. \end{cases}$$

1. (5 pts) Derive the cumulative distribution function $F_T(t)$.
 2. (5 pts) Using your result from part (1), compute $P(1 \leq T \leq 4)$ and leave the answer in exponential form.
 3. (9 pts) Compute $E[T]$ and $\text{Var}(T)$ directly from integrals. Show the essential integration steps leading to exact values.
 4. (6 pts) Define $Y = 3T + 2$. Compute $E[Y]$ and $\text{Var}(Y)$ using properties of expectation and variance. Provide exact values.
-

Question 4 — Interpreting Summaries and Convergence (25 points)

Table 1 summarises the results of a simulation: for each sample size n , 1,000 independent samples of size n were drawn from an exponential distribution with mean 4, and the sample mean was recorded for each run.

Table 1: Behaviour of Sample Means

Sample size n	Average of the 1,000 sample means	SD of the 1,000 sample means
5	4.08	1.80
20	4.02	0.90
80	4.01	0.44

Another analysis compared a standardised dataset (mean 0, variance 1) to the standard normal distribution. Selected points from the QQ-plot are listed below.

Answer the following:

Table 2: Selected QQ-Plot Quantiles

Normal quantile	Sample quantile
-2.0	-3.1
-1.0	-1.6
0.0	0.2
1.0	1.9
2.0	3.4

1. (9 pts) Use Table 1 to explain how the Law of Large Numbers manifests in this simulation. Refer explicitly to the numerical values.
2. (8 pts) Still relying on Table 1, discuss how the variability of the sample mean changes with n and connect your explanation to the Central Limit Theorem.
3. (8 pts) Interpret Table 2. Describe the shape of the corresponding QQ-plot, assess the plausibility of a normal model, and recommend a modelling or diagnostic action before relying on normal-based methods.

End of exam. Ensure answers are clearly labelled and presented legibly.