Does Electrocution Improve Roller-Crimper Cover Crop Termination?

Sougata Roy[1*], Ryan Hamberg[1], Muthukumar Bagavathiannan[1]

*[1]Department of Soil and Crop Sciences, Texas A&M University, College Station, TX · *Contact: sroy_agronomy@tamu.edu; muthu@tamu.edu*

Poster board 1251

Introduction

- Organic growers cannot use herbicides, so cover crops must be killed mechanically; incomplete kill causes regrowth, weed pressure and planting problems **(Creamer and Dabney 2002)**.
- The roller crimper (RC) kills cereal rye reliably only after anthesis, and surviving tillers regrow **(Mirsky et al. 2009; Kornecki 2015)**.
- Electrical weed control (EWC) kills plants with high-voltage current **(Slaven et al. 2023)**, but one pass only top-killed rye **(Sullivan 2025)**.
- **Gap:** whether EWC can kill what the crimper leaves alive, and whether pass order matters, is unknown.

Objective and Hypothesis

Objective: Determine whether EWC, before or after roller crimping, improves cereal rye termination over either tool alone.

Hypothesis: EWC + RC terminates rye better than either tool alone, in either order: crimping crushes the stems, and current kills the tillers the crimper misses.

Materials and Methods

Experimental design

- College Station, TX, 2026; RCBD, 4 reps
- Cereal rye (*Secale cereale* L.), Zadoks 85
- 30 × 10 ft plots; same-day passes (Fig. 1)

5 TREATMENTS
Control · EWC · RC
RC→EWC · EWC→RC

COVER CROP
Cereal rye, soft dough
≈ 615 g m^{-2} dry biomass

EWC: ZASSO EH60
2.5 m, 3.2 km h^{-1}
One pass

ROLLER CRIMPER
Commercial, 3 m wide
One pass

Measurements

- Termination, survival (visual): 3, 7, 15, 28 DAT
- Biomass (0, 28 DAT); soil moisture (HydroGO sensor)

EWC energy dose

- P = 3 × 255 V × 40–45 A ≈ 31–34 kW
- **E/A = P / (v · w) ≈ 13.8–15.5 kJ m^{-2}**

Data analysis

- RCBD ANOVA on raw %; Tukey's HSD (α = 0.05)
- Python 3.11, statsmodels 0.15 **(Seabold and Perktold 2010)**





Fig 1. EWC (Zasso EH60) and roller crimper in cereal rye.

Results and Discussion



Fig 2. Plots 15 days after treatment (DAT).




Fig 3. Visual termination score at 28 DAT (0 = none, 10 = complete; mean ± SE, n = 4); P < 0.001. Letters: Tukey's HSD (α = 0.05).

Key results

- **Termination:** at 28 DAT, EWC→RC (9.0) and RC→EWC (8.0) beat RC (6.5) and EWC (2.0) (Fig. 3); order did not differ (P = 0.08), consistent with crimping + flaming **(Frasconi et al. 2019)**
- **Regrowth:** 79% of the rye survived EWC alone and 34% survived RC, but only 6–11% survived the sequences (Fig. 5), consistent with EWC top-kill of rye **(Sullivan 2025)**
- **Biomass and soil water:** the sequences cut biomass by 57–62% (Fig. 6); crimped plots held 27–28% soil moisture vs 17% in the control, as with rolled rye mulch **(Kornecki et al. 2012)**



Fig 4. Proposed mechanism (conceptual, not measured): EWC cannot terminate rye completely and the crimper leaves some tillers standing; together, few tillers survive.

Conclusions

- **Hypothesis supported:** in either order, one same-day EWC pass raised the RC termination score from 6.5 to 8.0–9.0 (of 10) and cut surviving rye from 34% to 6–11%.
- **EWC alone failed** even at ≈ 10× the dose that killed an oat–clover mix **(WSU 2025)**, likely because mature, lignified rye (Zadoks 85) conducts current poorly.
- **For growers:** one extra same-day pass finishes a crimper kill without herbicides; tested at one site in one year.

TAKEAWAY

Add a same-day EWC pass to roller crimping, in either order: termination score rose to 8–9 of 10 and surviving rye fell to 6–11%.




Fig 5. Surviving stand at 15 DAT (mean ± SE, n = 4); P < 0.001. Letters: Tukey's HSD (α = 0.05).

What it shows: share of rye plants still alive and regrowing 15 days after termination; lower means a more complete kill.




Fig 6. Dry biomass before and 28 DAT (mean ± SE); initial P = 0.75. Letters compare 28 DAT.

Why biomass fell: killed, flattened rye lies on moist soil where microbes decompose it, while surviving tillers keep adding biomass **(Peterson et al. 2025)**.

Future Research

- Test another growth stage, e.g. anthesis
- Compare different effective EWC doses
- Measure yield of the following cash crop
- Track weed suppression through the season
- Repeat across years, sites and cover crops

References

Creamer NG, Dabney SM (2002) Am J Altern Agric 17:32–40
Frasconi C et al. (2019) PLoS ONE 14:e0211573
Kornecki TS et al. (2012) HortScience 47:217–223
Kornecki TS (2015) Appl Eng Agric 31:849–856
Mirsky SB et al. (2009) Agron J 101:1589–1596
Peterson CM, Tully KL, Mirsky SB (2025) Agron J, doi:10.1002/agj2.70201
Seabold S, Perktold J (2010) Proc 9th Python Sci Conf, 92–96
Slaven MJ et al. (2023) Weed Sci 71:403–421
Sullivan A (2025) GROW, growiwm.org
WSU (2025) Weeders of the West, smallgrains.wsu.edu

Acknowledgements

Funding: USDA-NIFA; TAMU Weed Science Lab