

# Numerical Analysis: Homework #2

Due on January 19, 2015

*Professor Mohler MWF 2:15*

**Rick Sullivan**

## Problem 1

Let  $f(x) = -x^3 - \cos x$  and  $p_0 = 1$ . Use two iterations of Newton's method to approximate the root of  $f(x)$ .

### Solution

Iteration one:

$$f'(x) = -3x^2 + \sin x$$

Use the slope at  $x = p_0 = 1$  to approximate the root for our next iteration.

$$\begin{aligned} f(1) &\approx -1.5403 \\ f'(1) &\approx -2.15853 \\ p_1 &= x_0 - \frac{f(x_0)}{f'(x_0)} \\ &= 1 - \frac{-1.5403}{-2.15853} \\ &= 0.2864 \end{aligned}$$

Iteration two with  $p_1 = 0.2864$ :

$$\begin{aligned} f(0.2864) &\approx -0.982759 \\ f'(0.2864) &\approx 0.0364182 \\ p_2 &= x_1 - \frac{f(x_1)}{f'(x_1)} \\ &= 0.2864 - \frac{-0.982759}{0.0364182} \\ &= 27.98538 \end{aligned}$$

Our final approximation is 27.9854. This is very inaccurate because the slope at our initial choice is nearly 0, pushing our estimation far from the actual root.

## Problem 2

The derivative of a function can be approximated by

$$f'(x) \approx \frac{-3f(x) + 4f(x+h) - f(x+2h)}{2h}$$

Show that the error of this approximation is  $O(h^2)$ .

### Solution

The Taylor series for  $f(x+h)$  and  $f(x+2h)$  are as follows.

$$\begin{aligned} f(x+h) &= f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(x) + Ch^4 \\ f(x+2h) &= f(x) + 2hf'(x) + 2h^2f''(x) + (4/3)h^3f'''(x) + Ch^4 \end{aligned}$$

Using those expansions to calculate the error of the approximation

$$\begin{aligned}
err_h &= f'(x) - \frac{-3f(x) + 4f(x+h) - f(x+2h)}{2h} \\
err_h &= f'(x) - \frac{-3f(x) + 4 \left[ f(x) + hf'(x) + \frac{h^2}{2}f''(x) + \frac{h^3}{6}f'''(x) + Ch^4 \right]}{2h} \\
&\quad - \frac{[f(x) + 2hf'(x) + 2h^2f''(x) + (4/3)h^3f'''(x) + Ch^4]}{2h} \\
&= f'(x) - \frac{4hf'(x) + 2h^2f''(x) + \frac{2h^3}{3}f'''(x) - 2hf'(x) - 2h^2f''(x) - \frac{4h^3}{3}f'''(x) + Ch^4}{2h} \\
&= f'(x) - \frac{2hf'(x) - \frac{2h^3}{3}f'''(x) + Ch^4}{2h} \\
&= f'(x) - f'(x) - \frac{h^2}{3}f'''(x) + Ch^3 \\
&= \frac{h^2}{3}f'''(x) + Ch^3
\end{aligned}$$

If  $h$  is small, the first term will be much more significant than any other term in the expansion. Therefore, the error of this approximation is  $O(h^2)$ .

### Problem 3

Use the approximation for  $f''(x)$  given in class to approximate the second derivative of

$$f(x) = \cos e^{x^2} + x^3$$

at  $x = 1$  using  $h = .1, .05, .025$ . How does the decrease in error compare to the order of accuracy of the method (is the decrease what we should expect)?

#### Solution

The equation for  $f''(x)$  given in class:

$$f''(x) = \frac{f(x+h) - 2f(x) + f(x-h)}{h^2}$$

Using the specified inputs:

$$\begin{aligned}
f''(1)_{0.1} &= \frac{f(1+0.1) - 2f(1) + f(1-0.1)}{0.1^2} \\
&\approx 0.3534 \\
f''(1)_{0.05} &= \frac{f(1+0.05) - 2f(1) + f(1-0.05)}{0.05^2} \\
&\approx 0.1661 \\
f''(1)_{0.025} &= \frac{f(1+0.025) - 2f(1) + f(1-0.025)}{0.025^2} \\
&\approx 0.1164
\end{aligned}$$

In class, we determined that the calculated error of this approach is  $O(h^2)$ . Therefore, estimation with choice  $h_1$  should have 4 times the error of an estimation with choice  $h_2 = (1/2)h_1$ .

$$\begin{aligned}
\frac{f''(1)_{0.1}}{f''(1)_{0.05}} &= 2.1276 \\
\frac{f''(1)_{0.05}}{f''(1)_{0.025}} &= 1.4270
\end{aligned}$$

ADD SOMETHING HERE. ARE THESE RESULTS CORRECT?

## Problem 4

Use the composite trapezoidal rule and Simpson's rule with  $n = 4$  to approximate

$$\int_0^1 x^2 e^{-x}$$

What is the error of each method?

### Solution

With  $n = 4$ , we will divide the interval  $[0, 1]$  into 4 segments with  $h = 0.25$ . Using the composite trapezoidal rule:

$$\begin{aligned} \int_0^1 x^2 e^{-x} &= \sum_{i=0}^3 \frac{1}{4} \frac{f(x_i) + f(x_{i+1})}{2} \\ &= \frac{1}{8} [f(0) + 2f(1/4) + 2f(1/2) + 2f(3/4) + f(1)] \\ &= \frac{1}{8} \left[ 1 + \frac{1}{8e^{1/4}} + \frac{1}{2e^{1/2}} + \frac{9}{8e^{3/4}} + \frac{1}{e} \right] \\ &\approx 0.2848 \end{aligned}$$

Using the composite Simpson's rule:

$$\begin{aligned} \int_0^1 x^2 e^{-x} &= \sum_{i=0}^1 \frac{0.25}{3} [f(x_{2i}) + 4f(x_{2i+1}) + f(x_{2i+2})] \\ &= \frac{1}{12} \left[ \left( 1 + \frac{1}{4e^{1/4}} + \frac{1}{4e^{1/2}} \right) + \left( \frac{1}{4e^{1/2}} + \frac{9}{4e^{3/4}} + \frac{1}{e} \right) \right] \\ &\approx 0.2441 \end{aligned}$$

TODO: ENTER ERROR OF THIS METHOD

## Problem 5

Determine  $c_0$ ,  $c_1$ , and  $c_2$  such that

$$\int_0^2 p(x) dx = c_0 p(0) + c_1 p(1) + c_2 p(2)$$

for all polynomials  $p(x)$  of degree 2 or less.

### Solution

Simpson's rule provides exact estimation for polynomials up to degree 3.

$$\begin{aligned} \int_0^2 p(x) dx &\approx \frac{1}{3} [p(0) + 4p(1) + p(2)] \\ &= \frac{1}{3} p(0) + \frac{4}{3} p(1) + \frac{1}{3} p(2) \end{aligned}$$

Therefore,  $c_0 = 1/3$ ,  $c_1 = 4/3$ , and  $c_2 = 1/3$ .