

Coisotropic reduction in different phase spaces

XVIII International Young Researchers Workshop in
Geometry, Dynamics and Field Theory

Rubén Izquierdo, Manuel De León

Wednesday 21st February, 2024

UCM-ICMAT

Coisotropic reduction in non-dissipative mechanics

Coisotropic reduction in dissipative mechanics

Coisotropic reduction in non-dissipative mechanics

The canonical phase spaces

- If Q is the configuration space of a mechanical system, the phase space $M := T^*Q$ inherits a canonical **symplectic structure** (M, ω) ,

$$\omega := \omega_Q = -d\lambda_Q = dq^i \wedge dp_i.$$

- The phase space $M := T^*Q \times \mathbb{R}$ inherits a canonical cosymplectic structure, (M, ω, θ) ,

$$\omega = \omega_Q = dq^i \wedge dp_i, \quad \theta = dt.$$

The canonical phase spaces

- If Q is the configuration space of a mechanical system, the phase space $M := T^*Q$ inherits a canonical symplectic structure (M, ω) ,

$$\omega := \omega_Q = -d\lambda_Q = dq^i \wedge dp_i.$$

- The phase space $M := T^*Q \times \mathbb{R}$ inherits a canonical **cosymplectic structure**, (M, ω, θ) ,

$$\omega = \omega_Q = dq^i \wedge dp_i, \quad \theta = dt.$$

The Poisson bracket

Symplectic and cosymplectic manifolds are **Poisson manifolds** with the bracket

$$\{f, g\} = \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i}.$$

In each of these cases, the bracket is induced by the Poisson bivector

$$\Lambda = \frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}, \quad \{f, g\} = \Lambda(df, dg).$$

We have an induced map

$$\sharp_\Lambda : T^*M \rightarrow TM, \quad \alpha \mapsto \iota_\alpha \Lambda.$$

Denote

$$\mathcal{H} := \text{im } \sharp_\Lambda = \left\langle \frac{\partial}{\partial q^i}, \frac{\partial}{\partial p_i} \right\rangle.$$

In symplectic manifolds, $\mathcal{H} = TM$.

The Poisson bracket

Symplectic and cosymplectic manifolds are Poisson manifolds with the bracket

$$\{f, g\} = \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i}.$$

In each of these cases, the bracket is induced by the **Poisson bivector**

$$\Lambda = \frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}, \quad \{f, g\} = \Lambda(df, dg).$$

We have an induced map

$$\sharp_\Lambda : T^*M \rightarrow TM, \quad \alpha \mapsto \iota_\alpha \Lambda.$$

Denote

$$\mathcal{H} := \text{im } \sharp_\Lambda = \left\langle \frac{\partial}{\partial q^i}, \frac{\partial}{\partial p_i} \right\rangle.$$

In symplectic manifolds, $\mathcal{H} = TM$.

The Poisson bracket

Symplectic and cosymplectic manifolds are Poisson manifolds with the bracket

$$\{f, g\} = \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i}.$$

In each of these cases, the bracket is induced by the Poisson bivector

$$\Lambda = \frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}, \quad \{f, g\} = \Lambda(df, dg).$$

We have an induced map

$$\sharp_\Lambda : T^*M \rightarrow TM, \quad \alpha \mapsto \iota_\alpha \Lambda.$$

Denote

$$\mathcal{H} := \text{im } \sharp_\Lambda = \left\langle \frac{\partial}{\partial q^i}, \frac{\partial}{\partial p_i} \right\rangle.$$

In symplectic manifolds, $\mathcal{H} = TM$.

The Poisson bracket

Symplectic and cosymplectic manifolds are Poisson manifolds with the bracket

$$\{f, g\} = \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} - \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i}.$$

In each of these cases, the bracket is induced by the Poisson bivector

$$\Lambda = \frac{\partial}{\partial q^i} \wedge \frac{\partial}{\partial p_i}, \quad \{f, g\} = \Lambda(df, dg).$$

We have an induced map

$$\sharp_\Lambda : T^*M \rightarrow TM, \quad \alpha \mapsto \iota_\alpha \Lambda.$$

Denote

$$\mathcal{H} := \text{im } \sharp_\Lambda = \left\langle \frac{\partial}{\partial q^i}, \frac{\partial}{\partial p_i} \right\rangle.$$

In symplectic manifolds, $\mathcal{H} = TM$.

Coisotropic and Lagrangian submanifolds

If $\Delta \subseteq T_x M$, we define the **orthogonal** as

$$\Delta^{\perp_\Delta} := \sharp_\Delta(\Delta^0),$$

where $\Delta^0 \subseteq T_x^* M$ is the annihilator of Δ . We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Delta} \subseteq \Delta,$$

- Lagrangian, if

$$\Delta^{\perp_\Delta} = \Delta \cap \mathcal{H}.$$

These definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Lagrangian submanifolds

If $\Delta \subseteq T_x M$, we define the orthogonal as

$$\Delta^{\perp_\Delta} := \sharp_\Delta(\Delta^0),$$

where $\Delta^0 \subseteq T_x^* M$ is the annihilator of Δ . We say that Δ is

- **Coisotropic**, if

$$\Delta^{\perp_\Delta} \subseteq \Delta,$$

- Lagrangian, if

$$\Delta^{\perp_\Delta} = \Delta \cap \mathcal{H}.$$

These definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Lagrangian submanifolds

If $\Delta \subseteq T_x M$, we define the orthogonal as

$$\Delta^{\perp_\Lambda} := \sharp_\Lambda(\Delta^0),$$

where $\Delta^0 \subseteq T_x^* M$ is the annihilator of Δ . We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Lagrangian, if

$$\Delta^{\perp_\Lambda} = \Delta \cap \mathcal{H}.$$

These definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Lagrangian submanifolds

If $\Delta \subseteq T_x M$, we define the orthogonal as

$$\Delta^{\perp_\Lambda} := \sharp_\Lambda(\Delta^0),$$

where $\Delta^0 \subseteq T_x^* M$ is the annihilator of Δ . We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Lagrangian, if

$$\Delta^{\perp_\Lambda} = \Delta \cap \mathcal{H}.$$

These definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic reduction in symplectic geometry

Let (M, ω) be a **symplectic manifold** and $i : N \hookrightarrow M$ be a **coisotropic submanifold**.

Proposition

$(TN)^{\perp_{\omega}} \subseteq TN$ is an involutive distribution.

Define \mathcal{F} to be the maximal foliation associated to $(TN)^{\perp_{\omega}}$. We will assume that N/\mathcal{F} admits a manifold structure such that the canonical projection $\pi : N \rightarrow N/\mathcal{F}$ is a submersion.

Theorem ([?]Weinstein)

There exists a unique symplectic form ω_N defined on N/\mathcal{F} such that

$$\pi^* \omega_N = i^* \omega.$$

Coisotropic reduction in symplectic geometry

Let (M, ω) be a symplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\omega}} \subseteq TN$ is an involutive distribution.

Define \mathcal{F} to be the maximal foliation associated to $(TN)^{\perp_{\omega}}$. We will assume that N/\mathcal{F} admits a manifold structure such that the canonical projection $\pi : N \rightarrow N/\mathcal{F}$ is a submersion.

Theorem ([?]Weinstein)

There exists a unique symplectic form ω_N defined on N/\mathcal{F} such that

$$\pi^* \omega_N = i^* \omega.$$

Coisotropic reduction in symplectic geometry

Let (M, ω) be a symplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\Lambda}} \subseteq TN$ is an involutive distribution.

Define \mathcal{F} to be the maximal foliation associated to $(TN)^{\perp_{\Lambda}}$. We will assume that N/\mathcal{F} admits a manifold structure such that the canonical projection $\pi : N \rightarrow N/\mathcal{F}$ is a submersion.

Theorem ([?]Weinstein)

There exists an unique symplectic form ω_N defined on N/\mathcal{F} such that

$$\pi^* \omega_N = i^* \omega.$$

Coisotropic reduction in symplectic geometry

Let (M, ω) be a symplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\Lambda}} \subseteq TN$ is an involutive distribution.

Define \mathcal{F} to be the maximal foliation associated to $(TN)^{\perp_{\Lambda}}$. We will assume that N/\mathcal{F} admits a manifold structure such that the canonical projection $\pi : N \rightarrow N/\mathcal{F}$ is a summersion.

Theorem ([?]Weinstein)

There exists an unique *symplectic* form ω_N defined on N/\mathcal{F} such that

$$\pi^* \omega_N = i^* \omega.$$

Coisotropic reduction in cosymplectic geometry

Let (M, ω, θ) be a **cosymplectic manifold** and $i : N \hookrightarrow M$ be a **coisotropic submanifold**.

Proposition

$(TN)^{\perp_{\theta}} \subseteq TN$ is an involutive distribution.

Suppose N/\mathcal{F} admits a manifold structure such that $\pi : N \rightarrow N/\mathcal{F}$ defines a submersion.

Theorem (?? RIL-MLR)

- If $TN \subseteq \mathcal{H}$, N/\mathcal{F} admits an unique symplectic structure compatible with the structure defined on M .
- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} admits an unique cosymplectic structure compatible with the one defined on M .

Coisotropic reduction in cosymplectic geometry

Let (M, ω, θ) be a cosymplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\theta}} \subseteq TN$ is an involutive distribution.

Suppose N/\mathcal{F} admits a manifold structure such that $\pi : N \rightarrow N/\mathcal{F}$ defines a submersion.

Theorem (?? RIL-MLR)

- If $TN \subseteq \mathcal{H}$, N/\mathcal{F} admits a unique symplectic structure compatible with the structure defined on M .
- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} admits a unique cosymplectic structure compatible with the one defined on M .

Coisotropic reduction in cosymplectic geometry

Let (M, ω, θ) be a cosymplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\theta}} \subseteq TN$ is an involutive distribution.

Suppose N/\mathcal{F} admits a manifold structure such that $\pi : N \rightarrow N/\mathcal{F}$ defines a summersion.

Theorem (?? RIL-MLR)

- If $TN \subseteq \mathcal{H}$, N/\mathcal{F} admits an unique symplectic structure compatible with the strcuture defined on M .
- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} admits an unique cosymplectic sturcture compatible with the one defined on M .

Coisotropic reduction in cosymplectic geometry

Let (M, ω, θ) be a cosymplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\Lambda}} \subseteq TN$ is an involutive distribution.

Suppose N/\mathcal{F} admits a manifold structure such that $\pi : N \rightarrow N/\mathcal{F}$ defines a submersion.

Theorem (?? RIL-MLR)

- If $TN \subseteq \mathcal{H}$, N/\mathcal{F} admits a unique **symplectic structure** compatible with the structure defined on M .
- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} admits a unique **cosymplectic structure** compatible with the one defined on M .

Coisotropic reduction in cosymplectic geometry

Let (M, ω, θ) be a cosymplectic manifold and $i : N \hookrightarrow M$ be a coisotropic submanifold.

Proposition

$(TN)^{\perp_{\Lambda}} \subseteq TN$ is an involutive distribution.

Suppose N/\mathcal{F} admits a manifold structure such that $\pi : N \rightarrow N/\mathcal{F}$ defines a summersion.

Theorem (?? RIL-MLR)

- If $TN \subseteq \mathcal{H}$, N/\mathcal{F} admits an unique symplectic structure compatible with the strcuture defined on M .
- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} admits an unique **cosymplectic sturcture** compatible with the one defined on M .

Coisotropic reduction in dissipative mechanics

The canonical phase spaces

- The phase space of an autonomous dissipative system is $T^*Q \times \mathbb{R}$, with its canonical **contact structure**

$$\eta = dz - p_i dq^i.$$

- If we want to study time-dependent dissipative mechanics, the phase space is $T^*Q \times \mathbb{R} \times \mathbb{R}$ endowed with its cocontact structure

$$\eta = dz - p_i dq^i; \quad \theta = dt.$$

The canonical phase spaces

- The phase space of an autonomous dissipative system is $T^*Q \times \mathbb{R}$, with its canonical contact structure

$$\eta = dz - p_i dq^i.$$

- If we want to study time-dependent dissipative mechanics, the phase space is $T^*Q \times \mathbb{R} \times \mathbb{R}$ endowed with its **cocontact structure**

$$\eta = dz - p_i dq^i; \quad \theta = dt.$$

The Jacobi bracket

In both of these phase spaces there is a **Jacobi bracket** which is locally given by

$$\{f, g\} = \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i} - \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} + p_i \left(\frac{\partial f}{\partial p_i} \frac{\partial g}{\partial z} - \frac{\partial g}{\partial p_i} \frac{\partial f}{\partial z} \right) + g \frac{\partial f}{\partial z} - f \frac{\partial g}{\partial z}.$$

This Jacobi bracket is defined through the Jacobi bivector and a vector field

$$\Lambda = \frac{\partial}{\partial p_i} \wedge \frac{\partial}{\partial q^i} + p_i \frac{\partial}{\partial p_i} \wedge \frac{\partial}{\partial z},$$

$$E = -\frac{\partial}{\partial z},$$

as

$$\{f, g\} = \Lambda(df, dg) + fE(g) - gE(f).$$

The Jacobi bracket

In both of these phase spaces there is a Jacobi bracket which is locally given by

$$\{f, g\} = \frac{\partial f}{\partial p_i} \frac{\partial g}{\partial q^i} - \frac{\partial f}{\partial q^i} \frac{\partial g}{\partial p_i} + p_i \left(\frac{\partial f}{\partial p_i} \frac{\partial g}{\partial z} - \frac{\partial g}{\partial p_i} \frac{\partial f}{\partial z} \right) + g \frac{\partial f}{\partial z} - f \frac{\partial g}{\partial z}.$$

This Jacobi bracket is defined through the **Jacobi bivector** and a vector field

$$\Lambda = \frac{\partial}{\partial p_i} \wedge \frac{\partial}{\partial q^i} + p_i \frac{\partial}{\partial p_i} \wedge \frac{\partial}{\partial z},$$

$$E = -\frac{\partial}{\partial z},$$

as

$$\{f, g\} = \Lambda(df, dg) + fE(g) - gE(f).$$

Coisotropic and Legendrian submanifolds

The orthogonal of a distribution $\Delta \subseteq TM$ is defined as

$$\Delta^{\perp_\Lambda} = \sharp_\Lambda(\Delta^0),$$

where

$$\sharp_\Lambda : T^*M \rightarrow TM; \quad \alpha \mapsto \iota_\alpha \Lambda.$$

We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Legendrian, if

$$\Delta^{\perp_\Lambda} = \Delta.$$

The same definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Legendrian submanifolds

The orthogonal of a distribution $\Delta \subseteq TM$ is defined as

$$\Delta^{\perp_\Lambda} = \sharp_\Lambda(\Delta^0),$$

where

$$\sharp_\Lambda : T^*M \rightarrow TM; \quad \alpha \mapsto \iota_\alpha \Lambda.$$

We say that Δ is

- **Coisotropic**, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Legendrian, if

$$\Delta^{\perp_\Lambda} = \Delta.$$

The same definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Legendrian submanifolds

The orthogonal of a distribution $\Delta \subseteq TM$ is defined as

$$\Delta^{\perp_\Lambda} = \sharp_\Lambda(\Delta^0),$$

where

$$\sharp_\Lambda : T^*M \rightarrow TM; \quad \alpha \mapsto \iota_\alpha \Lambda.$$

We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Legendrian, if

$$\Delta^{\perp_\Lambda} = \Delta.$$

The same definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic and Legendrian submanifolds

The orthogonal of a distribution $\Delta \subseteq TM$ is defined as

$$\Delta^{\perp_\Lambda} = \sharp_\Lambda(\Delta^0),$$

where

$$\sharp_\Lambda : T^*M \rightarrow TM; \quad \alpha \mapsto \iota_\alpha \Lambda.$$

We say that Δ is

- Coisotropic, if

$$\Delta^{\perp_\Lambda} \subseteq \Delta,$$

- Legendrian, if

$$\Delta^{\perp_\Lambda} = \Delta.$$

The same definitions apply to submanifolds $N \hookrightarrow M$ as well.

Coisotropic reduction

Proposition

If $N \hookrightarrow M$ is a **coisotropic** submanifold, then $(TN)^{\perp_{\Lambda}}$ is involutive and thus arises from a maximal foliation \mathcal{F} .

We assume that $\frac{\partial}{\partial z} \in TN$.

Theorem

If M is a contact manifold ([2]), N/\mathcal{F} admits a unique contact structure compatible with the one on M .

If M is a cocontact manifold ([1]):

- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} inherits a unique cocontact structure from M .*
- If $TN \subseteq \text{im } \sharp_{\Lambda} \oplus \langle \frac{\partial}{\partial z} \rangle$, N/\mathcal{F} inherits a unique contact structure from M .*

Coisotropic reduction

Proposition

If $N \hookrightarrow M$ is a coisotropic submanifold, then $(TN)^{\perp_{\Lambda}}$ is involutive and thus arises from a maximal foliation \mathcal{F} .

We assume that $\frac{\partial}{\partial z} \in TN$.

Theorem

If M is a **contact** manifold ([2]), N/\mathcal{F} admits a unique **contact** structure compatible with the one on M .

If M is a cocontact manifold ([1]):

- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} inherits a unique cocontact structure from M .
- If $TN \subseteq \text{im } \sharp_{\Lambda} \oplus \langle \frac{\partial}{\partial z} \rangle$, N/\mathcal{F} inherits a unique contact structure from M .

Coisotropic reduction

Proposition

If $N \hookrightarrow M$ is a coisotropic submanifold, then $(TN)^{\perp_{\Lambda}}$ is involutive and thus arises from a maximal foliation \mathcal{F} .

We assume that $\frac{\partial}{\partial z} \in TN$.

Theorem

If M is a contact manifold ([2]), N/\mathcal{F} admits a unique contact structure compatible with the one on M .

If M is a **cocontact** manifold ([1]):

- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} inherits a unique cocontact structure from M .
- If $TN \subseteq \text{im } \sharp_{\Lambda} \oplus \langle \frac{\partial}{\partial z} \rangle$, N/\mathcal{F} inherits a unique contact structure from M .

Coisotropic reduction

Proposition

If $N \hookrightarrow M$ is a coisotropic submanifold, then $(TN)^{\perp_{\Lambda}}$ is involutive and thus arises from a maximal foliation \mathcal{F} .

We assume that $\frac{\partial}{\partial z} \in TN$.

Theorem

If M is a contact manifold ([2]), N/\mathcal{F} admits a unique contact structure compatible with the one on M .

If M is a cocontact manifold ([1]):

- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} inherits a unique **cocontact** structure from M .*
- If $TN \subseteq \text{im } \sharp_{\Lambda} \oplus \langle \frac{\partial}{\partial z} \rangle$, N/\mathcal{F} inherits a unique contact structure from M .*

Coisotropic reduction

Proposition

If $N \hookrightarrow M$ is a coisotropic submanifold, then $(TN)^{\perp_{\Lambda}}$ is involutive and thus arises from a maximal foliation \mathcal{F} .

We assume that $\frac{\partial}{\partial z} \in TN$.

Theorem

If M is a contact manifold ([2]), N/\mathcal{F} admits a unique contact structure compatible with the one on M .

If M is a cocontact manifold ([1]):

- If $\frac{\partial}{\partial t} \in TN$, N/\mathcal{F} inherits a unique cocontact structure from M .*
- If $TN \subseteq \text{im } \sharp_{\Lambda} \oplus \langle \frac{\partial}{\partial z} \rangle$, N/\mathcal{F} inherits a unique **contact** structure from M .*

References



R. Abraham and J.E. Marsden.

Foundations of mechanics.

Number 364. The Benjamin/Cummings Publishing Company, Reading, Massachusetts, 19788.



Manuel de León and Rubén Izquierdo-López.

A review on coisotropic reduction in symplectic, cosymplectic, contact and co-contact hamiltonian systems, 2023.



Manuel de León and Manuel Lainz Valcázar.

Contact hamiltonian systems.

Journal of Mathematical Physics, 60(10), October 2019.