



Universität Hamburg

DER FORSCHUNG | DER LEHRE | DER BILDUNG

Bachelorthesis

Title

vorgelegt von

TJARK SIEVERS

Fakultät: Mathematik, Informatik und Naturwissenschaften

Fachbereich: Physik

Studiengang: Physik

Matrikelnummer: 7147558

Erstgutachter:

Zweitgutachter:

Kurzzusammenfassung

Danksagung

Motivation

Inhaltsverzeichnis

| | |
|--|----------|
| Kurzzusammenfassung | i |
| Danksagung | iii |
| Motivation | v |
| I Physical and Computational Basics | 1 |
| I.1 The Schrödinger Equation | 1 |
| I.2 Density Functional Theory | 1 |
| I.2.1 Hohenberg-Kohn theorems | 1 |
| I.2.2 Kohn-Sham equations | 1 |

Conventions

Throughout the text of this thesis scalars are written in italic s , vectors in bold italic \boldsymbol{v} and matrices in bold \mathbf{M} fonts. The summation/multiplication over nearest neighbour sites i and j is $\langle ij \rangle$ as a subscript to \sum/\prod . Furthermore, Hartree atomic units are used in general and only in selected instances it is deviated from this: $\hbar = m_e = e = 4\pi/\epsilon_0 = 1$.

I Physical and Computational Basics

I.1 The Schrödinger Equation

$$\hat{H}\Psi(\mathbf{r}_1, \dots, \mathbf{r}_N) = E\Psi(\mathbf{r}_1, \dots, \mathbf{r}_N) \quad (\text{I.1})$$

The Hamiltonian for an isolated system of N interacting electrons in an external potential, so the

I.2 Density Functional Theory

A direct solution to the electronic structure problem, this meaning obtaining the ground-state many-body wavefunction $\Psi(\mathbf{r}_1, \dots, \mathbf{r}_N)$ for a given potential is analytically impossible even for a small number of electrons. As such, the need for good approximations to obtain results for real world systems is high. One particularly successful approach is [Density Functional Theory](#) (DFT) [Density Functional Theory](#)

I.2.1 Hohenberg-Kohn theorems

I.2.2 Kohn-Sham equations

The success of DFT

