

I EG-X Model - Trying BCS theory

I.1 BCS Theory on the EG-X Model

I.1.1 Self-consistent calculation of the superconducting gaps

This does not really work! I neglect interband pairing at some point, so i throw away 6 out of 9 gap equations I have, also the GF ansatz works with diagonal Matsubara GFs, so I dont think it works here.

Compare [Bruus_Flensberg_2004]. Notable here: Multiple bands, and the gaps in each band depend in a complicated manner on the parameters U_α and the orbital Green's functions.

Define normal Green's function:

$$\mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) = -\langle T_\tau d_{n\mathbf{k}\uparrow}(\tau) d_{n\mathbf{k}\uparrow}^\dagger(0) \rangle \quad (\text{I.1})$$

Anomalous Green's function:

$$\mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) = -\langle T_\tau d_{n-\mathbf{k}\downarrow}(\tau) d_{n\mathbf{k}\uparrow}^\dagger(0) \rangle \quad (\text{I.2})$$

Equations of motion (Heisenberg equation), follow [Bruus_Flensberg_2004]:

$$\partial_\tau \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) = -\delta(\tau) + \langle T_\tau [d_{n\mathbf{k}\uparrow}, H_{BdG}] (\tau) d_{n\mathbf{k}\uparrow}^\dagger(0) \rangle \quad (\text{I.3})$$

$$\partial_\tau \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) = \langle T_\tau [d_{n-\mathbf{k}\downarrow}, H_{BdG}] (\tau) d_{n\mathbf{k}\uparrow}^\dagger(0) \rangle \quad (\text{I.4})$$

To calculate the commutators, use the relation (for operators A, B, C):

$$[A, BC] = ABC - BCA = (\{A, B\} - BA)C - B(\{C, A\} - AC) \quad (\text{I.5})$$

$$[d_{n-\mathbf{k}\downarrow}^\dagger, H_0] = \sum_{n'\mathbf{k}'\sigma'} \xi_{n'\mathbf{k}'} [d_{n-\mathbf{k}\downarrow}^\dagger, d_{n'\mathbf{k}'\sigma'}^\dagger d_{n'\mathbf{k}'\sigma'}] \quad (\text{I.6})$$

$$= \sum_{n'\mathbf{k}'\sigma'} \xi_{n'\mathbf{k}'} \left(\{d_{n-\mathbf{k}\downarrow}^\dagger, d_{n'\mathbf{k}'\sigma'}^\dagger\} - d_{n'\mathbf{k}'\sigma'}^\dagger d_{n-\mathbf{k}\downarrow}^\dagger \right) d_{n'\mathbf{k}'\sigma'} \quad (\text{I.7})$$

$$- d_{n'\mathbf{k}'\sigma'}^\dagger \left(\{d_{n'\mathbf{k}'\sigma'}, d_{n-\mathbf{k}\downarrow}^\dagger\} - d_{n-\mathbf{k}\downarrow}^\dagger d_{n'\mathbf{k}'\sigma'} \right) \quad (\text{I.8})$$

$$= \sum_{n'\mathbf{k}'\sigma'} \xi_{n'\mathbf{k}'} \left(-d_{n'\mathbf{k}'\sigma'}^\dagger d_{n-\mathbf{k}\downarrow}^\dagger d_{n'\mathbf{k}'\sigma'} - d_{n'\mathbf{k}'\sigma'}^\dagger \delta_{n'\mathbf{k}'\sigma', n-\mathbf{k}\uparrow} + d_{n'\mathbf{k}'\sigma'}^\dagger d_{n-\mathbf{k}\downarrow}^\dagger d_{n'\mathbf{k}'\sigma'} \right) \quad (\text{I.9})$$

$$= -\xi_{n\mathbf{k}} d_{n\mathbf{k}\uparrow}^\dagger \quad (\text{I.10})$$

$$\left[d_{n-\mathbf{k}\downarrow}, -\sum_{m\mathbf{k}'} \Delta_m^* d_{m-\mathbf{k}'\downarrow} d_{m\mathbf{k}'\uparrow} \right] \quad (\text{I.11})$$

$$= -\sum_{m\mathbf{k}'} \Delta_m^* \left(\{d_{n-\mathbf{k}\downarrow}, d_{m-\mathbf{k}'\downarrow}\} - d_{m-\mathbf{k}'\downarrow} d_{n-\mathbf{k}\downarrow} \right) d_{m\mathbf{k}'\uparrow} \quad (\text{I.12})$$

$$- d_{m-\mathbf{k}'\downarrow} \left(\{d_{m\mathbf{k}'\uparrow}, d_{n-\mathbf{k}\downarrow}\} - d_{n-\mathbf{k}\downarrow} d_{m\mathbf{k}'\uparrow} \right) \quad (\text{I.13})$$

$$= -\sum_{m\mathbf{k}'} \Delta_m^* \left(\delta_{n-\mathbf{k}\downarrow, m-\mathbf{k}'\downarrow} - d_{m-\mathbf{k}'\downarrow} d_{n-\mathbf{k}\downarrow} \right) d_{m\mathbf{k}'\uparrow} + d_{m-\mathbf{k}'\downarrow} d_{n-\mathbf{k}\downarrow} d_{m\mathbf{k}'\uparrow} \quad (\text{I.14})$$

$$= -\Delta_n^* d_{n\mathbf{k}\uparrow} \quad (\text{I.15})$$

$$\partial_\tau \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) = -\xi_{n\mathbf{k}} \langle T_\tau (d_{n-\mathbf{k}\downarrow}^\dagger(\tau) d_{n\mathbf{k}\uparrow}^\dagger(0)) \rangle - \Delta_n^* \langle T_\tau (d_{n\mathbf{k}\uparrow}(\tau) d_{n\mathbf{k}\uparrow}^\dagger(0)) \rangle \quad (\text{I.16})$$

$$= \xi_{n\mathbf{k}} \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) + \Delta_n^* \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) \quad (\text{I.17})$$

Similarly:

$$[d_{n-\mathbf{k}\uparrow}, H_0] = \sum_{n'\mathbf{k}'\sigma'} \xi_{n'\mathbf{k}'} [d_{n-\mathbf{k}\uparrow}, d_{n'\mathbf{k}'\sigma'}^\dagger d_{n'\mathbf{k}'\sigma'}] \quad (\text{I.18})$$

$$= \xi_n d_{n\mathbf{k}\uparrow}^\dagger \quad (\text{I.19})$$

$$\left[d_{n-\mathbf{k}\uparrow}, -\sum_{m\mathbf{k}'} \Delta_m d_{m-\mathbf{k}'\uparrow}^\dagger d_{m-\mathbf{k}'\downarrow}^\dagger \right] \quad (\text{I.20})$$

$$= -\Delta_n d_{n-\mathbf{k}\downarrow}^\dagger \quad (\text{I.21})$$

$$\partial_\tau \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) = -\delta(\tau) + \xi_{n\mathbf{k}} \langle T_\tau d_{n\mathbf{k}\uparrow}(\tau) d_{n\mathbf{k}\uparrow}^\dagger \rangle - \Delta_n \langle T_\tau d_{n-\mathbf{k}\downarrow}(\tau) d_{n\mathbf{k}\uparrow}^\dagger(0) \rangle \quad (\text{I.22})$$

$$= -\delta(\tau) - \xi_{n\mathbf{k}} \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) + \Delta_n \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) \quad (\text{I.23})$$

$$(\text{I.24})$$

All in all:

$$\partial_\tau \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) = -\delta(\tau) - \xi_{n\mathbf{k}} \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) + \Delta_n \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) \quad (\text{I.25})$$

$$\partial_\tau \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) = \xi_{n\mathbf{k}} \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, \tau) + \Delta_n^* \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, \tau) \quad (\text{I.26})$$

Fourier transform:

$$(-i\omega_n + \xi_{n\mathbf{k}}) \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, i\omega_n) = -1 + \Delta_n \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) \quad (\text{I.27})$$

$$(-i\omega_n - \xi_{n\mathbf{k}}) \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) = \Delta_n^* \mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, i\omega_n) \quad (\text{I.28})$$

This algebraic expression can be easily solved:

$$(-i\omega_n - \xi_{n\mathbf{k}}) \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) = \frac{\Delta_n^*}{-i\omega_n + \xi_{n\mathbf{k}}} (-1 + \Delta_n \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n)) \quad (\text{I.29})$$

$$(-i\omega_n - \xi_{n\mathbf{k}} - \frac{|\Delta_n|^2}{-i\omega_n + \xi_{n\mathbf{k}}}) \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) = \frac{-\Delta_n^*}{-i\omega_n + \xi_{n\mathbf{k}}} \quad (\text{I.30})$$

$$\left(\frac{(-i\omega_n - \xi_{n\mathbf{k}})(-i\omega_n + \xi_{n\mathbf{k}}) - |\Delta_n|^2}{-i\omega_n + \xi_{n\mathbf{k}}} \right) \mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) = \frac{-\Delta_n^*}{-i\omega_n + \xi_{n\mathbf{k}}} \quad (\text{I.31})$$

$$\mathcal{F}_{n\downarrow n\uparrow}(\mathbf{k}, i\omega_n) = \frac{-\Delta_n^*}{(-i\omega_n - \xi_{n\mathbf{k}})(-i\omega_n + \xi_{n\mathbf{k}}) - |\Delta_n|^2} \quad (\text{I.32})$$

$$= \frac{-\Delta_n^*}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.33})$$

$$= \frac{-\Delta_n^*}{(i\omega_n)^2 - E_{n\mathbf{k}}} \quad (\text{I.34})$$

$$(-i\omega_n + \xi_{n\mathbf{k}})\mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, i\omega_n) = -1 + \frac{-|\Delta_n|^2}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.35})$$

$$= \frac{-(i\omega_n)^2 + \xi_{n\mathbf{k}}^2 + |\Delta_n|^2 - |\Delta_n|^2}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.36})$$

$$= \frac{-(i\omega_n)^2 + \xi_{n\mathbf{k}}^2}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.37})$$

$$= \frac{(i\omega_n + \xi_{n\mathbf{k}})(-i\omega_n + \xi_{n\mathbf{k}})}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.38})$$

$$\mathcal{G}_{n\uparrow n\uparrow}(\mathbf{k}, i\omega_n) = \frac{i\omega + \xi_{n\mathbf{k}}}{(i\omega_n)^2 - \xi_{n\mathbf{k}}^2 - |\Delta_n|^2} \quad (\text{I.39})$$

$$= \frac{i\omega + \xi_{n\mathbf{k}}}{(i\omega_n)^2 - E_{n\mathbf{k}}} \quad (\text{I.40})$$

with the energies $E_{n\mathbf{k}} = \pm\sqrt{\xi_{n\mathbf{k}}^2 + |\Delta_n|^2}$.

To calculate the band gap in band n :

$$\Delta_n(\mathbf{k}) = -\sum_{\alpha} [G_{k\uparrow}]_{\alpha n}^* \Delta_{\alpha} [G_{-k\downarrow}]_{\alpha n}^* \quad (\text{I.41})$$

$$= \sum_{\alpha\mathbf{k}'} U_{\alpha} [G_{k\uparrow}]_{\alpha n}^* \langle c_{-k'\alpha\downarrow} c_{k'\alpha\uparrow} \rangle [G_{-k\downarrow}]_{\alpha n}^* \quad (\text{I.42})$$

$$= \sum_{\alpha\mathbf{k}'} U_{\alpha} [G_{k\uparrow}]_{\alpha n}^* [G_{-k\downarrow}]_{\alpha n}^* \sum_m [G_{-k'\downarrow}]_{\alpha m} [G_{k'\uparrow}]_{\alpha m} \langle d_{-k'm\downarrow} d_{k'm\uparrow} \rangle \quad (\text{I.43})$$

Can now use \mathcal{F} and fourier-transform:

$$\langle d_{-k'm\downarrow} d_{k'm\uparrow} \rangle = \mathcal{F}_{m\downarrow m\uparrow}^*(\mathbf{k}', \tau = 0^+) \quad (\text{I.44})$$

$$= \frac{1}{\beta} \sum_{i\omega_n} e^{-i\omega_n 0^+} \mathcal{F}_{m\downarrow m\uparrow}^*(\mathbf{k}', i\omega_n) \quad (\text{I.45})$$

The summation over the Matsubara frequencies can be solved via the Residue theorem (the poles z_0 of \mathcal{F} are the energies $\pm E_{m\mathbf{k}}$):

$$\frac{1}{\beta} \sum_{i\omega_n} e^{-i\omega_n 0^+} \mathcal{F}_{m\downarrow m\uparrow}^*(\mathbf{k}', i\omega_n) \quad (\text{I.46})$$

$$= \sum_{z_0 \text{ poles of } \mathcal{F}} e^{-z_0 0^+} n_F(z_0) \text{Res}_{z_0} \mathcal{F}_{m\downarrow m\uparrow}^*(\mathbf{k}', z_0) \quad (\text{I.47})$$

$$= e^{-E_{m\mathbf{k}} 0^+} n_F(E_{m\mathbf{k}}) \text{Res}_{E_{m\mathbf{k}}} \frac{-\Delta_m}{(i\omega_n)^2 - E_{m\mathbf{k}}} + e^{E_{m\mathbf{k}} 0^+} n_F(-E_{m\mathbf{k}}) \text{Res}_{-E_{m\mathbf{k}}} \frac{-\Delta_m}{(i\omega_n)^2 - E_{m\mathbf{k}}} \quad (\text{I.48})$$

with residue:

$$\text{Res}_{E_{m\mathbf{k}}} \frac{1}{(i\omega_n)^2 - z_0^2} = \frac{1}{\partial_z |_{z_0=E_{m\mathbf{k}}} ((i\omega)^2 - z_0^2)} = \frac{1}{2E_{m\mathbf{k}}} \quad (\text{I.49})$$

So we have

$$\langle d_{-k'm\downarrow} d_{k'm\uparrow} \rangle = -\Delta_m \left(\frac{n_F(E_{m\mathbf{k}})}{2E_{m\mathbf{k}}} - \frac{n_F(-E_{m\mathbf{k}})}{2E_{m\mathbf{k}}} \right) \quad (\text{I.50})$$

The n_F term can be written as:

$$n_F(E_{m\mathbf{k}'}) - n_F(-E_{m\mathbf{k}'}) = \frac{1}{e^{\beta E_{m\mathbf{k}'}} + 1} - \frac{1}{e^{-\beta E_{m\mathbf{k}'}} + 1} \quad (\text{I.51})$$

$$= \frac{e^{-\frac{1}{2}\beta E_{m\mathbf{k}'}}}{e^{-\frac{1}{2}\beta E_{m\mathbf{k}'}} + 1} \frac{1}{e^{\beta E_{m\mathbf{k}'}} + 1} - \frac{e^{\frac{1}{2}\beta E_{m\mathbf{k}'}}}{e^{\frac{1}{2}\beta E_{m\mathbf{k}'}} + 1} \frac{1}{e^{-\beta E_{m\mathbf{k}'}} + 1} \quad (\text{I.52})$$

$$= \frac{e^{-\frac{1}{2}\beta E_{m\mathbf{k}'}} - e^{\frac{1}{2}\beta E_{m\mathbf{k}'}}}{e^{\frac{1}{2}\beta E_{m\mathbf{k}'}} + e^{-\frac{1}{2}\beta E_{m\mathbf{k}'}}} \quad (\text{I.53})$$

$$= -\tanh\left(\frac{\beta E_{m\mathbf{k}'}}{2}\right) \quad (\text{I.54})$$

This results in the self-consistency equation for the gap:

$$\Delta_n(\mathbf{k}) = \sum_{\alpha m \mathbf{k}'} U_\alpha [G_{k\uparrow}]_{\alpha n}^* [G_{-k\downarrow}]_{\alpha n}^* [G_{-k'\downarrow}]_{\alpha m} [G_{k'\uparrow}]_{\alpha m} \Delta_m(\mathbf{k}') \frac{\tanh\left(\frac{\beta E_{m\mathbf{k}'}}{2}\right)}{2E_{m\mathbf{k}'}} \quad (\text{I.55})$$

Using time-reversal symmetry $[G_{-\mathbf{k}\downarrow}]_{\alpha m}^* = [G_{\mathbf{k}\uparrow}]_{\alpha m}$ this expression gets a bit simpler:

$$\Delta_n(\mathbf{k}) = \sum_{\alpha m \mathbf{k}'} U_\alpha |G_{\mathbf{k}\uparrow}]_{\alpha n}|^2 |G_{\mathbf{k}'\uparrow}]_{\alpha m}|^2 \Delta_m(\mathbf{k}') \frac{\tanh(\frac{\beta E_{m\mathbf{k}'}}{2})}{2E_{m\mathbf{k}'}} \quad (\text{I.56})$$

I.1.2 Computational Implementation

Use scipys `fixed_point` solver to solve the gap equation self-consistently.

Flatten $\Delta_n(\mathbf{k})$ the following way, to put it into the solver (\mathbf{k} discretized in some way):

$$x = \begin{pmatrix} \Re(\Delta_1(\mathbf{k}_1)) \\ \Re(\Delta_1(\mathbf{k}_2)) \\ \vdots \\ \Re(\Delta_2(\mathbf{k}_1)) \\ \vdots \\ \Re(\Delta_3(\mathbf{k}_1)) \\ \vdots \\ \Im(\Delta_1(\mathbf{k}_1)) \\ \vdots \\ \Im(\Delta_2(\mathbf{k}_1)) \\ \vdots \\ \Im(\Delta_3(\mathbf{k}_1)) \\ \vdots \end{pmatrix} \quad (\text{I.57})$$

so that accessing a certain element takes the form:

$$\Re \Delta_n(\mathbf{k}) = x \left[\text{index}(\mathbf{k}) + \frac{\text{len}(x) \cdot n}{6} \right] \quad (\text{I.58})$$

$$\Im \Delta_n(\mathbf{k}) = x \left[\text{index}(\mathbf{k}) + \frac{\text{len}(x) \cdot n}{6} + \frac{1}{2} \text{len}(x) \right] \quad (\text{I.59})$$