



Universität Hamburg  
DER FORSCHUNG | DER LEHRE | DER BILDUNG



UPPSALA  
UNIVERSITET

Master thesis

Title

Title Deutsch

vorgelegt von

TJARK SIEVERS

Fakultät: Mathematik, Informatik und Naturwissenschaften

Fachbereich: Physik

Studiengang: Physik

Matrikelnummer: 7147558

Erstgutachter: Prof. Dr. Annica Black-Schaffer (Uppsala University)

Zweitgutachterin: Prof. Dr. Tim Wehling



## **Kurzzusammenfassung**

## **Abstract**



# Contents

<b>Motivation</b>	<b>vii</b>
<b>I EG-X model</b>	<b>1</b>
I.1 Lattice Structure of Graphene . . . . .	1
I.2 Band structure . . . . .	3
<b>II Conclusion</b>	<b>5</b>
<b>A EG-X Hamiltonian in Reciprocal Space</b>	<b>7</b>
<b>Bibliography</b>	<b>11</b>
<b>List of Symbols</b>	<b>13</b>
<b>Acknowledgement</b>	<b>15</b>



# Motivation

Test

---

Test source

## Conventions

Throughout the text of this thesis scalars are written in italic  $s$ , vectors in bold italic  $\mathbf{s}$  and matrices in bold  $\mathbf{M}$  fonts. A matrix element is denoted as  $[\mathbf{M}]_{\alpha,\beta}$ . The summation/multiplication over nearest neighbour sites  $i$  and  $j$  is  $\langle ij \rangle$  as a subscript to  $\sum/\prod$ . Furthermore, Hartree atomic units are used in general and only in selected instances it is deviated from this:  $\hbar = m_e = e = 4\pi/\epsilon_0 = 1$ .



# I EG-X model

Test

---

Test source

The tight-binding Hamiltonian for the model reads

$$\begin{aligned} H_0 = & -t_X \sum_{\langle ij \rangle, \sigma} d_{i, \sigma}^\dagger d_{j, \sigma} - t_{Gr} \sum_{\langle ij \rangle, \sigma} \left( c_{i, \sigma}^{(A), \dagger} c_{j, \sigma}^{(B)} + c_{j, \sigma'}^{(B), \dagger} c_{i, \sigma}^{(A)} \right) \\ & + V \sum_{i, \sigma} d_{i, \sigma}^\dagger c_{i, \sigma}^{(A)} + \text{h.c.} \end{aligned} \quad (\text{I.1})$$

with:


- $d$  operators on the X atom
- $c^{(\epsilon)}$  operators on the Graphene site ( $\epsilon = A, B$ )
- $t_X$  next-nearest hopping for the X atoms
- $t_{Gr}$  next-nearest hopping on the Graphene
- $V$  hopping between X and Graphene B sites

This describes the

In material terms, this can be thought of as a sheet of graphene on top of another material, providing the additional X atoms, but in this thesis the model will be taken as a toy model, providing certain favorable aspects.

## I.1 Lattice Structure of Graphene

This section reviews the lattice structure of graphene and by extension also of the flat-band model, following the review [1].



Monolayer graphene forms a hexagonal lattice as seen in fig. I.1b. This is formed by two triangular sub lattices, so the unit cell of the hexagonal lattice has two atoms. The primitive vectors of the hexagonal lattice are

$$\begin{aligned} \mathbf{a}_1 &= \frac{a}{2} \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \\ \mathbf{a}_2 &= \frac{a}{2} \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix}, \end{aligned} \quad (\text{I.2})$$

with lattice constant  $a \approx 2.46 \text{ \AA}$  for graphene. The distance between nearest neighbor is

$$a = \sqrt{3}a_0. \quad (\text{I.3})$$

The primitive reciprocal lattice vectors  $\mathbf{b}_1, \mathbf{b}_2$  fulfill

$$\begin{aligned} \mathbf{a}_1 \cdot \mathbf{b}_1 &= \mathbf{a}_2 \cdot \mathbf{b}_2 = 2\pi \\ \mathbf{a}_1 \cdot \mathbf{b}_2 &= \mathbf{a}_2 \cdot \mathbf{b}_1 = 0, \end{aligned} \quad (\text{I.4})$$

so

$$\begin{aligned} \mathbf{b}_1 &= \frac{2\pi}{a} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{3}} \end{pmatrix} \\ \mathbf{b}_2 &= \frac{2\pi}{a} \begin{pmatrix} 1 \\ -\frac{1}{\sqrt{3}} \end{pmatrix}. \end{aligned} \quad (\text{I.5})$$




Figure I.2: Graphene Brillouin Zone

ic for primi-  
Z

in primitive

in: what sym-  
es does EG-X  
? How does  
influence BZ?

Points of high symmetry in the Brillouin zone are:

$$\Gamma = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (\text{I.6})$$

$$\text{M} = \frac{\pi}{a} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{3}} \end{pmatrix} \quad (\text{I.7})$$

$$\text{K} = \frac{4\pi}{3a} \begin{pmatrix} 1 \\ 0 \end{pmatrix} . \quad (\text{I.8})$$

## I.2 Band structure

aka why is the model interesting?

The detailed derivation can be found in appendix A.

$$H_0 = \sum_{\mathbf{k}, \sigma} \begin{pmatrix} c_{\mathbf{k}, \sigma}^{A, \dagger} & c_{\mathbf{k}, \sigma}^{B, \dagger} & d_{\mathbf{k}, \sigma}^{\dagger} \end{pmatrix} \begin{pmatrix} 0 & f_{Gr} & V \\ f_{Gr}^* & 0 & 0 \\ V & 0 & f_X \end{pmatrix} \begin{pmatrix} c_{\mathbf{k}, \sigma}^A \\ c_{\mathbf{k}, \sigma}^B \\ d_{\mathbf{k}, \sigma} \end{pmatrix} \quad (\text{I.9})$$

## II Conclusion



# A EG-X Hamiltonian in Reciprocal Space

In the following chapter, the model Hamiltonian

$$\begin{aligned}
 H_0 = & -t_X \sum_{\langle ij \rangle, \sigma} d_{i, \sigma}^\dagger d_{j, \sigma} - t_{Gr} \sum_{\langle ij \rangle, \sigma} \left( c_{i, \sigma}^{(A), \dagger} c_{j, \sigma}^{(B)} + c_{j, \sigma'}^{(B), \dagger} c_{i, \sigma}^{(A)} \right) \\
 & + V \sum_{i, \sigma} d_{i, \sigma}^\dagger c_{i, \sigma}^{(A)} + \text{h.c.}
 \end{aligned} \tag{A.1}$$

will be treated via Fourier transform.

The first step is to write out the sums over nearest neighbors  $\langle i, j \rangle$  explicitly, writing  $\delta_X, \delta_\epsilon$  ( $\epsilon = A, B$ ) for the vectors to the nearest neighbors of the X atoms and Graphene A, B sites. Doing the calculation for example of the X atoms:

$$-t_X \sum_{\langle ij \rangle, \sigma} (d_{i, \sigma}^\dagger d_{j, \sigma} + d_{j, \sigma}^\dagger d_{i, \sigma}) = -\frac{t_X}{2} \sum_{i, \sigma} \sum_{\delta_X} d_{i, \sigma}^\dagger d_{i+\delta_X, \sigma} - \frac{t_X}{2} \sum_{j, \sigma} \sum_{\delta_X} d_{j, \sigma}^\dagger d_{j+\delta_X, \sigma} \tag{A.2}$$

$$= -t_X \sum_{i, \sigma} \sum_{\delta_X} d_{i, \sigma}^\dagger d_{i+\delta_X, \sigma} \tag{A.3}$$

The factor 1/2 in eq. (A.2) is to account for double counting when going to the sum over all lattice sites  $i$ . By relabeling  $j \rightarrow i$  in the second sum, we then get eq. (A.3). Using now the discrete Fourier transform

$$\begin{aligned}
 c_i &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}} \\
 c_i^\dagger &= \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}}^\dagger
 \end{aligned} \tag{A.4}$$

with the completeness relation:

$$\sum_i e^{i\mathbf{k}\mathbf{r}_i} e^{-i\mathbf{k}'\mathbf{r}_i} = N\delta_{\mathbf{k},\mathbf{k}'}, \quad (\text{A.5})$$

eq. (A.3) reads:

$$-t_X \frac{1}{N} \sum_{i,\sigma} \sum_{\delta_X} d_{i,\sigma}^\dagger d_{i+\delta_X,\sigma'} = -t_X \frac{1}{N} \sum_{i,\sigma,\sigma'} \sum_{\delta_X} \sum_{\mathbf{k},\mathbf{k}'} e^{-i\mathbf{k}\mathbf{r}_i} d_{\mathbf{k},\sigma}^\dagger e^{i\mathbf{k}'\mathbf{r}_i} e^{i\mathbf{k}'\delta_X} d_{\mathbf{k}',\sigma'} \quad (\text{A.6})$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\sigma,\sigma'} \sum_{\delta_X} d_{\mathbf{k},\sigma}^\dagger e^{i\mathbf{k}'\delta_X} d_{\mathbf{k}',\sigma'} \sum_i e^{-i\mathbf{k}\mathbf{r}_i} e^{i\mathbf{k}'\mathbf{r}_i} \quad (\text{A.7})$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\sigma,\sigma'} \sum_{\delta_X} d_{\mathbf{k},\sigma}^\dagger e^{i\mathbf{k}'\delta_X} d_{\mathbf{k}',\sigma'} N\delta_{\mathbf{k},\mathbf{k}'} \quad (\text{A.8})$$

$$= -t_X \sum_{\mathbf{k},\sigma,\sigma'} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k},\sigma'} \sum_{\delta_X} e^{i\mathbf{k}\delta_X} \quad (\text{A.9})$$

The nearest neighbours for X atoms are the vectors  $\delta_{AA,i}$  from section I.1. With that, we can calculate:

$$f_X(\mathbf{k}) = -t_X \sum_{\delta_X} e^{i\mathbf{k}\delta_X} \quad (\text{A.10})$$

$$= -t_X \left( e^{ia(\frac{k_x}{2} + \frac{\sqrt{3}k_y}{2})} + e^{iak_x} + e^{ia(\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} \right) \quad (\text{A.11})$$

$$+ e^{ia(-\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} + e^{-iak_x} + e^{ia(-\frac{k_x}{2} + \frac{\sqrt{3}k_y}{2})} \quad (\text{A.12})$$

$$= -t_X \left( 2 \cos(ak_x) + 2e^{ia\frac{\sqrt{3}k_y}{2}} \cos\left(\frac{a}{2}k_x\right) + 2e^{-ia\frac{\sqrt{3}k_y}{2}} \cos\left(\frac{a}{2}k_x\right) \right) \quad (\text{A.13})$$

$$= -2t_X \left( \cos(ak_x) + 2 \cos\left(\frac{a}{2}k_x\right) \cos\left(\sqrt{3}\frac{a}{2}k_y\right) \right) \quad (\text{A.14})$$

We can do the same for the hopping between Graphene sites, for example :

$$-t_{\text{Gr}} \sum_{\langle ij \rangle, \sigma\sigma'} c_{i,\sigma}^{(A)\dagger} c_{j,\sigma'}^{(B)} = -t_{\text{Gr}} \sum_{i,\sigma,\sigma'} \sum_{\delta_{AB}} c_{i,\sigma}^{(A)\dagger} c_{i+\delta_{AB},\sigma'}^{(B)} \quad (\text{A.15})$$

$$= -t_{\text{Gr}} \sum_{\mathbf{k},\sigma,\sigma'} c_{\mathbf{k},\sigma}^{(A)\dagger} c_{\mathbf{k},\sigma'}^{(B)} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (\text{A.16})$$



We note

$$\sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} = \left( \sum_{\delta_{BA}} e^{i\mathbf{k}\delta_{BA}} \right)^* = \sum_{\delta_{BA}} e^{-i\mathbf{k}\delta_{BA}} \quad (\text{A.17})$$

and calculate

$$f_{Gr} = -t_{Gr} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (\text{A.18})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{i\frac{a}{2\sqrt{3}}(\sqrt{3}k_x - k_y)} + e^{i\frac{a}{2\sqrt{3}}(-\sqrt{3}k_x - k_y)} \right) \quad (\text{A.19})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{-i\frac{a}{2\sqrt{3}}k_y} \left( e^{i\frac{a}{2}k_x} + e^{-i\frac{a}{2}k_x} \right) \right) \quad (\text{A.20})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + 2e^{-i\frac{a}{2\sqrt{3}}k_y} \cos\left(\frac{a}{2}k_x\right) \right) \quad (\text{A.21})$$

All together, we get:

$$H_0 = \sum_{\mathbf{k}, \sigma, \sigma'} \begin{pmatrix} c_{\mathbf{k}, \sigma}^{A, \dagger} & c_{\mathbf{k}, \sigma}^{B, \dagger} & d_{\mathbf{k}, \sigma}^{\dagger} \end{pmatrix} \begin{pmatrix} 0 & f_{Gr} & V \\ f_{Gr}^* & 0 & 0 \\ V & 0 & f_X \end{pmatrix} \begin{pmatrix} c_{\mathbf{k}, \sigma}^A \\ c_{\mathbf{k}, \sigma}^B \\ d_{\mathbf{k}, \sigma} \end{pmatrix} \quad (\text{A.22})$$



## Bibliography

- [1] G. Yang et al. “Structure of Graphene and Its Disorders: A Review”. In: *Science and Technology of Advanced Materials* 19.1 (Aug. 2018), pp. 613–648. ISSN: 1468-6996. DOI: 10.1080/14686996.2018.1494493.



## List of Symbols



# Acknowledgement

I thank my supervisor Professor Tim Wehling for allowing me to get a glimpse into the fascinating research he and his research group do and for the opportunity to learn a lot about condensed matter in the past 5 months.

I also thank Michael Winter and Jan Berges for direct support, scripts, data, L<sup>A</sup>T<sub>E</sub>X templates and much more as well as the whole group ‘Computational Condensed Matter Theory’ for welcoming me into their open and productive atmosphere.

I thank all developers of free software, especially the ones I used in this thesis: L<sup>A</sup>T<sub>E</sub>X, PYTHON, NUMPY, MATPLOTLIB, GIT.

I thank my family for support through my all my studies up until now.

I especially thank Liv for massive amounts of love and support during the writing of this thesis.