

# Dressed Graphene Hamiltonian in Reciprocal Space

1

In this chapter, the Hamiltonian

$$H_0 = -t_X \sum_{\langle ij \rangle, \sigma} d_{i, \sigma}^\dagger d_{j, \sigma} - t_{Gr} \sum_{\langle ij \rangle, \sigma} c_{i, \sigma}^{(A), \dagger} c_{j, \sigma}^{(B)} + V \sum_{i, \sigma \sigma'} d_{i, \sigma}^\dagger c_{i, \sigma'}^{(A)} + \text{h.c.} \quad (1.1)$$

from ?? will be treated to obtain the electronic band structure shown in the chapter. The first step is to write out the sums over nearest neighbors  $\langle i, j \rangle$  explicitly, writing  $\delta_X, \delta_\epsilon$  ( $\epsilon = A, B$ ) for the vectors to the nearest neighbors of the X atoms and Graphene A, B sites. For example, for the X atoms this gives:

$$-t_X \sum_{\langle ij \rangle, \sigma} (d_{i, \sigma}^\dagger d_{j, \sigma} + d_{j, \sigma}^\dagger d_{i, \sigma}) = -\frac{t_X}{2} \sum_{i, \delta_X, \sigma} d_{i, \sigma}^\dagger d_{i+\delta_X, \sigma} - \frac{t_X}{2} \sum_{j, \delta_X, \sigma} d_{j, \sigma}^\dagger d_{j+\delta_X, \sigma} \quad (1.2)$$

$$= -t_X \sum_{i, \sigma} \sum_{\delta_X} d_{i, \sigma}^\dagger d_{i+\delta_X, \sigma} . \quad (1.3)$$

The factor 1/2 in eq. (1.2) is to account for double counting when going to the sum over all lattice sites  $i$ . By relabeling  $j \rightarrow i$  in the second sum, the two sum are the same and eq. (1.3) is obtained. Now, using the discrete Fourier transform

$$c_i = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}}, \quad c_i^\dagger = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}}^\dagger \quad (1.4)$$

with the completeness relation

$$\sum_i e^{i\mathbf{k}\mathbf{r}_i} e^{-i\mathbf{k}'\mathbf{r}_i} = N \delta_{\mathbf{k}, \mathbf{k}'} , \quad (1.5)$$

eq. (1.3) reads:

$$-t_X \frac{1}{N} \sum_{i,\sigma} \sum_X d_{i,\sigma}^\dagger d_{i+\delta_X,\sigma} = -t_X \frac{1}{N} \sum_{i,\sigma} \sum_{\mathbf{k},\mathbf{k}',\delta_X} (e^{-i\mathbf{k}\mathbf{r}_i} d_{\mathbf{k},\sigma}^\dagger) (e^{i\mathbf{k}'\mathbf{r}_i} e^{i\mathbf{k}'\delta_X} d_{\mathbf{k}',\sigma}) \quad (1.6)$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\delta_X,\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k}',\sigma} e^{i\mathbf{k}'\delta_X} \sum_i e^{-i\mathbf{k}\mathbf{r}_i} e^{i\mathbf{k}'\mathbf{r}_i} \quad (1.7)$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k}',\sigma} \sum_{\delta_X} e^{i\mathbf{k}'\delta_X} (N \delta_{\mathbf{k},\mathbf{k}'}) \quad (1.8)$$

$$= -t_X \sum_{\mathbf{k},\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k},\sigma} \sum_{\delta_X} e^{i\mathbf{k}\delta_X} \quad (1.9)$$

This is now diagonal in  $\mathbf{k}$  space. The sum over  $\delta_X$  can be explicitly calculated using the fact that the nearest neighbours vectors  $\delta_X$  for the X atoms are the vectors  $\delta_{AA,i}$  from ??, for example

$$\mathbf{k} \cdot \delta_{AA,1} = \begin{pmatrix} k_x \\ k_y \end{pmatrix} \cdot \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} = k_x + \sqrt{3}k_y \quad (1.10)$$

Clear up  
definition  
NN vectors  
and results

$$f_X(\mathbf{k}) = -t_X \sum_{\delta_X} e^{i\mathbf{k}\delta_X} \quad (1.11)$$

$$= -t_X \left[ e^{ia(\frac{k_x}{2} + \frac{\sqrt{3}k_y}{2})} + e^{iak_x} + e^{ia(\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} \right] \quad (1.12)$$

$$+ e^{ia(-\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} + e^{-iak_x} + e^{ia(-\frac{k_x}{2} + \frac{\sqrt{3}k_y}{2})} \quad (1.13)$$

$$= -t_X \left( 2 \cos(ak_x) + 2e^{ia\frac{\sqrt{3}k_y}{2}} \cos(\frac{a}{2}k_x) + 2e^{-ia\frac{\sqrt{3}k_y}{2}} \cos(\frac{a}{2}k_x) \right) \quad (1.14)$$

$$= -2t_X \left( \cos(ak_x) + 2 \cos(\frac{a}{2}k_x) \cos(\sqrt{3}\frac{a}{2}k_y) \right) . \quad (1.15)$$

The same can be done for the hopping between Graphene sites, for example :

$$-t_{Gr} \sum_{\langle ij \rangle, \sigma \sigma'} c_{i,\sigma}^{(A)\dagger} c_{j,\sigma'}^{(B)} = -t_{Gr} \sum_{i,\sigma \sigma'} \sum_{\delta_{AB}} c_{i,\sigma}^{(A)\dagger} c_{i+\delta_{AB},\sigma'}^{(B)} \quad (1.16)$$

$$= -t_{Gr} \sum_{\mathbf{k},\sigma,\sigma'} c_{\mathbf{k},\sigma}^{(A)\dagger} c_{\mathbf{k},\sigma'}^{(B)} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (1.17)$$

with again the sum over  $\delta_{AB}$

$$f_{\text{Gr}}(\mathbf{k}) = -t_{\text{Gr}} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (1.18)$$

$$= -t_{\text{Gr}} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{i\frac{a}{2\sqrt{3}}(\sqrt{3}k_x - k_y)} + e^{i\frac{a}{2\sqrt{3}}(-\sqrt{3}k_x - k_y)} \right) \quad (1.19)$$

$$= -t_{\text{Gr}} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{-i\frac{a}{2\sqrt{3}}k_y} \left( e^{i\frac{a}{2}k_x} + e^{-i\frac{a}{2}k_x} \right) \right) \quad (1.20)$$

$$= -t_{\text{Gr}} \left( e^{i\frac{a}{\sqrt{3}}k_y} + 2e^{-i\frac{a}{2\sqrt{3}}k_y} \cos\left(\frac{a}{2}k_x\right) \right) \quad (1.21)$$

We note

Show that!

$$\sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} = \left( \sum_{\delta_{BA}} e^{i\mathbf{k}\delta_{BA}} \right)^* = \sum_{\delta_{BA}} e^{-i\mathbf{k}\delta_{BA}} \quad (1.22)$$

All in all:

$$H_0 = \sum_{\mathbf{k}, \sigma, \sigma'} \begin{pmatrix} c_{\mathbf{k}, \sigma}^{A, \dagger} & c_{\mathbf{k}, \sigma}^{B, \dagger} & d_{\mathbf{k}, \sigma}^{\dagger} \end{pmatrix} \begin{pmatrix} 0 & f_{\text{Gr}} & V \\ f_{\text{Gr}}^* & 0 & 0 \\ V & 0 & f_X \end{pmatrix} \begin{pmatrix} c_{\mathbf{k}, \sigma}^A \\ c_{\mathbf{k}, \sigma}^B \\ d_{\mathbf{k}, \sigma} \end{pmatrix} \quad (1.23)$$