

Superconductivity

1

In this chapter I review theoretical concepts needed for understanding superconductivity and introduce the tools used to study superconductivity in the later chapters. There are many textbooks covering these topics which can be referenced for a more detailed treatment, such as refs. [1–5].

Macroscopically, the superconducting state can be described by a spontaneous breaking of a $U(1)$ phase rotation symmetry that is associated with an order parameter. The theory of spontaneous symmetry breaking and associated phase transitions is Ginzburg-Landau theory discussed in section 1.1, following refs. [1, 6]. Ginzburg-Landau theory introduces two length scales: the coherence length ξ_0 describing the length scale of amplitude variations of the order parameter and the London penetration depth λ_L , which is connected to energy cost of phase variations of the order parameter. They also connect to the energy gap Δ and the condensate stiffness D_S , which are often competing energy scales in superconductors. The interplay of these length (energy) scales determine the macroscopic properties of a superconductors, so there is a great interest in accessing them in computational ways. To this end, section 1.1 also introduces a theoretical framework based on Cooper pairs with finite momentum [7] that will be used in later chapters to calculate these length scales from microscopic theories.

Ginzburg Landau theory is a macroscopic theory, but it can be connected to microscopic theories: if a theory finds an expression for the order parameter describing the breakdown of symmetry, it can be connected to quantities expressed by Ginzburg-Landau theory. One such theory to describe superconductivity from a microscopic perspective is BCS (Bardeen-Cooper-Schrieffer) theory in section 1.2, which is A method to treat local interactions non-perturbatively is DMFT (Dynamical Mean Field Theory). Section 1.3 briefly introduces the Greens function method to treat many-body problems and outlines the DMFT self-consistency cycle.

Furthermore, section 1.4 introduces an emerging perspective in the study of novel superconductors: it turns out that the superfluid weight is connected to

a quantity of the electronic band structure called the quantum metric [8, 9], which is connected to

1.1 Ginzburg-Landau Theory of Superconductivity

Spontaneous Symmetry Breaking and Order Parameter

Symmetries are a powerful concept in physics. Noethers theorem [10] connects the symmetries of physical theories to associated conservation laws. An interesting facet of symmetries in physical theories is the fact, that a ground state of a system must not necessarily obey the same symmetries of its Hamiltonian, i.e. for a symmetry operation that is described by a unitary operator U , the Hamiltonian commutes with U (which results in expectation values of the Hamiltonian being invariant under the symmetry operation) but the states $|\phi\rangle$ and $U|\phi\rangle$ are different. This phenomenon is called spontaneous symmetry breaking and the state $|\phi\rangle$ is said to be symmetry-broken.

One consequence of this fact is that for a given symmetry-broken state $|\phi\rangle$, there exists multiple states that can be reached by repeatedly applying U to $|\phi\rangle$ and all have the same energy. To differentiate the symmetry-broken states an operator can be defined that has all these equivalent states as eigenvectors with different eigenvalues and zero expectation value for symmetric states. This is the microscopic notion of an order parameter.

The original notion of an order parameter was motivated from macroscopic observables that can then be related to the microscopic order parameter operator introduced above. Macroscopically I characterize the symmetry breaking by an order parameter Ψ which generally can be a complex-valued vector that becomes non-zero below the transition temperature T_C

$$|\Psi| = \begin{cases} 0 & T > T_C \\ |\Psi_0| > 0 & T < T_C \end{cases} . \quad (1.1)$$

In the example of a ferromagnet, a finite magnetization of a material is associated with a finite expectation value for the z-component of the spin operator, $m_z = \langle \hat{S}_z \rangle$. The order parameter describes the ‘degree of order’ [11]. Similarly to a magnetically ordered state, the SC state is characterized by an order parameter. The theory of phase transitions in superconductors was developed by Ginzburg and Landau [12]. Landau theory and conversely

Ginzburg-Landau theory is not concerned with the microscopic properties of the order parameter, but describes the changes in thermodynamic properties of matter with the development of an order parameter.

Still explain a little bit what the physical interpretation of the OP is

Landau and Ginzburg-Landau Theory

The free energy is a thermodynamic quantity:

$$F = E - TS \quad (1.2)$$

with the energy of the system E , temperature T and entropy S . A system in thermodynamic equilibrium has minimal free energy. The fundamental idea underlying Landau theory is to write the free energy $F[\Psi]$ as function of the order parameter Ψ and expand it as a polynomial:

$$F_L[\Psi] = \int d^d x f_L[\Psi], \quad (1.3)$$

where

$$f_L[\Psi] = \frac{r}{2}\Psi^2 + \frac{u}{4}\Psi^4 \quad (1.4)$$

is called the free energy density. Provided the parameters r and u are greater than 0, there is a minimum of $f_L[\Psi]$ that lies at $\Psi = 0$. Landau theory assumes that at the phase transition temperature T_C the parameter r changes sign, so it can be written in first order as

$$r = a(T - T_C). \quad (1.5)$$

Figure 1.1a shows the free energy as a function of a single-component, real order parameter Ψ and it illustrates the essence of Landau theory: there are two cases for the minima of the free energy f

$$\Psi = \begin{cases} 0 & T \geq T_C \\ \pm \sqrt{\frac{a(T_C - T)}{u}} & T < T_C \end{cases}, \quad (1.6)$$

so there is a for $T < T_C$ there are two minima corresponding to ground states with broken symmetry. When the order parameter can be calculated from some microscopic theory, the critical temperature T_C can be extracted from the behavior of the order parameter near T_C via a linear fit of

$$|\Psi|^2 \propto T_C - T. \quad (1.7)$$




Figure 1.1: (a) Landau free energy and (b) Mexican hat potential

Generalizing this from a one to an n -component order parameters is straightforward. One example is the complex or two component order parameter that will become important for

$$\Psi = \Psi_1 + i\Psi_2 = |\Psi|e^{i\phi}. \quad (1.8)$$

The Landau free energy then takes the form

$$f_L[\Psi] = r\Psi^*\Psi + \frac{u}{2}(\Psi^*\Psi)^2 = r|\Psi|^2 + \frac{u}{2}|\Psi|^4 \quad (1.9)$$

with again

$$r = a(T_C - T). \quad (1.10)$$

Instead of the two minima, the free energy here is rotational symmetry, because it is independent of the phase of the order parameter:

$$f_L[\Psi] = f_L[e^{i\phi}\Psi]. \quad (1.11)$$

This gives the so called ‘Mexican hat’ potential shown in fig. 1.1b. In this potential, the order parameter can be rotated continuously from one symmetry-broken state to another.

In 1950, Ginzburg and Landau published their theory of superconductivity, based on Landau’s theory of phase transitions [12]. Where Landau theory as

described above has an uniform order parameters, Ginzburg-Landau theory accounts for it being inhomogeneous, so an order parameter with spatially varying amplitude or direction. This in turn leads to the order parameter developing a fixed phase, which is the underlying mechanism of the superflow in superconductors.

Ginzburg-Landau theory can be developed for a general n -component order parameter, but in superfluids and superconductors the order parameter is complex, i.e. two-component. The Ginzburg-Landau free energy for a complex order parameter is

$$f_{GL}[\Psi, \Delta\Psi] = \frac{\hbar^2}{2m^*} |\Delta\Psi|^2 + r|\Psi|^2 + \frac{u}{2} |\Psi|^4, \quad (1.12)$$

where the gradient term $\Delta\Psi$ is added in comparison to the Landau free energy. The prefactor $\frac{\hbar^2}{2m^*}$ is chosen to illustrate the interpretation of the Ginzburg-Landau free energy as the energy of a condensate of bosons, where the gradient term $|\Delta\Psi|^2$ is the kinetic energy. The free energy in eq. (1.12) is sensitive to a twist of the phase of the order parameter. Substituting the expression $\Psi = |\Psi|e^{i\phi}$, the gradient term reads

$$\Delta\Psi = (\Delta|\Psi| + i\Delta\phi|\Psi|)e^{i\phi}. \quad (1.13)$$

With that, eq. (1.12) becomes

$$f_{GL} = \frac{\hbar^2}{2m^*} |\Psi|^2 (\Delta\phi)^2 + \left[\frac{\hbar^2}{2m^*} (\Delta|\Psi|)^2 + r|\Psi|^2 + \frac{u}{2} |\Psi|^4 \right]. \quad (1.14)$$

Now the contributions of phase and amplitude variations are split up: the first term describes energy cost of variations in the phase of the order parameter and the second term describes energy cost of variations in the magnitude of the order parameter. Twisting the phase of the condensate is associated with an energy cost. This energy cost is characterized by the stiffness D_s

The dominating fluctuation is determined by the ratio of the factors $\frac{\hbar^2}{2m^*}$ and r , which has the dimension Length^2 , from which define the correlation length.

$$\xi = \sqrt{\frac{\hbar^2}{2m^*|r|}} = \xi_0 \left(1 - \frac{T}{T_C}\right)^{-\frac{1}{2}} \quad (1.15)$$

Introduce superfluid density properly

Energy cost of phase bending via GL energy

Explanation phase stiffness

Explanation superfluid weight as linear response

where I define the zero temperature value as the coherence length $\xi_0 = \xi(T = 0) = \sqrt{\frac{\hbar^2}{2maT_C}}$. On length scales above ξ , the physics is entirely controlled by the phase degrees of freedom, i.e.

$$f_{GL} = \frac{\hbar^2}{2m^*} |\Psi|^2 (\Delta\phi)^2 + \text{const.} \quad (1.16)$$

Take case of frozen amplitude fluctuations, i.e. $\Delta|\Psi(\mathbf{r})| = 0$. Stationary point condition for eq. (1.14) gives:

$$|\Psi| = |\Psi_0| \sqrt{1 - \xi^2 |\Delta\phi(\mathbf{r})|^2} \quad (1.17)$$

This shows that the superconducting order gets suppressed and eventually destroyed by short-ranged (below ξ) phase fluctuations. By introducing a particular form of phase fluctuations $\phi = \mathbf{q} \cdot \mathbf{r}$ into a microscopic model, it is possible to probe this breakdown of superconductivity and thus gain insight into the nature of superconductivity, in particular this gives access to ξ .

Work over paragraph

Superconductors: charged superfluids, coupling to electromagnetic fields. Free energy with minimal coupling to an electromagnetic field:

$$f_{GL}[\Psi, \mathbf{A}] = \frac{\hbar^2}{2m^*} \left| \left(\Delta - \frac{ie^*}{\hbar} \mathbf{A} \right) \Psi \right|^2 + r|\Psi|^2 + \frac{u}{2} |\Psi|^4 \quad (1.18)$$

Describes really two intertwined Ginzburg-Landau theories for Ψ and \mathbf{A} respectively. This mean there are two length scales, the coherence length ξ governing amplitude fluctuations of Ψ and the London penetration depth λ_L , which determines the distance magnetic fields penetrate into the superconductor. Can get the current density from the stationary point condition of the free energy for the vector potential \mathbf{A} :

$$\frac{\delta f_{GL}}{\delta \mathbf{A}} = 0 = -\mathbf{j} + \frac{1}{\mu_0} \nabla \times \mathbf{B} \quad (1.19)$$

with the supercurrent density

$$\mathbf{j} = -i \frac{e\hbar}{m^*} (\Psi^* \Delta \Psi - \Psi \Delta \Psi^*) - \frac{4e^2}{m^*} |\Psi|^2 \mathbf{A} . \quad (1.20)$$

Introducing the OP with phase $\Psi = |\Psi|e^{i\phi}$:

$$\mathbf{j} = 2e|\Psi|^2 \frac{\hbar}{m^*} \left(\nabla\phi - \frac{2\pi}{\Phi_0} \mathbf{A} \right) \quad (1.21)$$

with the magnetic flux quantum $\Phi_0 = \frac{\pi\hbar}{e}$. Shows that not only an applied field \mathbf{A} can induce a supercurrent, but also the twist of the phase of the condensate. Rewriting using $|\Psi|^2 = n_s/2$ and introducing the superfluid velocity \mathbf{v}_S :

$$\mathbf{j} = en_s \mathbf{v}_S \quad (1.22)$$

where even for $\mathbf{A} = 0$, there is finite current because of the phase twist of the condensate ground state (which is quite remarkable).

Superconducting Length Scales

One of the challenges in achieving high-temperature superconductivity is the fact that the two intrinsic energy scales of superconductors i.e. the pairing amplitude and the phase coherence often compete. Can be seen in the phenomenon of BCS-BEC crossover physics [13]. The picture of this crossover is the following: for a small attractive interaction, pairs of electrons are very loosely bound and mobile, while for a stronger interaction the pairs are bound together stronger and are not mobile, because hopping of a pair would involve a virtual hopping, thus breaking up the pair. This is highly suppressed. The crossover region between these two regimes is the BCS-BEC crossover. The energy scales characterizing the crossover are the superconducting gap/order parameter describing how strong the order is and the superfluid weight describing how mobile the Cooper pairs are. These two energy scales are equivalently defined via the coherence length ξ_0 and the London penetration depth introduced in section 1.1. To understand the physics of the BCS-BEC crossover and eventually use it to optimize current and future superconductors it is thus imperative to have access to the superconducting length scales in ab initio approaches Witt et al. introduced a framework for doing this [7]. As already discussed in the context of eq. (1.17), strong phase fluctuations destroy superconducting order. A particular choice of phase fluctuations would be

$$\phi(\mathbf{r}) = \mathbf{q} \cdot \mathbf{r} \quad (1.23)$$

Section about BCS-BEC crossover? Can especially talk about what is optimised!

Write introduction better

which corresponds to Cooper pairs with a finite momentum \mathbf{q} . In most materials: Cooper pairs do not carry finite center-of-mass momentum. In presence of e.g. external fields or magnetism: SC states with FMP might arise [14–16]

Procedure in the paper: enforce FMP states via constraints on pair-center-of-mass momentum \mathbf{q} , access characteristic length scales ξ_0, λ_L through analysis of the momentum and temperature-dependent OP. FF-type pairing with Cooper pairs carrying finite momentum:

$$\Psi_{\mathbf{q}}(\mathbf{r}) = |\Psi_{\mathbf{q}}|e^{i\mathbf{q}\cdot\mathbf{r}} \quad (1.24)$$

Then the free energy density eq. (1.12) is

$$f_{GL}[\Psi_{\mathbf{q}}] = r|\Psi_{\mathbf{q}}|^2 + \frac{u}{2}|\Psi_{\mathbf{q}}|^4 + \frac{\hbar^2 q^2}{2m^*}|\Psi_{\mathbf{q}}|^2 \quad (1.25)$$

Stationary point of the system:

$$\frac{\delta f_{GL}}{\delta \Psi_{\mathbf{q}}^*} = 2\Psi_{\mathbf{q}} [r(1 - \xi^2 q^2) + u|\Psi_{\mathbf{q}}|^2] = 0 \quad (1.26)$$

which results in the \mathbf{q} -dependence of the OP

$$|\Psi_{\mathbf{q}}|^2 = |\Psi_0|^2 (1 - \xi(T)^2 q^2) \quad (1.27)$$

For some value, SC order breaks down, $\psi_{\mathbf{q}_c} = 0$, because the kinetic energy from phase modulation exceeds the gain in energy from pairing. In GL theory: $q_c = \xi(T)^{-1}$. The temperature dependence of the OP and extracted $\xi(T)$ gives access to the coherence length via eq. (1.15)


$$\xi(T) = \xi_0 \left(1 - \frac{T}{T_C}\right)^{-\frac{1}{2}} \quad (1.28)$$

The Cooper pair

The momentum of the Cooper pairs entails a charge supercurrent $\mathbf{j}_{\mathbf{q}}$. For small q

The depairing current is an upper boundary for the maximal current that can flow through a material, also called the critical current \mathbf{j}_c . The value of \mathbf{j}_c is strongly dependent on the geometry of the sample [17, 18], so \mathbf{j}_{dp} is not

Depairing
current from
FMP

Figure 1.2: *a and b*

necessarily experimentally available, but it can be used to calculate the London penetration depth [2]

$$\lambda_L(T) = \sqrt{\frac{\Phi_0}{3\sqrt{3}\pi\mu_0\zeta(T)j_{dp}(T)}} = \lambda_{L,0} \left(1 - \left(\frac{T}{T_C}\right)^4\right)^{-\frac{1}{2}} \quad (1.29)$$

1.2 Bardeen-Cooper-Schrieffer Theory

It took nearly 50 years after the first discovery of superconductivity in mercury by Heike Kamerlingh Onnes in 1911 [19] for the first microscopic description of this phenomenon to be published in 1957 by John Bardeen, Leon Cooper and J. Robert Schrieffer [20].

The BCS (Bardeen-Cooper-Schrieffer) description of superconductivity is based on the fact that the Fermi sea is unstable towards development of bound pairs under arbitrarily small attraction [21]. The origin of the attractive interaction $V_{\mathbf{k},\mathbf{k}'}$, which Bardeen, Cooper and Schrieffer identified as a retarded electron-phonon interaction [20].

There exist many textbooks tackling BCS theory from different angles, such as refs. [1, 2]. This section gives an introduction to the relevant physics of BCS theory as originally proposed, then derives the

BCS Hamiltonian

Where does this formula come from? Second London equation

D_S

What else can be done with the FMP method?

Connection of the FMP method to linear response techniques

Better introduction

Work over paragraph

BCS-Hamiltonian:

$$H_{\text{BCS}} = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \quad (1.30)$$

This Hamiltonian can be solved exactly using a mean field approach, because it involves an interaction at zero momentum and thus infinite range. Order parameter in mean field BCS theory is the pairing amplitude

$$\Delta = -\frac{U}{N_{\mathbf{k}}} \sum_{\mathbf{k}} \langle c_{-\mathbf{k}\downarrow} c_{\mathbf{k}\uparrow} \rangle = -U \langle c_{-\mathbf{r}=0\downarrow} c_{\mathbf{r}=0\uparrow} \rangle \simeq U\Psi. \quad (1.31)$$

A finite Δ corresponds to the pairing introduced above: there is a finite expectation value for a coherent creation/annihilation of a pair of electrons with opposite momentum and spin. A finite Δ also introduces a band gap into the spectrum. BCS theory brings multiple aspects together: concept of paired electrons with the pairing amplitude being the order parameter in SC, an explanation for the attractive interaction overcoming Coulomb repulsion and a model Hamiltonian that very elegantly captures the essential physics.

It is very successful in two ways: on the one hand it could quantitatively predict effects in the SCs known at the time, for example the Hebel-Slichter peak that was measured in 1957 [22, 23] and the band gap measured by Giaever in 1960 [24]. On the other hand, it established electronic pairing, i.e. the picture of a quantum-mechanical wave function with a defined phase as already described by Fritz London in 1937 [25] as the microscopic mechanism behind SC. This picture still holds today even for high T_C /unconventional superconductors, so SCs that cannot be described by BCS theory [26].

Multiband BCS Theory

The Hubbard model is the simplest model for interacting electron systems. It goes back to works by Hubbard [27], Kanamori [28] and Gutzweiler [29].

$$H_{\text{int}} = U \sum_i c_{i,\uparrow}^\dagger c_{i,\downarrow}^\dagger c_{i,\downarrow} c_{i,\uparrow} \quad (1.32)$$

where $U > 0$. It describes a repulsive interaction between electrons of different spin at the same lattice site.

Besides

Other pairing interactions can be taken, gives explanations for a lot of different SCs

Kinetic term as well

[30]

The Hubbard model in the form of eq. (1.32) can be extended in a multitude of ways to model a variety of physical system. Here: extension to multiple orbitals (i.e. atoms in the unit cell for lattice systems) and an attractive interaction, i.e. a negative U . Physical motivation for taking a negative- U Hubbard model: electrons can experience a local attraction interaction, for example through electrons coupling with phononic degrees of freedom or with electronic excitations that can be described as bosons [31]. The form of the interaction term is then:

$$H_{\text{int}} = - \sum_{i,\alpha} U_{\alpha} c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \quad (1.33)$$

where α counts orbitals and the minus sign in front is taken so that $U > 0$ now corresponds to an attractive interaction (this is purely convention).

There are a multitude of ways to derive a mean field description of a given interacting Hamiltonian. Very rigorous in path integral formulations as saddle points, given for example in ref. [1]. The review follows ref. [32]. A more intuitive way based on ref. [3] discussed here looks at the operators and which one are small.

Look at interaction term eq. (1.33). Mean-field approximation (here specifically for superconductivity i.e. pairing): operators do not deviate much from their average value, i.e. the deviation operators

$$d_{i,\alpha} = c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} - \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle \quad (1.34)$$

$$e_{i,\alpha} = c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} - \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle \quad (1.35)$$

are small (don't contribute much to expectation values and correlation functions), so that in the interaction part of the Hamiltonian

$$H_{\text{int}} = - \sum_{i,\alpha} U_{\alpha} c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \quad (1.36)$$

$$= - \sum_{i,\alpha} U_{\alpha} (d_{i,\alpha}^{\dagger} + \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle) (e_{i,\alpha} + \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle) \quad (1.37)$$

$$= - \sum_{i,\alpha} U_{\alpha} (d_{i,\alpha} e_{i,\alpha} + d_{i,\alpha} \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle + e_{i,\alpha} \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle$$

$$+ \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle) \quad (1.39)$$

Some relevance of the repulsive Hubbard model

There are some more specific papers to the specific mechanisms (and also some more mechanism), could cite these here and say some more things

Order of operators? -> also in all other equations!

there are other combinations, talk about that

deviations with small deltas

the first term is quadratic in the deviation and can be neglected. Thus arrive at the approximation

$$H_{\text{int}} \approx - \sum_{i,\alpha} U_{\alpha} \left(d_{i,\alpha} \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle + e_{i,\alpha} \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle + \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle \right) \quad (1.40)$$

$$= - \sum_{i,\alpha} U_{\alpha} (c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle + c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle) \quad (1.41)$$

$$- \langle c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} \rangle \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle) \quad (1.42)$$

$$= \sum_{i,\alpha} \left(\Delta_{i,\alpha} c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} + \Delta_{i,\alpha}^* c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} - \frac{|\Delta_{i,\alpha}|^2}{U_{\alpha}} \right) \quad (1.43)$$

with the expectation value

$$\Delta_{i,\alpha} = -U_{\alpha} \langle c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \rangle \quad (1.44)$$

which is called the superconducting gap and is the order parameter introduced in Ginzburg-Landau theory in section 1.1. To include finite momentum in BCS theory, take the ansatz of a Fulde-Ferrel (FF) type pairing [33]:

$$\Delta_{i,\alpha} = \Delta_{\alpha} e^{i\mathbf{q}\mathbf{r}_{i\alpha}} \quad (1.45)$$

Using the Fourier transform

$$c_{i\alpha\sigma} = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}_{i\alpha}} c_{\mathbf{k}\alpha\sigma} \quad (1.46)$$

can write mean-field interaction term as

$$\sum_{i,\alpha} \left(\Delta_{i,\alpha} c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} + \Delta_{i,\alpha}^* c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} - \frac{|\Delta_{i,\alpha}|^2}{U_{\alpha}} \right) \quad (1.47)$$

can write

$$H_{\text{MF}} = \sum_{\mathbf{k}\alpha\beta\sigma} [H_{0,\sigma}(\mathbf{k})]_{\alpha\beta} c_{\mathbf{k}\alpha\sigma}^{\dagger} c_{\mathbf{k}\beta\sigma} + \sum_{\alpha,\mathbf{k}} (\Delta_{\alpha} c_{\mathbf{k}\alpha\uparrow}^{\dagger} c_{-\mathbf{k}\alpha\downarrow}^{\dagger} + \Delta_{\alpha}^* c_{-\mathbf{k}\alpha\downarrow} c_{\mathbf{k}\alpha\uparrow}) \quad (1.48)$$

The Hamiltonian in eq. (1.48) can be written as

How to include finite momentum, rewrite equations

there are phase factors introduced by the orbital positions

Get the remaining terms here

$$H_{\text{MF}} = \sum_{\mathbf{k}} \mathbf{C}_{\mathbf{k}}^\dagger H_{\text{BdG}}(\mathbf{k}) \mathbf{C}_{\mathbf{k}} \quad (1.49)$$

$$\mathbf{C}_{\mathbf{k}} = \left(c_{\mathbf{k}1\uparrow} \quad c_{\mathbf{k}2\uparrow} \quad \dots \quad c_{\mathbf{k}n_{\text{orb}}\uparrow} \quad c_{-\mathbf{k}1\downarrow}^\dagger \quad c_{-\mathbf{k}2\downarrow}^\dagger \quad \dots \quad c_{-\mathbf{k}n_{\text{orb}}\downarrow}^\dagger \right)^T \quad (1.50)$$

with the so-called Bogoliubov-de Gennes (BdG) matrix

$$H_{\text{BdG}}(\mathbf{k}) = \begin{pmatrix} H_{0,\uparrow}(\mathbf{k}) - \mu & \Delta \\ \Delta^\dagger & -H_{0,\downarrow}^*(-\mathbf{k}) + \mu \end{pmatrix} \quad (1.51)$$

with $H_{0,\sigma}$ being the F.T. of the kinetic term and $\Delta = \text{diag}(\Delta_1, \Delta_2, \dots, \Delta_{n_{\text{orb}}})$. Problem is now reduced to diagonalization of the BdG matrix. Write

$$H_{\text{BdG}} = U_{\mathbf{k}} \epsilon_{\mathbf{k}} U_{\mathbf{k}}^\dagger \quad (1.52)$$

and

$$H_{\text{MF}} = \sum_{\mathbf{k}} \gamma_{\mathbf{k}} \epsilon_{\mathbf{k}} \gamma_{\mathbf{k}}^\dagger \quad (1.53)$$

with quasi-particle operators

$$\gamma_{\mathbf{k}} = U_{\mathbf{k}}^\dagger c_{\mathbf{k}} \quad (1.54)$$

Using the gap equation

$$\Delta_\alpha = -U \quad (1.55)$$

the order parameter can be determined self-consistently, i.e. starting from an initial value, the BdG matrix needs to be set up, diagonalized and then used to determine Δ_α again, until a converged value is found.

1.3 Dynamical Mean-Field Theory

Green's Function Formalism

Green's functions: method to encode influence of many-body effects on propagation of particles in a system. Depending on the context, different kinds of Green's functions are employed. Following [3]. The Matsubara formalism naturally includes finite This is done via a so-called Wick rotation of the time variable t into imaginary time

$$t \rightarrow -i\tau \quad (1.56)$$

Write indices everywhere without comma

gap equation

SC current in BCS

Introduction DMFT, citing what has been achieved with it so far, what is the basic idea etc.

Give an introduction

Work over the paragraph

where τ is real and has the dimension time. This enables the simultaneous expansion of exponential $e^{-\beta H}$ coming from the thermodynamic average and e^{-iHt} coming from the time evolution of operators.

For our context (translationally invariant and systems), Matsubara GF are defined as

$$G(\mathbf{k}, \tau, 0) = -\langle T_\tau(A(\tau)B(0)) \rangle \quad (1.57)$$

with time-ordering operator in imaginary time:

$$T_\tau(A(\tau)B(\tau')) = \Theta(\tau - \tau')A(\tau)B(\tau') \pm \Theta(\tau' - \tau)B(\tau')A(\tau) \quad (1.58)$$

so that operators with later 'times' go to the left.

Can prove from properties of Matsubara GF, that they are only defined for

$$-\beta < \tau < \beta \quad (1.59)$$

Due to this, the Fourier transform of the Matsubara GF is defined on discrete values:

$$C_{AB}(i\omega_n) = \int_0^\beta d\tau \quad (1.60)$$

with fermionic/bosonic Matsubara frequencies

$$\omega_n = \begin{cases} \frac{2n\pi}{\beta} & \text{for bosons} \\ \frac{(2n+1)\pi}{\beta} & \text{for fermions} \end{cases} \quad (1.61)$$

It turns out that Matsubara GFs and retarded GFs can be generated from a common function $C_{AB}(z)$ that is defined on the entire complex plane except for the real axis. So I can get the retarded GF $C_{AB}^R(\omega)$ by analytic continuation:

$$C_{AB}^R(\omega) = C_{AB}(i\omega_n \rightarrow \omega + i\eta) \quad (1.62)$$

So in particular the extrapolation of the Matsubara GF to zero is proportional to the density of states at the chemical potential. Gapped: density is zero (Matsubara GF goes to 0), metal: density is finite (Matsubara GF goes to finite value).

What is the eta there -
> need to define it in retarded GF

Dyson Equation

Dyson equation:

$$G_\sigma(\mathbf{k}, i\omega_n) = \frac{G_\sigma^0(\mathbf{k}, i\omega_n)}{1 - G_\sigma^0(\mathbf{k}, i\omega_n)\Sigma_\sigma(\mathbf{k}, i\omega_n)} = \frac{1}{i\omega_n - \tilde{\epsilon}_{\mathbf{k}-\Sigma_\sigma(\mathbf{k}, i\omega_n)}} \quad (1.63)$$

Dyson equation

Self energy

DMFT

Following [34].

Most general non-interacting electronic Hamiltonian in second quantization:

$$H_0 = \sum_{i,j,\sigma} \quad (1.64)$$

with lattice coordinates i, j and spin σ .

One particle Green's function (many-body object, coming from the Hubbard model):

$$G(\mathbf{k}, i\omega_n) = \frac{1}{i\omega_n + \mu - \epsilon_{\mathbf{k}} - \Sigma(\mathbf{k}, i\omega_n)} \quad (1.65)$$

with the self energy $\Sigma(i\omega_n)$ coming from the solution of the effect on-site problem:

The Dyson equation

$$G(\mathbf{k}, i\omega_n) = (G_0(\mathbf{k}, i\omega_n) - \Sigma(\mathbf{k}, i\omega_n))^{-1} \quad (1.66)$$

relates the non-interacting Greens function $G_0(\mathbf{k}, i\omega_n)$ and the fully-interacting Greens function $G(\mathbf{k}, i\omega_n)$ (inversion of a matrix!).

Nambu-Gorkov Green's Functions

Introduction following [1, ch. 14.7]

Order parameter can be chosen as the anomalous GF:

$$\Psi = F^{\text{loc}}(\tau = 0^-) \quad (1.67)$$

or the superconducting gap

$$\Delta = Z\Sigma^{\text{AN}} \quad (1.68)$$

that can be calculated from the anomalous self-energy Σ^{AN} and quasiparticle weight Z

More general introduction into NG GFs, how they look like, what they describe etc.

Sources for these?

How to get quasiparticle weight?

See what is specific to this paper, see that I can derive that myself

1.4 Quantum Metric

First formulated in [35]

Following Cheng - a pedagogical Introduction

Parameter dependent Hamiltonian $\{H(\lambda)\}$, smooth dependence on parameter
 $\lambda = (\lambda_1, \lambda_2, \dots) \in \mathcal{M}$ (base manifold)
 Hamiltonian acts on parametrized Hilbert space $\mathcal{H}(\lambda)$
 Eigenenergies $E_n(\lambda)$, eigenstates $|\phi_n(\lambda)\rangle$
 System state $|\psi(\lambda)\rangle$ is linear combination of $|\phi_n(\lambda)\rangle$ at every point in \mathcal{M}
 Infinitesimal variation of the parameter $d\lambda$:

Dont get it
here

$$ds^2 = \|\psi(\lambda + d\lambda) - \psi(\lambda)\|^2 = \langle \delta\psi | \delta\psi \rangle = \langle \partial_\mu \psi | \partial_\nu \psi \rangle d\lambda^\mu d\lambda^\nu = (\gamma_{\mu\nu} + i\sigma_{\mu\nu}) d\lambda^\mu d\lambda^\nu \quad (1.69)$$

Last part is splitting up into real and imaginary part

Quantum Metric and Superfluid Weight

[8]

Write up
notes about
quantum
metric and
superfluid
weight