

# CONTENTS

I	INTRODUCTION	1
II	SUPERCONDUCTIVITY	3
II.1	Ginzburg-Landau Theory of Superconductivity . . . . .	3
II.2	Superconducting length scales . . . . .	8
II.3	Bardeen-Cooper-Schrieffer Theory . . . . .	9
II.3.1	BCS Hamiltonian . . . . .	9
II.3.2	Attractive Hubbard Model . . . . .	10
II.4	Dynamical Mean-Field Theory . . . . .	12
II.4.1	Green's Function Formalism . . . . .	12
II.4.2	Perturbation theory, Dyson equation . . . . .	14
II.4.3	Nambu-Gorkov GF . . . . .	15
II.4.4	DMFT . . . . .	15
III	QUANTUM METRIC	17
III.1	Quantum Metric and superfluid weight . . . . .	17
IV	DRESSED GRAPHENE MODEL	19
IV.1	Lattice Structure . . . . .	19
IV.2	EG-X Model . . . . .	22
V	RESULTS	25
V.1	Gaps . . . . .	25
V.2	Superfluid Weight . . . . .	25
V.3	Breakdown of SC with finite momentum . . . . .	25
V.4	Coherence length etc. . . . .	25
A	EG-X HAMILTONIAN IN RECIPROCAL SPACE	27
B	NOTES ON THE COMPUTATIONAL IMPLEMENTATION	31

ii CONTENTS

---

BIBLIOGRAPHY	33
NOT CITED	39
LIST OF SYMBOLS	41

# TODO LIST

What to put in introduction? . . . . .	1
More extensive introduction . . . . .	3
Introduce spontaneous symmetry breaking . . . . .	3
Work over paragraph . . . . .	3
Work over paragraph . . . . .	4
Make graphic for Landau free energy . . . . .	4
Make graphic for Landau OP and BCS OP . . . . .	4
Make my own graphic for mexican hat potential . . . . .	4
Work over paragraph . . . . .	5
Work over paragraph . . . . .	6
I dont know why that is. Can I support that somehow better? -> See Niklas thesis . . . . .	7
Phase rigidity and superflow . . . . .	8
Better introduction . . . . .	8
Depairing current from FMP . . . . .	9
Full formula for supercurrent, with sum over orbitals . . . . .	9
DS from FMP . . . . .	9
Write more about the connection between all the things here . . . . .	9
Microscopic OP in BCS . . . . .	10
What is explained by phononic pairing? . . . . .	10
Other pairing interactions can be taken, gives explanations for a lot of different SCs . . . . .	10
Introduce mean field here . . . . .	10
What is the connection between mean field theory and BCS theory? . . .	10
Some relevance of the repulsive Hubbard model . . . . .	10
There are some more specific papers to the specific mechanisms (and also some more mechanism), could cite these here and say some more things . . . . .	11
Define: what exactly are orbitals in this context? . . . . .	11
General multi-band mean field theory theory . . . . .	12

How to solve mean field theory self-consistently . . . . .	12
How to include finite momentum . . . . .	12
Show GFs can be related to observables . . . . .	13
How to resolve ambiguity at borders of integral . . . . .	14
single-particle Matsubara GF . . . . .	14
equations of motion for Matsubara GF . . . . .	14
Short introduction to diagrams . . . . .	14
Self energy . . . . .	14
Dyson equation . . . . .	14
More general introduction into NG GFs, how they look like, what they describe etc. . . . .	15
Sources for these? . . . . .	15
How to get quasiparticle weight? . . . . .	15
See what is specific to this paper, see that I can derive that myself . . . .	17
Dont get it here . . . . .	17
Write up notes about quantum metric and superfluid weight . . . . .	17
Write introduction to the model and what is done in this chapter . . . . .	19
First BZ vs this Gamma centered one . . . . .	20
Write the Hamiltonian with orbital indices . . . . .	23
Clean up this section . . . . .	27
Correct exp expressions . . . . .	28
Example for a vector product . . . . .	28
Show that! . . . . .	28
Data availability . . . . .	31
What software for what? . . . . .	31

# I INTRODUCTION

In 1894, Albert Michelson remarked that “it seems probable that most of the grand underlying principles have been firmly established”<sup>1</sup> [1, p. 159].

What to put in introduction?

At the end of the 19th and the beginning of the 20th century, cooling technology made great progress. Liquifying gases, were able to reach temperatures as low as 4 K (the boiling point of Helium). Using that, SC was discovered in mercury in 1911 by Heike Onnes [2]. Superconductivity describes the phenomenon of the electrical resistance of a material suddenly dropping to zero below a critical temperature  $T_C$ . Discovery of Meissner effect, perfect expulsion of external magnetic fields in 1933 [3]. This started almost half a century of intensive theoretical research, which culminated in John Bardeen, Leon Cooper and J. Robert Schrieffer developing the microscopic theory now known as BCS theory [4]. 1986 and 1987: discovery of superconductivity with very high  $T_C$  found in cuprates [5, 6]. Cuprate superconductors are made up of layers of copper oxide and charge reservoirs in between. The specific charge reservoir layers determine the properties of the SC and varying them lead to a rich zoo of materials with high  $T_C$  [7].

Largest commercial application to date is in magnetic resonance imaging, a medical technique using strong magnetic fields and field gradients [8]. Enabled due to the fact, that SCs can carry much stronger currents and thus generate much higher magnetic field strength. Technical applications in research are much wider, ranging from strong superconducting magnets in the LHC [9] and other particle accelerators over detectors of single photons in astrophysics [10] to extremely sensitive measurement devices for magnetic fields [11] and voltages [12] based on the Josephson effect [13].

Since the first discovery of SC in cuprates, there has been a lot of work to develop superconductors with higher transition temperatures. One interesting development is in twisted multilayer systems, first realized as twisted bilayer

---

<sup>1</sup>Variations of this quote have been attributed to Lord Kelvin, although the poetry having it said by someone, whose experiment would eventually lead to the development of special relativity is undeniable.

Graphene [14]. In comparison to the complex crystal structure of e.g. the Cuprates, twisted multilayer systems have a very simple structure and can be tuned very easily: the angle of twist between the layers can be easily accessed experimentally. The defining feature of these systems are flat electronic bands due to folding of the Brilluoin zone. Superconductivity in these systems is enhanced due to the fact that in the flat bands, interactions between the electrons are very strongly enhanced. Thus these systems are a very interesting playground to study strongly correlation effects in general and superconductivity in particular.

## II SUPERCONDUCTIVITY

In this chapter: review theoretical concepts needed for describing SC.

Macroscopically, SC state can be described by a spontaneous breaking of a  $U(1)$  phase rotation symmetry, that is associated with an order parameter. Theory of this: GL theory section II.1.

One tool to describe superconductivity from a microscopic perspective: BCS theory section II.3.

Taking fluctuations beyond mean field into account: DMFT section II.4.

There are many textbooks covering these topics which can be referenced for a more detailed treatment, such as refs. [15–18].

### II.1 GINZBURG-LANDAU THEORY OF SUPERCONDUCTIVITY

For this review, follow chapter 11 in ref. [15].

#### ORDER PARAMETER [19]

Landau theory: phase transitions (e.g. iron becomes magnetic, water freezes, superfluidity/superconductivity) are associated with the development of an order parameter when the temperature drops below the transition temperature  $T_C$ .

$$|\psi| = \begin{cases} 0, & T > T_C \\ |\psi_0| > 0, & T < T_C \end{cases} \quad (\text{II.1})$$

Landau theory does not need microscopic expression for order parameter, it provides coarse-grained description of the properties of matter. The order parameter description is good at length scales above  $\xi_0$ , the coherence length (e.g. size of Cooper pairs for SC). On length scales above  $\xi_0$ , the order parameter behaves as a smoothly varying function.

More extensive introduction

Introduce spontaneous symmetry breaking

Work over paragraph

## LANDAU THEORY

Work over p

Basic idea of Landau theory: write free energy as function  $F[\psi]$  of the order parameter. Region of small  $\psi$ , expand free energy of many-body system as simple polynomial:

$$f_L = \frac{1}{V}F[\psi] = \frac{r}{2}\psi^2 + \frac{u}{4}\psi^4 \quad (\text{II.2})$$

Provided  $r$  and  $u$  are greater than 0: minimum of  $f_L[\psi]$  lies at  $\psi = 0$ . Landau theory assumes: at phase transition temperature  $r$  changes sign, so:

$$r = a(T - T_C) \quad (\text{II.3})$$

Minimum of free energy occurs for:

$$\psi = \begin{cases} 0 \\ \pm \sqrt{\frac{a(T_C - T)}{u}} \end{cases} \quad (\text{II.4})$$

Make graphic for  
Landau free energy

Make graphic for  
Landau OP and BCS  
OP

Two minima for free energy function for  $T < T_C$ . With this, we can extract  $T_C$  from the knowledge of the dependence of  $|\psi|^2$  on  $T$  via a linear fit. This is only valid for an area near  $T_C$  (where Landau theory holds), but can be used to get  $T_C$  from microscopic theories.

Going from a one to a  $n$ -component order parameters, OP acquires directions and magnitude. Particularly important example: complex or two component order parameter in superfluids and superconductors:

$$\psi = \psi_1 + i\psi_2 = |\psi|e^{i\phi} \quad (\text{II.5})$$

The Landau free energy takes the form:

$$f[\psi] = r(\psi^*\psi) + \frac{u}{2}(\psi^*\psi)^2 \quad (\text{II.6})$$

As before:

$$r = a(T - T_C) \quad (\text{II.7})$$


Make my own  
graphic for mexi-  
can hat potential

Figure II.1 shows the Landau free energy as function of  $\psi$ .

Rotational symmetry, because free energy is independent of the global phase of the OP:

$$f[\psi] = f[e^{ia}\psi] \quad (\text{II.8})$$





**Figure II.1:** Mexican hat potential

In this 'Mexican hat' potential: order parameter can be rotated continuously from one broken-symmetry state to another. If we want the phase to be rigid, we need to introduce an There is a topological argument for the fact that the phase is rigid. This leads to Ginzburg-Landau theory. Will see later: well-defined phase is associated with persistent currents or superflow.

#### GINZBURG-LANDAU THEORY

Work over paragraph

Landau theory: energy cost of a uniform order parameter, more general theory needs to account for inhomogenous order parameters, in which the amplitude varies or direction of order parameter is twisted -> GL theory. First: one-component, 'Ising' order parameter. GL introduces additional energy  $\delta f \propto |\Delta\psi|^2$ ,  $f_{GL}[\psi, \Delta\psi] = \frac{s}{2}|\Delta\psi|^2 + f_L[\psi(s)]$ , or in full:

$$f_{GL}[\psi, \Delta\psi, h] = \frac{s}{2}(\Delta\psi)^2 + \frac{r}{2}\psi^2 + \frac{u}{4}\psi^4 \quad (\text{II.9})$$

GL theory is only valid near critical point, where OP is small enough to permit leading-order expansion. Dimensional analysis shows:  $\frac{s}{r} = L^2$  has dimension of length squared. Length scale introduced by the gradient term: correlation

length

$$\xi(T) = \sqrt{\frac{s}{|r(T)|}} = \xi_0 \left| 1 - \frac{T}{T_C} \right|^{-\frac{1}{2}} \quad (\text{II.10})$$

sets characteristic length scale of order-parameter fluctuations, where

$$\xi_0 = \xi(T = 0) = \sqrt{\frac{s}{\alpha T_C}} \quad (\text{II.11})$$

is a measure of the microscopic coherence length. Near transition  $\xi(T)$  diverges, but far from transition it becomes comparable with the coherence length.

Work over paragraph

#### COMPLEX ORDER AND SUPERFLOW

Now: GL theory of complex or two-component order parameters, so superfluids and superconductors. Heart of discussion: emergence of a ‘macroscopic wavefunction’, where the microscopic field operators  $\hat{\psi}(x)$  acquire an expectation value:

$$\langle \hat{\psi}(x) \rangle = \psi(x) = |\psi(x)| e^{i\theta(x)} \quad (\text{II.12})$$

Reminder: Field operators are the real space representations of creation/annihilation operators. They can be thought of the super position of all ways of creating a particle at position  $x$  via the basis coefficients.

Magnitude determines density of particles in the superfluid:

$$|\psi(x)|^2 = n_s(x) \quad (\text{II.13})$$

Density operator is

$$\hat{\rho} = \hat{\psi}(x) \hat{\psi}^\dagger(x) \quad (\text{II.14})$$

so expectation value of that is the formula above.

Twist/gradient of phase determines superfluid velocity:

$$\mathbf{v}_s(x) = \frac{\hbar}{m} \Delta \phi(x) \quad (\text{II.15})$$

We will derive this later in the chapter. Counterintuitive from quantum mechanics: GL suggested that  $\Phi(x)$  is a macroscopic manifestation of a macroscopic number of particles condensed into precisely the same quantum state. Emergent phenomenon, collective properties of matter not a-priori evident from microscopic physics.

GL free energy density for superfluid (with one added term in comparison to Landau energy):

$$f_{GL}[\psi, \Delta\psi] = s|\Delta\psi|^2 + r|\psi|^2 + \frac{u}{2}|\psi|^4 \quad (\text{II.16})$$

Compare with the energy density of a bosonic field (with a quartic interaction):

$$H = \int d^Dx \frac{\hbar^2}{2m} |\Delta\psi|^2 + r|\psi|^2 + \frac{u}{2}|\psi|^4 \quad (\text{II.17})$$

Interpret GL free energy as energy density of a condensate of bosons in which the field operator behaves as a complex order parameter. Gives interpretation of gradient term as kinetic energy:

$$s|\Delta\psi|^2 = \frac{\hbar^2}{2m} \langle \Delta\hat{\psi}^\dagger \Delta\hat{\psi} \rangle \implies s = \frac{\hbar^2}{2m} \quad (\text{II.18})$$

As in Ising order: correlation length/GL-coherence length governs characteristic range of amplitude fluctuations of the order parameter:

$$\xi = \sqrt{\frac{s}{|r|}} = \sqrt{\frac{\hbar^2}{2m|r|}} = \xi_0 \left(1 - \frac{T}{T_C}\right)^{-\frac{1}{2}} \quad (\text{II.19})$$

where  $\xi_0 = \xi(T=0) = \sqrt{\frac{\hbar^2}{2maT_C}}$  is the coherence length. Beyond this length scale: only phase fluctuations survive.

Freeze out fluctuations in amplitude (no  $x$ -dependence in amplitude)  $\psi(x) = \sqrt{n_s} e^{i\phi(x)}$ , then  $\Delta\psi = i\Delta\phi\psi$  and  $|\Delta\psi|^2 = n_s(\Delta\phi)^2$ , dependency of kinetic energy on the phase twist is (bringing it into the form  $\frac{m}{2}v^2$ ):

$$\frac{\hbar^2 n_s}{2m} (\Delta\phi)^2 = \frac{mn_s}{2} \left(\frac{\hbar}{m} \Delta\phi\right)^2 \quad (\text{II.20})$$

So twist of phase results in increase in kinetic energy, associated with a superfluid velocity:

$$\mathbf{v}_s = \frac{\hbar}{m} \Delta\phi \quad (\text{II.21})$$

(this is explained in detail later).

I dont know why that is. Can I support that somehow better? -> See Niklas thesis

For interpretation of superfluid states: coherent states. These are eigenstates of the field operator

$$\hat{\psi}(x) |\psi\rangle = \psi(x) |\psi\rangle \quad (\text{II.22})$$

and don't have a definite particle number. Importantly, this small uncertainty in particle number enables a high degree of precision in phase (which is the property of a condensate).

**PHASE RIGIDITY AND SUPERFLOW** In GL theory, energy is sensitive to a twist of the phase. Substitute  $\psi = |\psi|e^{i\phi}$  into GL free energy, gradient term is:

$$\Delta\psi = (\Delta|\psi| + i\Delta\phi|\psi|)e^{i\phi} \quad (\text{II.23})$$

So:

$$f_{GL} = \frac{\hbar}{2m}|\psi|^2(\Delta\phi)^2 + \left[ \frac{\hbar}{2m}(\Delta|\psi|)^2 + r|\psi|^2 + \frac{u}{2}|\psi|^4 \right] \quad (\text{II.24})$$

The second term resembles GL functional for an Ising order parameter, describes energy cost of variations in the magnitude of the order parameter.

Phase rigidity and superflow

## II.2 SUPERCONDUCTING LENGTH SCALES

Better introduction

From [20].

In most materials: Cooper pairs do not carry finite center-of-mass momentum. In presence of e.g. external fields or magnetism: SC states with FMP might arise.

Theory/procedure in the paper: enforce FMP states via constraints on pair-center-of-mass momentum  $\mathbf{q}$ , access characteristic length scales  $\xi_0, \lambda_L$  through analysis of the momentum and temperature-dependent OP. FF-type pairing with Cooper pairs carrying finite momentum:

$$\psi_{\mathbf{q}}(\mathbf{r}) = |\psi_{\mathbf{q}}|e^{i\mathbf{q}\cdot\mathbf{r}} \quad (\text{II.25})$$

Then the free energy density is

$$f_{GL}[\psi_{\mathbf{q}}] = \alpha|\psi_{\mathbf{q}}|^2 + \frac{b}{2}|\psi_{\mathbf{q}}|^4 + \frac{\hbar^2 q^2}{2m^*}|\psi_{\mathbf{q}}|^2 \quad (\text{II.26})$$

Stationary point of the system:

$$\frac{\delta f_{GL}}{\delta \psi_{\mathbf{q}}^*} = 2\psi_{\mathbf{q}} [\alpha(1 - \xi^2 q^2) + b|\psi_{\mathbf{q}}|^2] = 0 \quad (\text{II.27})$$

which results in the  $\mathbf{q}$ -dependence of the OP

$$|\psi_{\mathbf{q}}|^2 = |\psi_0|^2(1 - \xi(T)^2 q^2) \quad (\text{II.28})$$

For some value, SC order breaks down,  $\psi_{\mathbf{q}_c} = 0$ , because the kinetic energy from phase modulation exceeds the gain in energy from pairing. In GL theory:  $q_c = \xi(T)^{-1}$ . The temperature dependence of the OP and extracted  $\xi(T)$  gives access to the coherence length via

$$\xi(T) = \xi_0(1 - \frac{T}{T_C})^{-\frac{1}{2}} \quad (\text{II.29})$$

Specifically: take

$$\xi(T) = \frac{1}{\sqrt{2}|\mathbf{Q}|} \quad (\text{II.30})$$

with  $\mathbf{Q}$  such that

$$|\frac{\psi_{\mathbf{Q}}(T)}{\psi_0(T)}| = \frac{1}{\sqrt{2}} \quad (\text{II.31})$$

The Cooper pair [21, 22]

## II.3 BARDEEN-COOOPER-SCHRIEFFER THEORY

### II.3.1 BCS HAMILTONIAN

Following [15, ch. 14].

First phenomenological description of SC: Fritz London in 1937 [23]. He was motivated by the discovery of the Meissner effect in 1933 [3], where magnetic flux inside of the superconductor is always pushed out in contrast to a perfectly conducting material, which would hold a ‘memory’ of the magnetic field at the time of the phase transition. This suggests that transition to the SC state is reversible and a SC is not just the limiting case of a conductor with infinite conductivity, in which according to the Maxwell equations, the magnetic flux

Depairing current from FMP

Full formula for supercurrent, with sum over orbitals

DS from FMP

Write more about the connection between all the things here

would not change. Londons first descriptions is based on a one-particle wave function  $\phi(x)$ . He proposed that persistent supercurrent is a property of the ground state associated with its rigidity against the application of a field.

In 1950 [19]: GL interpreted this wave function as a complex order parameter as explained in section II.1.

Microscopic description of SC: 1957 by John Bardeen, his postdoc Leon Cooper and the graduate in the group, J. Robert Schrieffer [4]. Description is based on the fact that the Fermi sea is unstable towards development of bound pairs under arbitrarily small attraction [24].

The BCS-Hamiltonian is:

$$H_{\text{BCS}} = \sum_{\mathbf{k}\sigma} \epsilon_{\mathbf{k}\sigma} c_{\mathbf{k}\sigma}^\dagger c_{\mathbf{k}\sigma} + \sum_{\mathbf{k}, \mathbf{k}'} V_{\mathbf{k}, \mathbf{k}'} c_{\mathbf{k}\uparrow}^\dagger c_{-\mathbf{k}\downarrow}^\dagger c_{-\mathbf{k}'\downarrow} c_{\mathbf{k}'\uparrow} \quad (\text{II.32})$$

The final element in this description was the origin of the attractive interaction  $V_{\mathbf{k}, \mathbf{k}'}$  between electrons, which Bardeen, Cooper and Schrieffer identified as a retarded electron-phonon interaction [4]. This so-called BCS-theory of superconductivity is very successful in explaining experimental results in many compounds.

BCS theory always comes in conjunction with a mean-field description, decoupling the

The reason for this: the interaction in the BCS Hamiltonian eq. (II.32) takes place at exactly zero momentum and as such is an infinite range interaction

A

### II.3.2 ATTRACTIVE HUBBARD MODEL

Simple choice for pairing interaction:

$$V_{\mathbf{k}, \mathbf{k}'} = U \quad (\text{II.33})$$

This is then just the (attractive) Hubbard model. The Hubbard model is the simplest model for interacting electron systems. It goes back to works by Hubbard [25], Kanamori [26] and Gutzwiller [27].

$$H_{\text{int}} = U \sum_i c_{i,\uparrow}^\dagger c_{i,\downarrow}^\dagger c_{i,\downarrow} c_{i,\uparrow} \quad (\text{II.34})$$

where  $U > 0$ .

Microscopic OP in BCS

What is explained by phononic pairing?

Other pairing interactions can be taken, gives explanations for a lot of different SCs

Introduce mean field here

What is the connection between mean field theory and BCS theory?

Some relevance of the repulsive Hubbard model

Besides

[28]

This simple Hubbard model can be extended in a multitude of ways to model a variety of physical system. In this work: extension to multiple orbitals and an attractive interaction, i.e. a negative  $U$ . Physical motivation for taking a negative- $U$  Hubbard model: electrons can experience a local attraction interaction, for example through electrons coupling with phononic degrees of freedom or with electronic excitations that can be described as bosons [29]. The form of the interaction term is then:

$$H_{\text{int}} = - \sum_{i,\alpha} U_{\alpha} c_{i,\alpha,\uparrow}^{\dagger} c_{i,\alpha,\downarrow}^{\dagger} c_{i,\alpha,\downarrow} c_{i,\alpha,\uparrow} \quad (\text{II.35})$$

where  $\alpha$  counts orbitals and the minus sign in front is taken so that  $U > 0$  now corresponds to an attractive interaction (this is purely convention).

**MULTI-BAND BCS THEORY** Look at interaction term eq. (II.35). Mean-field approximation: operators do not deviate much from their average value:

$$H_{\text{int}} \approx \sum_{\alpha,\mathbf{k}} (\Delta_{\alpha} c_{\mathbf{k}\alpha\uparrow}^{\dagger} c_{-\mathbf{k}\alpha\downarrow}^{\dagger} + \Delta_{\alpha}^{*} c_{-\mathbf{k}\alpha\downarrow} c_{\mathbf{k}\alpha\uparrow}) \quad (\text{II.36})$$

Fourier transformation:

$$H_{\text{int}} = - \frac{1}{N^2} \sum_{\alpha,\mathbf{k}_{1,2,3,4}} U_{\alpha} e^{i(\mathbf{k}_1 + \mathbf{k}_4 - \mathbf{k}_1 - \mathbf{k}_3) \cdot \mathbf{r}_{i\alpha}} c_{\mathbf{k}_1\alpha\uparrow}^{\dagger} c_{\mathbf{k}_3\alpha\downarrow}^{\dagger} c_{\mathbf{k}_2\alpha\downarrow} c_{\mathbf{k}_4\alpha\uparrow} \quad (\text{II.37})$$

Impose zero-momentum pairing:  $\mathbf{k}_1 + \mathbf{k}_3 = 0$  and  $\mathbf{k}_2 + \mathbf{k}_4 = 0$ :

$$H_{\text{int}} = - \sum_{\alpha,\mathbf{k},\mathbf{k}'} U_{\alpha} c_{\mathbf{k}\alpha\uparrow}^{\dagger} c_{-\mathbf{k}\alpha\downarrow}^{\dagger} c_{-\mathbf{k}'\alpha\downarrow} c_{\mathbf{k}'\alpha\uparrow} \quad (\text{II.38})$$

Mean-field approximation:

$$H_{\text{int}} \approx \sum_{\alpha,\mathbf{k}} (\Delta_{\alpha} c_{\mathbf{k}\alpha\uparrow}^{\dagger} c_{-\mathbf{k}\alpha\downarrow}^{\dagger} + \Delta_{\alpha}^{*} c_{-\mathbf{k}\alpha\downarrow} c_{\mathbf{k}\alpha\uparrow}) \quad (\text{II.39})$$

with

$$\Delta_{\alpha} = -U_{\alpha} \sum_{\mathbf{k}'} \langle c_{-\mathbf{k}'\alpha\downarrow} c_{\mathbf{k}'\alpha\uparrow} \rangle \quad (\text{II.40})$$

$$\Delta_{\alpha}^{*} = -U_{\alpha} \sum_{\mathbf{k}'} \langle c_{\mathbf{k}'\alpha\uparrow}^{\dagger} c_{-\mathbf{k}'\alpha\downarrow}^{\dagger} \rangle \quad (\text{II.41})$$

There are some more specific papers to the specific mechanisms (and also some more mechanism), could cite these here and say some more things

Define: what exactly are orbitals in this context?

This gives the BCS mean field Hamiltonian:

$$H_{BCS} = \sum_{\mathbf{k}\alpha\beta\sigma} [H_{0,\sigma}(\mathbf{k})]_{\alpha\beta} c_{\mathbf{k}\alpha\sigma}^\dagger c_{\mathbf{k}\beta\sigma} - \mu \sum_{\mathbf{k}\alpha\sigma} n_{\mathbf{k}\alpha\sigma} + \sum_{\alpha,\mathbf{k}} (\Delta_\alpha c_{\mathbf{k}\alpha\uparrow}^\dagger c_{-\mathbf{k}\alpha\downarrow}^\dagger + \Delta_\alpha^* c_{-\mathbf{k}\alpha\downarrow} c_{\mathbf{k}\alpha\uparrow}) \quad (\text{II.42})$$

with Nambu spinor

$$\Psi_{\mathbf{k}} = \begin{pmatrix} c_{1,\mathbf{k}\uparrow} \\ c_{2,\mathbf{k}\uparrow} \\ c_{3,\mathbf{k}\uparrow} \\ c_{1,-\mathbf{k}\downarrow}^\dagger \\ c_{2,-\mathbf{k}\downarrow}^\dagger \\ c_{3,-\mathbf{k}\downarrow}^\dagger \end{pmatrix} \quad (\text{II.43})$$

we have:

$$H_{MF} = \sum_{\mathbf{k}} \Psi_{\mathbf{k}}^\dagger \mathcal{H}(\mathbf{k}) \Psi_{\mathbf{k}} \quad (\text{II.44})$$

with

$$\mathcal{H}(\mathbf{k}) = \begin{pmatrix} H_{0,\uparrow}(\mathbf{k}) - \mu & \Delta \\ \Delta^\dagger & -H_{0,\downarrow}^*(-\mathbf{k}) + \mu \end{pmatrix} \quad (\text{II.45})$$

General multi-band  
mean field theory  
theory

with  $H_{0,\sigma}$  being the F.T. of the kinetic term and  $\Delta = \text{diag}(\Delta_1, \Delta_2, \Delta_3)$ .

#### SELF-CONSISTENT SOLUTION

How to solve mean  
field theory self-  
consistently

**FINITE MOMENTUM** To include finite momentum, take the ansatz of a Fulde-Ferrel (FF) type pairing [30]:

$$\Delta \quad (\text{II.46})$$

How to include fi-  
nite momentum

## II.4 DYNAMICAL MEAN-FIELD THEORY

### II.4.1 GREEN'S FUNCTION FORMALISM

Following [17]

Green's functions: method to encode influence of many-body effects on propagation of particles in a system.



Have different kinds of Green's functions, for example the retarded Green's function:

$$G^R(\mathbf{r}\sigma t, \mathbf{r}'\sigma' t') = -i\Theta(t - t') \langle \{c_{\mathbf{r}\sigma}(t), c_{\mathbf{r}'\sigma'}^\dagger(t')\} \rangle \quad (\text{II.47})$$

They give the amplitude of a particle inserted at point  $\mathbf{r}'$  at time  $t'$  to propagate to position  $\mathbf{r}$  at time  $t$ . For time-independent Hamiltonians and systems in equilibrium, the GFs only depend on time differences:

$$G^R(\mathbf{r}\sigma t, \mathbf{r}'\sigma' t') = G^R(\mathbf{r}\sigma, \mathbf{r}'\sigma', t - t') \quad (\text{II.48})$$

So we can take  $t' = 0$  and consider  $t$  as the only free variable:

$$G^R(\mathbf{r}\sigma, \mathbf{r}'\sigma', t) = -i\Theta(t) \langle \{c_{\mathbf{r}\sigma}(t), c_{\mathbf{r}'\sigma'}^\dagger(0)\} \rangle \quad (\text{II.49})$$

In a translation invariant system: can use  $\mathbf{k}$  as a natural basis set:

$$G^R(\mathbf{k}, \sigma, \sigma', t) = -i\Theta(t) \langle \{c_{\mathbf{k}\sigma}(t), c_{\mathbf{k}\sigma'}^\dagger(0)\} \rangle \quad (\text{II.50})$$

Define Fourier-transform:

$$G^R(\mathbf{k}, \sigma, \sigma', \omega) = \int_{-\infty}^{\infty} dt G^R(\mathbf{k}, \sigma, \sigma', t) \quad (\text{II.51})$$

Can define the spectral function from this:

$$A(\mathbf{k}\sigma, \omega) = -2\Im G^R(\mathbf{k}\sigma, \omega) \quad (\text{II.52})$$

Looking at the diagonal elements of  $G^R$  here. The spectral function can be thought of as the energy resolution of a particle with energy  $\omega$ . This mean, for non-interacting systems, the spectral function is a delta-function around the single-particle energies:

$$A_0(\mathbf{k}\sigma, \omega) = 2\pi\delta(\omega - \epsilon_{\mathbf{k}\sigma}) \quad (\text{II.53})$$

For interacting systems this is not true, but  $A$  can still be peaked.

Mathematical technique to calculate retarded GFs involves defining GFs on imaginary times  $\tau$ :

$$t \rightarrow -i\tau \quad (\text{II.54})$$

where  $\tau$  is real and has the dimension time. This enables the simultaneous expansion of exponential  $e^{-\beta H}$  coming from the thermodynamic average and  $e^{-iHt}$  coming from the time evolution of operators.

Show GFs can be related to observables

Define imaginary time/Matsubara GF  $C_{AB}(\tau, 0)$ :

$$C_{AB}(\tau, 0) = -\langle T_\tau(A(\tau)B(0)) \rangle \quad (\text{II.55})$$

with time-ordering operator in imaginary time:

$$T_\tau(A(\tau)B(\tau')) = \Theta(\tau - \tau')A(\tau)B(\tau') \pm \Theta(\tau' - \tau)B(\tau')A(\tau) \quad (\text{II.56})$$

so that operators with later 'times' go to the left.

Can prove from properties of Matsubara GF, that they are only defined for

$$-\beta < \tau < \beta \quad (\text{II.57})$$

Due to this, the Fourier transform of the Matsubara GF is defined on discrete values:

$$C_{AB}(i\omega_n) = \int_0^\beta d\tau \quad (\text{II.58})$$

with fermionic/bosonic Matsubara frequencies

$$\omega_n = \begin{cases} \frac{2n\pi}{\beta} & \text{for bosons} \\ \frac{(2n+1)\pi}{\beta} & \text{for fermions} \end{cases} \quad (\text{II.59})$$

How to resolve ambiguity at borders of integral

It turns out that Matsubara GFs and retarded GFs can be generated from a common function  $C_{AB}(z)$  that is defined on the entire complex plane except for the real axis. So we can get the retarded GF  $C_{AB}^R(\omega)$  by analytic continuation:

$$C_{AB}^R(\omega) = C_{AB}(i\omega_n \rightarrow \omega + i\eta) \quad (\text{II.60})$$

So in particular the extrapolation of the Matsubara GF to zero is proportional to the density of states at the chemical potential. Gapped: density is zero (Matsubara GF goes to 0), metal: density is finite (Matsubara GF goes to finite value) [17, p. 8.3.4].

single-particle Matsubara GF

equations of motion for Matsubara GF

Short introduction to diagrams

Self energy

Dyson equation

## II.4.2 PERTURBATION THEORY, DYSON EQUATION

Dyson equation:

$$G_\sigma(\mathbf{k}, i\omega_n) = \frac{G_\sigma^0(\mathbf{k}, i\omega_n)}{1 - G_\sigma^0(\mathbf{k}, i\omega_n)\Sigma_\sigma(\mathbf{k}, i\omega_n)} = \frac{1}{i\omega_n - \tilde{\epsilon}_{\mathbf{k}-\Sigma_\sigma(\mathbf{k}, i\omega_n)}} \quad (\text{II.61})$$

## II.4.3 NAMBU-GORKOV GF

Introduction following [15, ch. 14.7]

Order parameter can be chosen as the anomalous GF:

$$\Psi = F^{\text{loc}}(\tau = 0^-) \quad (\text{II.62})$$

or the superconducting gap

$$\Delta = Z\Sigma^{\text{AN}} \quad (\text{II.63})$$

that can be calculated from the anomalous self-energy  $\Sigma^{\text{AN}}$  and quasiparticle weight  $Z$

Sources for these?

## II.4.4 DMFT

Following [31].

Most general non-interacting electronic Hamiltonian in second quantization:

$$H_0 = \sum_{i,j,\sigma} \quad (\text{II.64})$$

with lattice coordinates  $i, j$  and spin  $\sigma$ .

One particle Green's function (many-body object, coming from the Hubbard model):

$$G(\mathbf{k}, i\omega_n) = \frac{1}{i\omega_n + \mu - \epsilon_{\mathbf{k}} - \Sigma(\mathbf{k}, i\omega_n)} \quad (\text{II.65})$$

with the self energy  $\Sigma(i\omega_n)$  coming from the solution of the effect on-site problem:

The Dyson equation

$$G(\mathbf{k}, i\omega_n) = (G_0(\mathbf{k}, i\omega_n) - \Sigma(\mathbf{k}, i\omega_n))^{-1} \quad (\text{II.66})$$

relates the non-interacting Greens function  $G_0(\mathbf{k}, i\omega_n)$  and the fully-interacting Greens function  $G(\mathbf{k}, i\omega_n)$  (inversion of a matrix!).

How to get quasiparticle weight?

General intro-  
to NG GFs,  
look like,  
describe



### III QUANTUM METRIC

First formulated in [32]

Following Cheng - a pedagogical Introduction

Parameter dependent Hamiltonian  $\{H(\lambda)\}$ , smooth dependence on parameter

$\lambda = (\lambda_1, \lambda_2, \dots) \in \mathcal{M}$  (base manifold)

Hamiltonian acts on parametrized Hilbert space  $\mathcal{H}(\lambda)$

Eigenenergies  $E_n(\lambda)$ , eigenstates  $|\phi_n(\lambda)\rangle$

System state  $|\psi(\lambda)\rangle$  is linear combination of  $|\psi_n(\lambda)\rangle$  at every point in  $\mathcal{M}$

Infinitesimal variation of the parameter  $d\lambda$  :

See what is specific to this paper, see that I can derive that myself

Dont get it here

$$ds^2 = \|\psi(\lambda+d\lambda) - \psi(\lambda)\|^2 = \langle \delta\psi | \delta\psi \rangle = \langle \partial_\mu \psi | \partial_\nu \psi \rangle d\lambda^\mu d\lambda^\nu = (\gamma_{\mu\nu} + i\sigma_{\mu\nu}) d\lambda^\mu d\lambda^\nu \quad (\text{III.1})$$

Last part is splitting up into real and imaginary part

#### III.1 QUANTUM METRIC AND SUPERFLUID WEIGHT

Write up notes about quantum metric and superfluid weight




## IV DRESSED GRAPHENE MODEL

Write introduction to the model and what is done in this chapter

### IV.1 LATTICE STRUCTURE

Structure of honeycomb lattice following [33].

Monolayer graphene forms a honeycomb lattice, which is a hexagonal Bravais lattice with a two atom basis.



**Figure IV.1:** *Graphene lattice structure*

Primitive lattice vectors of the hexagonal lattice:

$$\mathbf{a}_1 = \frac{a}{2} \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} \quad (\text{IV.1})$$

$$\mathbf{a}_2 = \frac{a}{2} \begin{pmatrix} 1 \\ -\sqrt{3} \end{pmatrix} \quad (\text{IV.2})$$

with lattice constant  $a \approx 2.46 \text{ \AA}$  (distance between unit cells). Have

$$a = \sqrt{3}a_0 \quad (\text{IV.3})$$

with the nearest-neighbour distance  $a_0$ .

Vectors to the nearest-neighbor  $B_i$  ( $i = 1, 2, 3$ ) atoms from atom  $A$ :

$$\delta_{AB,1} = \begin{pmatrix} 0 \\ \frac{a}{\sqrt{3}} \end{pmatrix}, \delta_{AB,2} = \begin{pmatrix} \frac{a}{2} \\ -\frac{a}{2\sqrt{3}} \end{pmatrix}, \delta_{AB,3} = \begin{pmatrix} -\frac{a}{2} \\ -\frac{a}{2\sqrt{3}} \end{pmatrix} \quad (\text{IV.4})$$

Vectors to the nearest-neighbor  $A_i$  ( $i = 1, 2, 3$ ) atoms from atom  $B$ :

$$\delta_{BA,1} = \begin{pmatrix} 0 \\ -\frac{a}{\sqrt{3}} \end{pmatrix}, \delta_{BA,2} = \begin{pmatrix} \frac{a}{2} \\ \frac{a}{2\sqrt{3}} \end{pmatrix}, \delta_{BA,3} = \begin{pmatrix} -\frac{a}{2} \\ \frac{a}{2\sqrt{3}} \end{pmatrix} \quad (\text{IV.5})$$

The vectors between the Graphene  $A$  atom and the six neighbours on the same sub lattice can be found by rotating  $\mathbf{a}_1$  six times by  $1/6 * 2\pi = \pi/3$ :

$$\delta_{AA,1} = \mathbf{a}_1 = \frac{a}{2} \begin{pmatrix} 1 \\ \sqrt{3} \end{pmatrix} = a \begin{pmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix} = a \begin{pmatrix} \sin(\frac{\pi}{6}) \\ \cos(\frac{\pi}{6}) \end{pmatrix} \quad (\text{IV.6})$$

$$\delta_{AA,2} = a \begin{pmatrix} \sin(\frac{3\pi}{6}) \\ \cos(\frac{3\pi}{6}) \end{pmatrix} = a \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{IV.7})$$

$$\delta_{AA,3} = a \begin{pmatrix} \sin(\frac{5\pi}{6}) \\ \cos(\frac{5\pi}{6}) \end{pmatrix} = a \begin{pmatrix} \frac{1}{2} \\ -\frac{\sqrt{3}}{2} \end{pmatrix} \quad (\text{IV.8})$$

$$\delta_{AA,4} = a \begin{pmatrix} \sin(\frac{7\pi}{6}) \\ \cos(\frac{7\pi}{6}) \end{pmatrix} = a \begin{pmatrix} -\frac{1}{2} \\ -\frac{\sqrt{3}}{2} \end{pmatrix} \quad (\text{IV.9})$$

$$\delta_{AA,5} = a \begin{pmatrix} \sin(\frac{9\pi}{6}) \\ \cos(\frac{9\pi}{6}) \end{pmatrix} = a \begin{pmatrix} -1 \\ 0 \end{pmatrix} \quad (\text{IV.10})$$

$$\delta_{AA,6} = a \begin{pmatrix} \sin(\frac{11\pi}{6}) \\ \cos(\frac{11\pi}{6}) \end{pmatrix} = a \begin{pmatrix} -\frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{pmatrix} \quad (\text{IV.11})$$


First BZ vs this  
Gamma centered  
one

The primitive reciprocal lattice vectors  $\mathbf{b}_1, \mathbf{b}_2$  fulfill

$$\mathbf{a}_1 \cdot \mathbf{b}_1 = \mathbf{a}_2 \cdot \mathbf{b}_2 = 2\pi \quad (\text{IV.12})$$

$$\mathbf{a}_1 \cdot \mathbf{b}_2 = \mathbf{a}_2 \cdot \mathbf{b}_1 = 0, \quad (\text{IV.13})$$





**Figure IV.2:** *Graphene Brillouin Zone*

so we have:

$$\mathbf{b}_1 = \frac{2\pi}{a} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{3}} \end{pmatrix} \quad (\text{IV.14})$$

$$\mathbf{b}_2 = \frac{2\pi}{a} \begin{pmatrix} 1 \\ -\frac{1}{\sqrt{3}} \end{pmatrix} \quad (\text{IV.15})$$

Points of high symmetry in the Brillouin zone are:


$$\Gamma = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad (\text{IV.16})$$

$$\mathbf{M} = \frac{\pi}{a} \begin{pmatrix} 1 \\ \frac{1}{\sqrt{3}} \end{pmatrix} \quad (\text{IV.17})$$

$$\mathbf{K} = \frac{4\pi}{3a} \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (\text{IV.18})$$

## IV.2 EG-X MODEL

Graphene lattice and a site X.



**Figure IV.3:** *EG-X model*

Without interaction:

$$H_0 = -t_X \sum_{\langle ij \rangle, \sigma} d_{i, \sigma}^\dagger d_{j, \sigma} - t_{Gr} \sum_{\langle ij \rangle, \sigma} c_{i, \sigma}^{(A)\dagger} c_{j, \sigma}^{(B)} + V \sum_{i, \sigma \sigma'} d_{i, \sigma}^\dagger c_{i, \sigma'}^{(A)} + \text{h.c.} \quad (\text{IV.19})$$

with:

- $d$  operators on the X atom
- $c^{(\epsilon)}$  operators on the graphene site ( $\epsilon = A, B$ )
- $t_X$  NN hopping for X
- $t_{Gr}$  NN hopping of Gr
- $V$  hybridization between X and Graphene B sites

We can also introduce an onsite Hubbard interaction:

$$H_{\text{int}} = U_X \sum_i d_{i, \uparrow}^\dagger d_{i, \downarrow}^\dagger d_{i, \downarrow} d_{i, \uparrow} + U_{Gr} \sum_{i, \epsilon=A, B} c_{i, \uparrow}^{(\epsilon)\dagger} c_{i, \downarrow}^{(\epsilon)\dagger} c_{i, \downarrow}^\epsilon c_{i, \uparrow}^\epsilon \quad (\text{IV.20})$$

Hamilto-  
n orbital

Define sublattice index

$$\alpha = 1, 2, 3 \quad (\text{IV.21})$$

with  $1 \cong \text{Gr}_1, 2 \cong \text{Gr}_2, 3 \cong \text{X}$ . Then we can write the non-interacting term as

$$H_0 = - \sum_{\langle i,j \rangle, \alpha, \beta, \sigma} [\mathbf{t}]_{i\alpha, j\beta} c_{i\alpha}^\dagger c_{j\beta} \quad (\text{IV.22})$$

with the matrix

$$\mathbf{t} = \begin{pmatrix} 0 & t_{\text{Gr}} & 0 \\ t_{\text{Gr}} & 0 & -V\delta_{ij} \\ 0 & -V\delta_{ij} & t_{\text{X}} \end{pmatrix} \quad (\text{IV.23})$$

Add chemical potential:

$$-\mu \sum_{i\alpha\sigma} n_{i\alpha\sigma} \quad (\text{IV.24})$$

Also write the interaction part with  $\alpha$  (with changed signs compared to Niklas, to keep in line with papers about the attractive Hubbard model):

$$H_{int} = - \sum_{i\alpha} U_\alpha c_{i\alpha\uparrow}^\dagger c_{i\alpha\downarrow}^\dagger c_{i\alpha\downarrow} c_{i\alpha\uparrow} \quad (\text{IV.25})$$



## V RESULTS

Parameter sweeps using [34].

### V.1 GAPS

### V.2 SUPERFLUID WEIGHT

### V.3 BREAKDOWN OF SC WITH FINITE MOMENTUM

### V.4 COHERENCE LENGTH ETC.



## A EG-X HAMILTONIAN IN RECIPROCAL SPACE

Clean up this section

In the following chapter, the model Hamiltonian

$$H_0 = -t_X \sum_{\langle ij \rangle, \sigma} d_{i,\sigma}^\dagger d_{j,\sigma} - t_{Gr} \sum_{\langle ij \rangle, \sigma} c_{i,\sigma}^{(A)\dagger} c_{j,\sigma}^{(B)} + V \sum_{i,\sigma} d_{i,\sigma}^\dagger c_{i,\sigma}^{(A)} + \text{h.c.} \quad (\text{A.1})$$

will be treated to obtain the band structure. The first step is to write out the sums over nearest neighbors  $\langle i, j \rangle$  explicitly, writing  $\delta_X, \delta_\epsilon$  ( $\epsilon = A, B$ ) for the vectors to the nearest neighbors of the X atoms and Graphene A, B sites. Doing the calculation for example of the X atoms:

$$-t_X \sum_{\langle ij \rangle, \sigma} (d_{i,\sigma}^\dagger d_{j,\sigma} + d_{j,\sigma}^\dagger d_{i,\sigma}) = -\frac{t_X}{2} \sum_{i,\sigma} \sum_{\delta_X} d_{i,\sigma}^\dagger d_{i+\delta_X,\sigma} - \frac{t_X}{2} \sum_{j,\sigma} \sum_{\delta_X} d_{j,\sigma}^\dagger d_{j+\delta_X,\sigma} \quad (\text{A.2})$$

$$= -t_X \sum_{i,\sigma} \sum_{\delta_X} d_{i,\sigma}^\dagger d_{i+\delta_X,\sigma} \quad (\text{A.3})$$

The factor  $1/2$  in eq. (A.2) is to account for double counting when going to the sum over all lattice sites  $i$ . By relabeling  $j \rightarrow i$  in the second sum, the two sum are the same and eq. (A.3) is obtained. Using now the discrete Fourier transform

$$c_i = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}}, \quad c_i^\dagger = \frac{1}{\sqrt{N}} \sum_{\mathbf{k}} e^{-i\mathbf{k}\mathbf{r}_i} c_{\mathbf{k}}^\dagger \quad (\text{A.4})$$

with the completeness relation

$$\sum_i e^{i\mathbf{k}\mathbf{r}_i} e^{-i\mathbf{k}'\mathbf{r}_i} = N \delta_{\mathbf{k},\mathbf{k}'}, \quad (\text{A.5})$$

eq. (A.3) reads:

$$-t_X \frac{1}{N} \sum_{i,\sigma} \sum_{\delta_X} d_{i,\sigma}^\dagger d_{i+\delta_X,\sigma} = -t_X \frac{1}{N} \sum_{i,\sigma} \sum_{\mathbf{k},\mathbf{k}',\delta_X} (e^{-i\mathbf{k}\mathbf{r}_i} d_{\mathbf{k},\sigma}^\dagger) (e^{i\mathbf{k}'\mathbf{r}_i} e^{i\mathbf{k}'\delta_X} d_{\mathbf{k}',\sigma}) \quad (\text{A.6})$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\delta_X,\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k}',\sigma} e^{i\mathbf{k}'\delta_X} \sum_i e^{-i\mathbf{k}\mathbf{r}_i} e^{i\mathbf{k}'\mathbf{r}_i} \quad (\text{A.7})$$

$$= -t_X \frac{1}{N} \sum_{\mathbf{k},\mathbf{k}',\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k}',\sigma} \sum_{\delta_X} e^{i\mathbf{k}'\delta_X} (N \delta_{\mathbf{k},\mathbf{k}'}) \quad (\text{A.8})$$

$$= -t_X \sum_{\mathbf{k},\sigma} d_{\mathbf{k},\sigma}^\dagger d_{\mathbf{k},\sigma} \sum_{\delta_X} e^{i\mathbf{k}\delta_X}. \quad (\text{A.9})$$

This part is now diagonal in  $\mathbf{k}$  space. The nearest neighbours vectors  $\delta_X$  for the X atoms are the vectors  $\delta_{AA,i}$  from ?? . With that, the sum over  $\delta_X$  can be explicitly calculated:

Correct exp expressions

Example for a vector product

$$f_X(\mathbf{k}) = -t_X \sum_{\delta_X} e^{i\mathbf{k}\delta_X} \quad (\text{A.10})$$

$$= -t_X \left[ \exp \left( ia \left( \frac{k_x}{2} + \frac{\sqrt{3}k_y}{2} \right) \right) + e^{iak_x} + e^{ia(\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} \right. \quad (\text{A.11})$$

$$\left. + e^{ia(-\frac{k_x}{2} - \frac{\sqrt{3}k_y}{2})} + e^{-iak_x} + e^{ia(-\frac{k_x}{2} + \frac{\sqrt{3}k_y}{2})} \right] \quad (\text{A.12})$$

$$= -t_X \left( 2 \cos(ak_x) + 2e^{ia\frac{\sqrt{3}k_y}{2}} \cos\left(\frac{a}{2}k_x\right) + 2e^{-ia\frac{\sqrt{3}k_y}{2}} \cos\left(\frac{a}{2}k_x\right) \right) \quad (\text{A.13})$$

$$= -2t_X \left( \cos(ak_x) + 2 \cos\left(\frac{a}{2}k_x\right) \cos\left(\sqrt{3}\frac{a}{2}k_y\right) \right). \quad (\text{A.14})$$

The same can be done for the hopping between Graphene sites, for example :

$$-t_{\text{Gr}} \sum_{\langle ij \rangle, \sigma, \sigma'} c_{i,\sigma}^{(A)\dagger} c_{j,\sigma'}^{(B)} = -t_{\text{Gr}} \sum_{i,\sigma,\sigma'} \sum_{\delta_{AB}} c_{i,\sigma}^{(A)\dagger} c_{i+\delta_{AB},\sigma'}^{(B)} \quad (\text{A.15})$$


$$= -t_{\text{Gr}} \sum_{\mathbf{k},\sigma,\sigma'} c_{\mathbf{k},\sigma}^{(A)\dagger} c_{\mathbf{k},\sigma'}^{(B)} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (\text{A.16})$$

Show that!

We note

$$\sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} = \left( \sum_{\delta_{BA}} e^{i\mathbf{k}\delta_{BA}} \right)^* = \sum_{\delta_{BA}} e^{-i\mathbf{k}\delta_{BA}} \quad (\text{A.17})$$





**Figure A.1:** Bands of the non-interacting EG-X model. All the bands are spin-degenerate.

and calculate

$$f_{Gr} = -t_{Gr} \sum_{\delta_{AB}} e^{i\mathbf{k}\delta_{AB}} \quad (\text{A.18})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{i\frac{a}{2\sqrt{3}}(\sqrt{3}k_x - k_y)} + e^{i\frac{a}{2\sqrt{3}}(-\sqrt{3}k_x - k_y)} \right) \quad (\text{A.19})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + e^{-i\frac{a}{2\sqrt{3}}k_y} \left( e^{i\frac{a}{2}k_x} + e^{-i\frac{a}{2}k_x} \right) \right) \quad (\text{A.20})$$

$$= -t_{Gr} \left( e^{i\frac{a}{\sqrt{3}}k_y} + 2e^{-i\frac{a}{2\sqrt{3}}k_y} \cos\left(\frac{a}{2}k_x\right) \right) \quad (\text{A.21})$$

All together, we get:

$$H_0 = \sum_{\mathbf{k}, \sigma, \sigma'} \begin{pmatrix} c_{\mathbf{k}, \sigma}^{A, \dagger} & c_{\mathbf{k}, \sigma}^{B, \dagger} & d_{\mathbf{k}, \sigma}^{\dagger} \end{pmatrix} \begin{pmatrix} 0 & f_{Gr} & V \\ f_{Gr}^* & 0 & 0 \\ V & 0 & f_X \end{pmatrix} \begin{pmatrix} c_{\mathbf{k}, \sigma}^A \\ c_{\mathbf{k}, \sigma}^B \\ d_{\mathbf{k}, \sigma} \end{pmatrix} \quad (\text{A.22})$$

The band structure for the non-interacting EG-X model is easily obtained by diagonalising the matrix in eq. ???. This was done in fig. A.1.

Values used for calculation:

- $a_0 = 1$
- $t_{Gr} = 1$
- $t_X = 0.01$

$V$  is the control parameter. A range from  $V = 0.1$  to  $V = 2$  can be mapped onto materials in experiment.



## B NOTES ON THE COMPUTATIONAL IMPLEMENTATION

All the code is available at [github.com/Ruberhauptmann/quant-met](https://github.com/Ruberhauptmann/quant-met).

All the data,

Data availability

For reproducibility, Datalad [35] is used.

The implementation relies on the work of many contributors of packages in Python's ecosystem, most important among them NumPy [36], SciPy [37], Matplotlib [38], Pandas [39, 40] and Parasweep [34].

What software for what?



## BIBLIOGRAPHY

- [1] U. of Chicago. *Annual Register*. 1893-1930., 1896. 462 pp. (cit. on p. 1).
- [2] H. K. Onnes. "Further Experiments with Liquid Helium. G. On the Electrical Resistance of Pure Metals, Etc. VI. On the Sudden Change in the Rate at Which the Resistance of Mercury Disappears." In: *Through Measurement to Knowledge: The Selected Papers of Heike Kamerlingh Onnes 1853–1926*. Ed. by K. Gavroglu and Y. Goudaroulis. Dordrecht: Springer Netherlands, 1991, pp. 267–272. ISBN: 978-94-009-2079-8. DOI: 10.1007/978-94-009-2079-8\_17 (cit. on p. 1).
- [3] W. Meissner and R. Ochsenfeld. "Ein neuer Effekt bei Eintritt der Supraleitfähigkeit". In: *Naturwissenschaften* 21.44 (Nov. 1, 1933), pp. 787–788. ISSN: 1432-1904. DOI: 10.1007/BF01504252 (cit. on pp. 1, 9).
- [4] J. Bardeen, L. N. Cooper, and J. R. Schrieffer. "Theory of Superconductivity". In: *Physical Review* 108.5 (Dec. 1, 1957), pp. 1175–1204. DOI: 10.1103/PhysRev.108.1175 (cit. on pp. 1, 10).
- [5] J. G. Bednorz and K. A. Müller. "Possible High T<sub>c</sub> Superconductivity in the Ba–La–Cu–O System". In: *Zeitschrift für Physik B Condensed Matter* 64.2 (June 1, 1986), pp. 189–193. ISSN: 1431-584X. DOI: 10.1007/BF01303701 (cit. on p. 1).
- [6] S.-i. Uchida et al. "High T<sub>c</sub> Superconductivity of La-Ba-Cu Oxides". In: *Japanese Journal of Applied Physics* 26 (1A Jan. 1, 1987), p. L1. ISSN: 1347-4065. DOI: 10.1143/JJAP.26.L1 (cit. on p. 1).
- [7] D. Rybicki et al. "Perspective on the Phase Diagram of Cuprate High-Temperature Superconductors". In: *Nature Communications* 7.1 (May 6, 2016), p. 11413. ISSN: 2041-1723. DOI: 10.1038/ncomms11413 (cit. on p. 1).
- [8] P. A. Rinck. *Magnetic Resonance in Medicine - A Critical Introduction*. ISBN: 978-628-01-2260-1. URL: <http://www.magnetic-resonance.org/> (cit. on p. 1).

- [9] A. Tollestrup and E. Todesco. “The Development of Superconducting Magnets for Use in Particle Accelerators: From the Tevatron to the LHC”. In: *Reviews of Accelerator Science and Technology* 1.01 (2008), pp. 185–210 (cit. on p. 1).
- [10] K. Irwin and G. Hilton. “Transition-Edge Sensors”. In: *Cryogenic Particle Detection*. Ed. by C. Enss. Berlin, Heidelberg: Springer, 2005, pp. 63–150. ISBN: 978-3-540-31478-3. DOI: 10.1007/10933596\_3 (cit. on p. 1).
- [11] M. I. Faley et al. “High-Tc SQUID Biomagnetometers”. In: *Superconductor Science and Technology* 30.8 (July 2017), p. 083001. ISSN: 0953-2048. DOI: 10.1088/1361-6668/aa73ad (cit. on p. 1).
- [12] A. M. Klushin et al. “Present and Future of High-Temperature Superconductor Quantum-Based Voltage Standards”. In: *IEEE Instrumentation & Measurement Magazine* 23.2 (Apr. 2020), pp. 4–12. ISSN: 1941-0123. DOI: 10.1109/MIM.2020.9062678 (cit. on p. 1).
- [13] B. D. Josephson. “Possible New Effects in Superconductive Tunnelling”. In: *Physics Letters* 1.7 (July 1, 1962), pp. 251–253. ISSN: 0031-9163. DOI: 10.1016/0031-9163(62)91369-0 (cit. on p. 1).
- [14] Y. Cao et al. “Unconventional Superconductivity in Magic-Angle Graphene Superlattices”. In: *Nature* 556.7699 (Apr. 2018), pp. 43–50. ISSN: 1476-4687. DOI: 10.1038/nature26160 (cit. on p. 2).
- [15] P. Coleman. *Introduction to Many-Body Physics*. Cambridge University Press, Nov. 2015. ISBN: 978-0-521-86488-6. DOI: 10.1017/CBO9781139020916 (cit. on pp. 3, 9, 15).
- [16] M. Tinkham. *Introduction to Superconductivity*. 2. ed. International Series in Pure and Applied Physics. New York: McGraw-Hill, 1996. 454 pp. ISBN: 978-0-07-064878-4 978-0-07-114782-8 (cit. on p. 3).
- [17] H. Bruus and K. Flensberg. *Many-Body Quantum Theory in Condensed Matter Physics: An Introduction*. Oxford Graduate Texts. Oxford, New York: Oxford University Press, Nov. 11, 2004. 466 pp. ISBN: 978-0-19-856633-5 (cit. on pp. 3, 12, 14).
- [18] A. I. Larkin and A. A. Varlamov. *Theory of Fluctuations in Superconductors*. Oxford Science Publications 127. Oxford Oxford: Clarendon Press Oxford University Press, 2005. ISBN: 978-0-19-852815-9 (cit. on p. 3).

- 
- [19] L. D. Landau. “On The Theory of Superconductivity”. In: *Collected Papers of L.D. Landau*. Ed. by D. Ter haar. Pergamon, Jan. 1, 1965, pp. 546–568. ISBN: 978-0-08-010586-4. DOI: 10.1016/B978-0-08-010586-4.50078-X (cit. on pp. 3, 10).
- [20] N. Witt et al. *Bypassing the Lattice BCS-BEC Crossover in Strongly Correlated Superconductors: Resilient Coherence from Multiorbital Physics*. July 11, 2024. DOI: 10.48550/arXiv.2310.09063. Pre-published (cit. on p. 8).
- [21] N. F. Q. Yuan and L. Fu. “Supercurrent Diode Effect and Finite-Momentum Superconductors”. In: *Proceedings of the National Academy of Sciences* 119.15 (Apr. 12, 2022), e2119548119. DOI: 10.1073/pnas.2119548119 (cit. on p. 9).
- [22] R. Shimano and N. Tsuji. “Higgs Mode in Superconductors”. In: *Annual Review of Condensed Matter Physics* 11 (Volume 11, 2020 Mar. 10, 2020), pp. 103–124. ISSN: 1947-5454, 1947-5462. DOI: 10.1146/annurev-conmatphys-031119-050813 (cit. on p. 9).
- [23] F. London. “A New Conception of Supraconductivity”. In: *Nature* 140.3549 (Nov. 1, 1937), pp. 793–796. ISSN: 1476-4687. DOI: 10.1038/140793a0 (cit. on p. 9).
- [24] L. N. Cooper. “Bound Electron Pairs in a Degenerate Fermi Gas”. In: *Physical Review* 104.4 (Nov. 15, 1956), pp. 1189–1190. DOI: 10.1103/PhysRev.104.1189 (cit. on p. 10).
- [25] J. Hubbard and B. H. Flowers. “Electron Correlations in Narrow Energy Bands”. In: *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences* 276.1365 (Nov. 26, 1963), pp. 238–257. DOI: 10.1098/rspa.1963.0204 (cit. on p. 10).
- [26] J. Kanamori. “Electron Correlation and Ferromagnetism of Transition Metals”. In: *Progress of Theoretical Physics* 30.3 (Sept. 1, 1963), pp. 275–289. ISSN: 0033-068X. DOI: 10.1143/PTP.30.275 (cit. on p. 10).
- [27] M. C. Gutzwiller. “Effect of Correlation on the Ferromagnetism of Transition Metals”. In: *Physical Review Letters* 10.5 (Mar. 1, 1963), pp. 159–162. DOI: 10.1103/PhysRevLett.10.159 (cit. on p. 10).

- [28] M. Qin et al. “The Hubbard Model: A Computational Perspective”. In: *Annual Review of Condensed Matter Physics* 13 (Volume 13, 2022 Mar. 10, 2022), pp. 275–302. issn: 1947-5454, 1947-5462. doi: 10.1146/annurev-conmatphys-090921-033948 (cit. on p. 11).
- [29] R. Micnas, J. Ranninger, and S. Robaszkiewicz. “Superconductivity in Narrow-Band Systems with Local Nonretarded Attractive Interactions”. In: *Reviews of Modern Physics* 62.1 (Jan. 1, 1990), pp. 113–171. doi: 10.1103/RevModPhys.62.113 (cit. on p. 11).
- [30] J. J. Kinnunen et al. “The Fulde–Ferrell–Larkin–Ovchinnikov State for Ultracold Fermions in Lattice and Harmonic Potentials: A Review”. In: *Reports on Progress in Physics* 81.4 (Feb. 2018), p. 046401. issn: 0034-4885. doi: 10.1088/1361-6633/aaa4ad (cit. on p. 12).
- [31] A. Georges et al. “Dynamical Mean-Field Theory of Strongly Correlated Fermion Systems and the Limit of Infinite Dimensions”. In: *Reviews of Modern Physics* 68.1 (Jan. 1, 1996), pp. 13–125. doi: 10.1103/RevModPhys.68.13 (cit. on p. 15).
- [32] J. P. Provost and G. Vallee. “Riemannian Structure on Manifolds of Quantum States”. In: *Communications in Mathematical Physics* 76.3 (Sept. 1, 1980), pp. 289–301. issn: 1432-0916. doi: 10.1007/BF02193559 (cit. on p. 17).
- [33] G. Yang et al. “Structure of Graphene and Its Disorders: A Review”. In: *Science and Technology of Advanced Materials* 19.1 (Aug. 29, 2018), pp. 613–648. issn: 1468-6996. doi: 10.1080/14686996.2018.1494493 (cit. on p. 19).
- [34] E. Bach. “Parasweep: A Template-Based Utility for Generating, Dispatching, and Post-Processing of Parameter Sweeps”. In: *SoftwareX* 13 (Jan. 1, 2021), p. 100631. issn: 2352-7110. doi: 10.1016/j.softx.2020.100631 (cit. on pp. 25, 31).
- [35] Y. O. Halchenko et al. “DataLad: Distributed System for Joint Management of Code, Data, and Their Relationship”. In: *Journal of Open Source Software* 6.63 (July 1, 2021), p. 3262. issn: 2475-9066. doi: 10.21105/joss.03262 (cit. on p. 31).



- 
- [36] C. R. Harris et al. “Array Programming with NumPy”. In: *Nature* 585.7825 (Sept. 2020), pp. 357–362. ISSN: 1476-4687. DOI: 10.1038/s41586-020-2649-2 (cit. on p. 31).
  - [37] P. Virtanen et al. “SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python”. In: *Nature Methods* 17.3 (Mar. 2020), pp. 261–272. ISSN: 1548-7105. DOI: 10.1038/s41592-019-0686-2 (cit. on p. 31).
  - [38] J. D. Hunter. “Matplotlib: A 2D Graphics Environment”. In: *Computing in Science & Engineering* 9.3 (May 2007), pp. 90–95. ISSN: 1558-366X. DOI: 10.1109/MCSE.2007.55 (cit. on p. 31).
  - [39] W. McKinney. “Data Structures for Statistical Computing in Python”. In: *Python in Science Conference*. Austin, Texas, 2010, pp. 56–61. DOI: 10.25080/Majora-92bf1922-00a (cit. on p. 31).
  - [40] T. pandas development team. *Pandas-Dev/Pandas: Pandas*. Version v2.2.3. Zenodo, Sept. 20, 2024. DOI: 10.5281/ZENODO.3509134 (cit. on p. 31).



## NOT CITED

- [41] K. H. Bennemann and J. B. Ketterson, eds. *Superconductivity*. Berlin, Heidelberg: Springer Berlin Heidelberg, 2008. ISBN: 978-3-540-73252-5 978-3-540-73253-2. DOI: 10.1007/978-3-540-73253-2.
- [42] S. Peotta and P. Törmä. “Superfluidity in Topologically Nontrivial Flat Bands”. In: *Nature Communications* 6.1 (Nov. 20, 2015), p. 8944. ISSN: 2041-1723. DOI: 10.1038/ncomms9944.



## LIST OF SYMBOLS

Symbol	Meaning	Definition
$\tau$	general imaginary time variable	13
$C_{AB}(\tau, 0)$	Correlation function in imaginary time	14