

PT Name: Age / Sex: Ref By:	Reg No: Barcode: INV: 17 OH Progesterone	Reg. Date: Rep. Date: Sample: Serum
-----------------------------------	--	---

17 OH Progesterone

TEST RESULT

TEST DESCRIPTION	RESULT	UNITS	
17 OH Progesterone METHOD: CLIA		ng/mL	

CATEGORY	RANGE (NG/ML)
< 1 yr	< 3.00
> 1 yr	< 2.00
Adult Males	0.63 - 2.15
Postmenopausal Females	0.19 - 0.71

PREMENOPAUSAL FEMALES	RANGE (NG/ML)
Follicular phase	0.32 - 1.47
Luteal phase	0.25 - 2.91
Contraception	0.20 - 1.90

CLINICAL SIGNIFICANCE

The adrenal glands, ovaries, testes, and placenta produce OHPG. It is hydroxylated at the 11 and 21 position to produce cortisol. Deficiency of either 11- or 21-hydroxylase results in decreased cortisol synthesis, and feedback inhibition of adrenocorticotrophic hormone (ACTH) secretion is lost. Consequent increased pituitary release of ACTH increases production of OHPG. But, if 17-alpha-hydroxylase or 3-beta-hydroxysteroid dehydrogenase type 2 are deficient, OHPG levels are low with possible increase in progesterone or pregnenolone respectively.

OHPG is bound to both corticosteroid binding globulin and albumin and total OHPG is measured in this assay. OHPG is converted to pregnanetriol, which is conjugated and excreted in the urine. In all instances, more specific tests are available to diagnose disorders or steroid metabolism than pregnanetriol measurement.