

Exploratory data analysis

Objective

- Get the **quick idea about data**
 - visualization is the easiest way
 - check descriptive statistics
- **Data cleaning process** to reduce the number of data problems in the future
 - handle missing data, outliers or typo etc.
 - need to be careful!
- **Explore your data** to determine whether the model assumptions are met etc.
 - E.g., check normality of data

Visualization and descriptive statistics

- **Visualization**
 - Histogram
 - Boxplot
 - Scatter plot (to find the relationship btw 2 variables)
- **Descriptive statistics**
 - Mean, median, variance, skewness, kurtosis etc.
 - Correlation (for 2 variables)
- Get rough idea about the distribution of data
- Check outliers or missingness
- In general, to check normality of data

Visualization and descriptive statistics

- Skewed right: $\text{mean} > \text{median}$
- Skewed left: $\text{mean} < \text{median}$
 - **Robustness** of median.
 - Able to guess its skewness based on mean and median values
- Normal distribution is symmetric, bell-shaped
 - Inverse is NOT true e.g., Cauchy distribution, t-distribution
- Able to check its normality (informally) based on visual and descriptive statistics

Example : airquality

- Daily air quality measurements in New York, May to September 1973. (R built-in data)
- 154 observations on 6 variables – Ozone, Solar R, Wind, ...


```
> head(airquality,10)
```

	Ozone	Solar.R	Wind	Temp	Month	Day
1	41	190	7.4	67	5	1
2	36	118	8.0	72	5	2
3	12	149	12.6	74	5	3
4	18	313	11.5	62	5	4
5	NA	NA	14.3	56	5	5
6	28	NA	14.9	66	5	6
7	23	299	8.6	65	5	7
8	19	99	13.8	59	5	8
9	8	19	20.1	61	5	9
10	NA	194	8.6	69	5	10


```
> |
```

```
hist(airquality$Ozone,main="Ozone",xlab="Ozone")
```

Provides the distribution of the data. This can also be used to assess potential outlier concerns.



```
boxplot(airquality$Ozone, ylab="Ozone")
points(mean(airquality$Ozone, na.rm=TRUE), col="red")
```



Example of descriptive statistics

```
summary(airquality$Ozone, na.rm=TRUE)
```

```
##      Min. 1st Qu.  Median    Mean 3rd Qu.    Max.      NA's  
##      1.00   18.00   31.50   42.13   63.25   168.00       37
```

```
mean(airquality$Ozone, na.rm=TRUE) -> mean
```

```
## [1] 42.12931
```

```
var(airquality$Ozone, na.rm=TRUE) -> variance
```

```
## [1] 1088.201
```

```
skewness(airquality$Ozone, na.rm=TRUE) -> skewness
```

```
## [1] 1.209866
```

```
range(airquality$Ozone, na.rm=TRUE) -> range [min,max]
```

```
## [1]    1 168
```


Important statistical assumptions

- Normality
 - Why normality check is important?
 - 1) When conducting a **t-test** or **ANOVA**, normality assumption is required
 - 2) When using **correlation** and **regression techniques**, lack of normality and outliers impact your conclusions
 - There are a lot of tests one can use to check for normality and outliers in the data.

Inference based on Normality


- Under normality assumption, we can perform following tests.
 - ✓ One-sample t-test
(e.g., test if iphone battery life span > 2 years)
 - ✓ Two-sample t-test
(e.g., test if iphone and galaxy have the same life span)
 - ✓ ANOVA test (simply speaking, comparing group means among more than two groups)
(e.g., test among iphone, galaxy and Android phone)

Detection of Normality

- How to check Normality?
 - ✓ **Qualitatively check** by looking at:
 - : histogram, boxplot, quantile-quantile plot (QQ plot) etc..
 - ✓ **Quantitative check** by formal test
 - : Sharpiro-Wilk test ...
- For a comparison among groups (e.g., t-test, ANOVA), **normality check should be conducted by groups**
- If **at least one group** does not follow normality, t-test or ANOVA conclusions may **NOT** be valid.

```
qqnorm(airquality$Ozone); qqline(airquality$Ozone, col = 2)
```

Normal Q-Q Plot



Quantile-Quantile Plots

(a.k.a., Q-Q plots): A useful diagnostics of how well a specified theoretical distribution fits your data. If the quantiles of the theoretical and data distributions agree, the plotted points fall on or near the line.

Shapiro-Wilk Normality test

```
shapiro.test(airquality$Ozone)

##
##  Shapiro-Wilk normality test
##
## data:  airquality$Ozone
## W = 0.87867, p-value = 2.79e-08
```

H0: Data follows normal distribution

H1: Data does not follow normal distribution

- If p-value is larger than significance level (in general $\alpha=0.05$), we do not enough evidence to reject the null hypothesis, thus our conclusion is - data follows normal distribution
- If p-value is smaller than significance level, we have enough evidence to reject the null hypothesis, thus our conclusion is – data does not follow Normal distribution