





গণপ্রজাতন্ত্রী বাংলাদেশ



I graduated in 1999 from Munkkiniemi high school. Mathematics was a subject I had naturally thrived in – so, after some bumps and turns, I found myself at the University of Helsinki studying mathematics. And yeah, indeed, I love mathematics—I love the apparent universality of it. This subject is without a doubt debatable, but, at least in some sense, I like to think that mathematical truths are universal in the truest sense of the word; they are eternal, and they are the same everywhere, regardless of the physical universe we live in. Aliens in another galaxy will end up with the same mathematical truths we do. Aliens in another universe will end up with the same mathematical truths we do. Mathematics has the power to explain what we see in our everyday life. Mathematics is not only natural science but a form of art and poetry. Mathematics is music—music is mathematics.

While studying mathematics, physics, and computer science, I took some courses on economics. That inspired me to write my bachelor's thesis on optimal control theory. I worked on the problem of how increasing public investments affects the GDP. I did not find any breakthrough, but it was an intriguing subject. I proceeded with my graduate studies studying applied mathematics. I studied subjects like partial differential equations, functional analysis, dynamical systems, and—the University of Helsinki's pride—complex analysis. (My thesis advisor said that, in a moral sense, you cannot graduate from the University of Helsinki without taking some courses on Complex analysis, because a lot of the discipline has been developed at the university.) In addition, as a more “practical” subject, I studied some inverse problems. Summa summarum, I studied a wide range of fields in mathematics.

During my graduate studies, I spent half a year in Utrecht, Netherlands, studying more applied analysis of varying subjects (searching periodic orbits in the Lorentz attractor as an example of a course—that I failed). At Utrecht University, I got the inspiration for the subject for my future master's thesis; the finite element method (FEM). After I got back to Helsinki from the exchange, I had a chance to study more about the FEM in Aalto University's courses. (Aalto University is a consortium of the Helsinki University of Technology, the Helsinki School of Economics, and the University of Art and Design Helsinki.) While writing my thesis I also taught basic mathematics courses at the University of Helsinki and gained valuable experience in the pedagogical area.

In the binge of graduation, I started to look for future opportunities. I looked for coding jobs in Helsinki and Tallinn, jobs for mathematicians in the mapping industry, continuing at some universities to pursue a Ph.D., etc. I am glad I had the chance to use my creativity and continue in Aalto University's Department of Signal Processing and Acoustics to research low earth orbit satellite communications. The research methodology was from a stochastic geometry perspective, which was well aligned with my mathematical background.

My professional ambitions are in improving the lives of people globally. Communications play an essential in the picture. (But contain some challenging problems also, as we have seen with social media.) Through effective communication, we can share knowledge, control resources, discuss issues, etc.—however, globally, the communication infrastructure is still not nearly complete. My interests contain, but are not limited to, communications, particularly wireless networks and signal processing. Modulation and demodulation, bandpass and passband. My dream is to share my knowledge in the process toward a free and honest world. (Pardon me for the cliches.)





PARADOXES







BEAR



1995

[illegible]





$$K_P(\tau) = k \exp \left\{ -D \tau^2 \log(2) \right\} = \frac{p_t^2}{2 \log(2)} \pi \lambda \left(\frac{h \varphi_{\text{RX}}}{\sin^2(\epsilon)} \right)^2 \exp \left\{ -v_{\text{sat}}^2 \tau^2 / (h^2 \varphi_{\text{RX}}^2) \log(2) \right\},$$















ESPRESSO













A pixelated, grayscale illustration of a stylized letter 'I'. The letter is composed of a thick vertical stem and a horizontal crossbar. To the left of the letter is a vertical bar with a gradient from light to dark gray. To the right of the letter is a small, abstract shape resembling a stylized 'L' or a corner bracket, also rendered in a pixelated, grayscale style. The entire image has a low-resolution, digital-art aesthetic.



expanding world + new world



Fig 1(a): True $P(t)$, GPR estimate & forecast, and 300-sample MA & forecast





Fig 1(b): Non-stationary Gaussian noise $I(t)$ with sampled observations









Qatar Airways









$$S_{RX}(t) = \frac{A}{l(d_0)} \sqrt{\frac{K}{K+1}} e^{-2i\pi\tau_0(t)f_c} e^{-2i\pi\theta} + \sum_{j=1}^{100} \frac{A}{l(d_j) \sqrt{100} \sqrt{K+1}} e^{-2i\pi\tau_j(t)f_c} e^{-2i\pi\theta},$$











2020 = 2020







QWERTY

$$v = \sqrt{\frac{GM}{h + R_{\oplus}}},$$

WORLDWIDE











2

=

12

.

100



$$T_c = \frac{1}{8D_g} \approx 10^{-3}$$
























$$1 = \frac{1}{e^{\frac{\pi^2}{2VP}}}$$






$$] = 1 - \left(1 - e^{-\frac{\pi^2}{2\sqrt{P/N}}} \right)^N,$$





192








$$t \rightarrow \text{order} \rightarrow \frac{\sin(2\pi B t)}{\pi t}$$





$$\mathbb{Z} \ni i \mapsto \frac{\sin\left(2\pi \frac{B_L}{f_c} i\right)}{\pi i},$$

$$i \mapsto \delta[i] - \frac{\sin(2\pi \frac{B_H}{f_c} i)}{\pi i},$$



WAVELENGTH



WINNERS

$$\mu = \exp\left(\mu_{\text{LIN}} + \frac{\sigma_{\text{LIN}}^2}{2}\right)$$

$$O(2\pi) \cong SO(2) \cong \mathbb{R}/2\pi\mathbb{Z}$$











FOR THE WIN





A pixelated, black and white representation of the text "Xmas is not a religion". The text is rendered in a jagged, digital font style, with each character composed of small, distinct pixel blocks. The letters are arranged in two rows: "Xmas is" on the top row and "not a religion" on the bottom row. The overall aesthetic is reminiscent of early computer graphics or low-resolution digital art.

$$S(t) = \sum_{n=0}^{N-1} x[n]$$
















$$H = \begin{pmatrix} -a & -0 & -1 & 0 & 0 & 0 & ar & 0 & r \\ 0 & 0 & 0 & -a & -0 & -1 & 0 & 0 & 0 \\ -c & -b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & -b^2/a & -1 & cr & rb^2/a & r \\ a & 0 & -1 & 0 & 0 & 0 & ar & 0 & -r \\ 0 & 0 & 0 & a & 0 & -1 & 0 & 0 & 0 \\ -c & b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & b^2/a & -1 & -rc & rb^2/ar & -r \end{pmatrix}$$

$$\lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X \geq x)}{x},$$







$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x}$$

$$= \lim_{x \rightarrow \infty} \frac{-x/\theta}{x} = 1/\theta.$$



$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(1 - \gamma(k, x/\theta)/\Gamma(k))}{x}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(k, x/\theta)/\Gamma(k))}{x}$$

$$\begin{aligned}
 &\stackrel{(a)}{=} \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(k, x/\theta))}{x} = \lim_{x \rightarrow \infty} - \frac{\log(x^{k-1} e^{-x/\theta})}{x}
 \end{aligned}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x} = 1/\theta.$$



$$\lim_{x \rightarrow \infty} \frac{\Gamma(s, x)}{x^{s-1} e^{-x}} = 1.$$







$$\begin{aligned}
&= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = - \lim_{x \rightarrow \infty} \frac{\log \operatorname{erfc} \left(\frac{x - \mu}{\sigma \sqrt{2}} \right)}{x} \stackrel{(b)}{\geq} - \lim_{x \rightarrow \infty} \frac{\log \left(\sqrt{\frac{e}{2\pi}} e^{-2 \left(\frac{x - \mu}{\sigma \sqrt{2}} \right)^2} \right)}{x}
\end{aligned}$$

$$\geq -\lim_{x \rightarrow \infty} \frac{\log \left(e^{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2} \right)}{x} = -\lim_{x \rightarrow \infty} \frac{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2}{x} = \frac{1}{\sigma^2} \lim_{x \rightarrow \infty} \frac{x^2 - 2x\mu + \mu^2}{x} = \infty,$$



$$\psi(x) \geq \sqrt{\frac{2e}{\pi}} \frac{\sqrt{\beta - 1}}{\beta} e^{-\beta x^2},$$









































Q1 2024

2023年12月21日



$$\mathbb{E}[(S_1 + S_2)^2] = \int_0^1 (\cos(2\pi t) + \cos(2\pi t + \pi))^2 dt = \int_0^1 0 dt = 0,$$



1992 + 1992

$$= \int_0^1 \cos^2(2\pi t) dt + \int_0^1 \cos^2(2\pi t + \pi) dt$$

$$= \int_0^1 2 \cos^2(2\pi t) dt = \int_0^1 \cos(4\pi t) dt + 1 = 1.$$





$$E[S_1^2 + S_2^2] = E[S_1^2] + E[S_2^2] = E[S_1^2] + 2E[S_1 S_2] + E[S_2^2]$$

1991 + 1991 + 1991

1992-93

$$\mathbb{E}[S_1 S_2] = \int_0^1 \cos(2\pi t) \cos(2\pi t + \pi) dt = \int_0^1 -\cos^2(2\pi t) dt = -\frac{1}{2} \int_0^1 \cos(4\pi t) dt - \frac{1}{2} = -\frac{1}{2},$$

Q1

2

Q2

E

Q3

4

Q4

2024 + 2025

2023-03-20

$$\mathbb{E} \phi_1, \phi_2 [\mathbb{E} [S_1^2 + S_2^2]] = \mathbb{E} \phi_1 [\mathbb{E} [S_1^2]] + \mathbb{E} \phi_2 [\mathbb{E} [S_2^2]] + \mathbb{E} \phi_1, \phi_2 [\mathbb{E} [S_1 S_2]]$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_1))^2 dt d\phi_1 + \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_2))^2 dt d\phi_2$$

$$+ \frac{1}{2\pi} \int_0^{2\pi} \int_0^{2\pi} \int_0^1 \cos(2\pi t + \phi_1) \cos(2\pi t + \phi_2) dt d\phi_1 d\phi_2$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_1))^2 dt d\phi_1 + \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_2))^2 dt d\phi_2 = 1/2 + 1/2 = 1.$$

1992

212122222





$${}_3F_2(1,1,1+b;2,2;x)=\sum_{n=0}^{\infty}\frac{(1)_n(1)_n(1+b)_n}{(2)_n(2)_n}\frac{x^n}{n!}$$

$$= \sum_{n=0}^{\infty} \frac{(1+b)_n}{(n+1)^2 n!} x^n = \frac{1}{b!} \sum_{n=0}^{\infty} \frac{(n+1)_b}{(n+1)^2} x^n$$

$$\underline{\underline{(a)}} \quad \frac{1}{b!} \sum_{n=0}^{\infty} \frac{\sum_{k=1}^b \begin{bmatrix} b \\ k \end{bmatrix} (n+1)^k}{(n+1)^2} x^n$$

$$\begin{aligned}
 &= \frac{1}{b!} \sum_{k=1}^b \left[\begin{matrix} b \\ k \end{matrix} \right] \sum_{n=0}^{\infty} \frac{x^n}{(n+1)^{2-k}} \stackrel{(b)}{=} \frac{1}{b!} \sum_{k=1}^b \left[\begin{matrix} b \\ k \end{matrix} \right] \frac{\operatorname{Li}_{2-k}(x)}{x},
 \end{aligned}$$

b

16

