



গণপ্রজাতন্ত্রী বাংলাদেশ

I graduated in 2009 from Munkkiniemi high school. Mathematics was a subject I had naturally thrived in – so, after some bumps and turns, I found myself at the University of Helsinki studying mathematics. And yeah, indeed, I love mathematics – I love the apparent universality of it. This subject is without a doubt debatable, but, at least in some sense, I like to think that mathematical truths are universal in the truest sense of the word; they are eternal, and they are the same everywhere, regardless of the physical universe we live in. Aliens in another galaxy will end up with the same mathematical truths we do. Aliens in another universe will end up with the same mathematical truths we do. Mathematics has the power to explain what we see in our everyday life. Mathematics is not only natural science but a form of art and poetry. Mathematics is music – music is mathematics.


While studying mathematics, physics, and computer science, I took some courses on economics. That inspired me to write my bachelor's thesis on optimal control theory. I worked on the problem of how increasing public investments affects the GDP. I did not find any breakthrough, but it was an intriguing subject.

I proceeded with my graduate studies studying applied mathematics. I studied subjects like partial differential equations, functional analysis, dynamical systems, and – the University of Helsinki's pride – complex analysis. (My thesis advisor said that, in a moral sense, you cannot graduate from the University of Helsinki without taking some courses on Complex analysis, because a lot of the discipline has been developed at the university.) In addition, as a more “practical” subject, I studied some inverse problems. Summa summarum, I studied a wide range of fields in mathematics.


During my graduate studies, I spent half a year in Utrecht, Netherlands, studying more applied analysis of varying subjects (searching periodic orbits in the Lorentz attractor as an example of a course – that I failed). At Utrecht University, I got the inspiration for the subject for my future master's thesis; the Finite Element Method. After I got back to Helsinki from the exchange, I had a chance to study more about the finite element method in Aalto University's courses. (Aalto University is a consortium of the Helsinki University of Technology, the Helsinki School of Economics, and the University of Art and Design Helsinki.) While writing my thesis I also taught basic mathematics courses at the University of Helsinki and gained valuable experience in the pedagogical area.

In the binge of graduation, I started to look for future opportunities. I looked for coding jobs in Helsinki and Tallinn, jobs for mathematicians in the mapping industry, continuing at some universities to pursue a Ph.D., etc. I am glad I had the chance to use my creativity and continue in Aalto University's Department of Signal Processing and Acoustics to research low earth orbit satellite communications. The research methodology was from a stochastic geometry perspective, which was well aligned with my mathematical background.

My professional ambitions are in improving the lives of people globally. Communications play an essential in the picture. (But contain some challenging problems also, as we have seen with social media.) Through effective communication, we can share knowledge, control resources, discuss issues, etc. – however, globally, the communication infrastructure is still not nearly complete. My interests contain, but are not limited to, communications, particularly wireless networks and signal processing. My dream is to share my knowledge in the process toward a free and honest world. (Pardon me for the cliches.)















$$1 = e^{-\frac{\pi^2}{2\sqrt{P}}}$$





$$] = 1 - \left(1 - e^{-\frac{\pi^2}{2\sqrt{P/N}}} \right)^N,$$



1000



1992

$$2 \sqrt{r_1 r_2} \frac{1}{\sqrt{2\pi}} e^{-y^2/2} dy.$$



$$= \mathbb{E} \exp \left[- \sum_i f(|Y_i|) \right] = \mathbb{E} \prod_i \exp \left[- f(|Y_i|) \right]$$


$$= \mathbb{E}_n \left[(2\pi)^{-n/2} \int_0^\infty \cdots \int_0^\infty \prod_{i=0}^n \exp[-f(y_i)] \frac{2}{\sqrt{2\pi}} e^{-y_i^2} dy_1 \cdots dy_n \right]$$

$$= \mathbb{E}_n \left[(2\pi)^{-n/2} \left(\int_0^\infty 2 \exp[-f(y) - y^2] dy \right)^n \right]$$

$$= \mathbb{E}_n \exp \left[\sum_{i=0}^n \log \left(\int_0^\infty \exp[-f(y)] \frac{2e^{-y^2}}{\sqrt{2\pi}} dy \right) \right]$$



$$g = -\log\left(\int_0^{\infty} \exp[-f(v)] \frac{e^{-v^2}}{\sqrt{2\pi}} dv\right)$$




$$= \exp \left[- \int_{\mathbb{R}^2} 1 - e^{\log \left(\int_0^\infty \exp[-f(y)] \frac{2e^{-y^2}}{\sqrt{2\pi}} dy \right)} \Lambda(dx) \right]$$

$$= \exp \left[- \int_{\mathbb{R}^2} \left(1 - \int_0^\infty \exp[-f(y)] \frac{2e^{-y^2}}{\sqrt{2\pi}} dy \right) \Lambda(dx) \right]$$

$$= \exp \left[- \int_{\mathbb{R}^2} \int_0^\infty (1 - \exp[-f(y)]) \frac{2e^{-y^2}}{\sqrt{2\pi}} dy \Lambda(dx) \right]$$

$$= \exp \left[- \int_0^{\infty} (1 - \exp[-f(y)]) \sqrt{\frac{2}{\pi}} e^{-y^2} N dy \right], \quad (1)$$


$$\sqrt{\frac{2}{\pi}} e^{-x^2} \mathcal{N}.$$





192







$$t \rightarrow \text{order} \rightarrow \frac{\sin(2\pi B t)}{\pi t}$$






$$\mathbb{Z} \ni i \mapsto \frac{\sin\left(2\pi \frac{B_L}{f_c} i\right)}{\pi i},$$

$$i \mapsto \delta[i] - \frac{\sin(2\pi \frac{B_H}{f_c} i)}{\pi i},$$





গণপ্রজাতন্ত্রী বাংলাদেশ

2, 10x, 10x, 10x





WAVELENGTH



WINNERS

$$\mu = \exp\left(\mu_{\text{LN}} + \frac{\sigma_{\text{LN}}^2}{2}\right)$$

$$0 \leq \frac{0^2}{1} \leq \frac{1^2}{2} \leq \frac{2^2}{3} \leq \dots$$









FOR THE WIN














A 16x16 grayscale pixelated image of the letter 'A'. The image is composed of a grid of 256 pixels. The letter 'A' is rendered in a dark gray/black color, with some lighter gray pixels used for shading and detail. The background is white. The 'A' is formed by two main vertical strokes and a horizontal crossbar. The top of the 'A' is slightly wider than the bottom. There are some small gaps and irregularities in the pixelation, giving it a hand-drawn or low-resolution digital appearance.



A 16x16 grayscale pixelated image of the number 75. The image is composed of a grid of 256 pixels, with varying shades of gray representing the digits. The number 7 is on the left and the number 5 is on the right. The style is reminiscent of early digital art or low-resolution computer graphics.

A 16x16 grayscale pixelated image of the number 77. The image is composed of 256 individual pixels, each represented by a small square. The color of each pixel is determined by its value in a 16x16 grid of numbers from 0 to 255. The number 77 is rendered in a dark gray/black color, with the background being a lighter gray. The image is centered on a white background.

$$S(t) = \sum_{n=0}^{N-1} x[n]$$

















Q1 2024

2021年12月



$$\mathbb{E}[(S_1 + S_2)^2] = \int_0^1 (\cos(2\pi t) + \cos(2\pi t + \pi))^2 dt = \int_0^1 0 dt = 0,$$



1992 + 1992

$$= \int_0^1 \cos^2(2\pi t) dt + \int_0^1 \cos^2(2\pi t + \pi) dt$$

$$= \int_0^1 2 \cos^2(2\pi t) dt = \int_0^1 \cos(4\pi t) dt + 1 = 1,$$







$$E[S_1^2 + S_2^2] = E[S_1^2] + E[S_2^2] = E[S_1^2] + E[S_2^2],$$

1991 + 1991 + 1991

1992-1993

$$\mathbb{E}[S_1 S_2] = \int_0^1 \cos(2\pi t) \cos(2\pi t + \pi) dt = \int_0^1 -\cos^2(2\pi t) dt = -\frac{1}{2} \int_0^1 \cos(4\pi t) dt - \frac{1}{2} = -\frac{1}{2},$$










$$H = \begin{pmatrix} -a & -0 & -1 & 0 & 0 & 0 & ar & 0 & r \\ 0 & 0 & 0 & -a & -0 & -1 & 0 & 0 & 0 \\ -c & -b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & -b^2/a & -1 & cr & rb^2/a & r \\ a & 0 & -1 & 0 & 0 & 0 & ar & 0 & -r \\ 0 & 0 & 0 & a & 0 & -1 & 0 & 0 & 0 \\ -c & b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & b^2/a & -1 & -rc & rb^2/ar & -r \end{pmatrix}$$

$$\lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X \geq x)}{x},$$







$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x}$$

$$= \lim_{x \rightarrow \infty} \frac{-x/\theta}{x} = 1/\theta.$$





$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(1 - \gamma(k, x/\theta)/\Gamma(k))}{x}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(\kappa, x/\theta)/\Gamma(\kappa))}{x}$$

$$\begin{aligned}
 &\stackrel{(a)}{=} \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(k, x/\theta))}{x} = \lim_{x \rightarrow \infty} - \frac{\log(x^{k-1} e^{-x/\theta})}{x}
 \end{aligned}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x} = 1/\theta.$$



$$\lim_{x \rightarrow \infty} \frac{\Gamma(s, x)}{x^{s-1} e^{-x}} = 1.$$







$$\begin{aligned}
&= \lim_{x \rightarrow \infty} -\frac{\log \mathbb{P}(X > x)}{x} = -\lim_{x \rightarrow \infty} \frac{\log \operatorname{erfc}\left(\frac{x-\mu}{\sigma\sqrt{2}}\right)}{x} \stackrel{(b)}{\geq} -\lim_{x \rightarrow \infty} \frac{\log\left(\sqrt{\frac{e}{2\pi}} e^{-2\left(\frac{x-\mu}{\sigma\sqrt{2}}\right)^2}\right)}{x}
\end{aligned}$$

$$\geq -\lim_{x \rightarrow \infty} \frac{\log \left(e^{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2} \right)}{x} = -\lim_{x \rightarrow \infty} \frac{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2}{x} = \frac{1}{\sigma^2} \lim_{x \rightarrow \infty} \frac{x^2 - 2x\mu + \mu^2}{x} = \infty,$$



$$\psi(x) \geq \sqrt{\frac{2e}{\pi}} \frac{\sqrt{\beta - 1}}{\beta} e^{-\beta x^2},$$




































2020年12月20日