



x

1

7

x

10

0

1

0

1








PARADOXES

ESPRESSO










BEAR





THE FIRST TWO



100%







2023-10-21

$$= \underbrace{\frac{p_t^2}{2 \log(2)} \pi \lambda \left(\frac{h \varphi_{\text{RX}}}{\sin^2(\epsilon)} \right)^2}_{k \triangleq} \exp \left\{ - \tau^2 \underbrace{v_{\text{sat}}^2 \sin^4(\epsilon) / (h^2 \varphi_{\text{RX}}^2)}_{D \triangleq} \log(2) / 2 \right\},$$



























Fig 1(a): True $P(t)$, GPR estimate & forecast, and 100-sample MA & forecast






Fig 1(b): Non-stationary Gaussian noise $I(t)$ with sampled observations













expanding world + new world







Qatar Airways









$$S_{\text{RX}}(t) = \frac{A}{l(d_0)} \sqrt{\frac{K}{K+1}} e^{-2i\pi\tau_0(t)f_c} e^{-2i\pi\theta} + \sum_{j=1}^{100} \frac{A}{l(d_j) \sqrt{100} \sqrt{K+1}} e^{-2i\pi\tau_j(t)f_c} e^{-2i\pi\theta},$$




















QWERTYUIOP

$$v = \sqrt{\frac{GM}{h + R_{\oplus}}},$$

WORLD OF









2

=

12

.

100



$$T_c = \frac{1}{8D_g} \approx 10^{-3}$$




























$$1 = e^{-\frac{\pi^2}{2\sqrt{P}}}$$






$$] = 1 - \left(1 - e^{-\frac{\pi^2}{2\sqrt{P/N}}} \right)^N,$$





192








$$t \rightarrow \text{order} \rightarrow \frac{\sin(2\pi B t)}{\pi t}$$





$$\mathbb{Z} \ni i \mapsto \frac{\sin\left(2\pi \frac{B_L}{f_c} i\right)}{\pi i},$$

$$i \mapsto \delta[i] - \frac{\sin(2\pi \frac{B_H}{f_c} i)}{\pi i},$$



গণপ্রজাতন্ত্রী বাংলাদেশ

WAVELENGTH



WINNERS


$$\mu = \exp\left(\mu_{\text{LN}} + \frac{\sigma_{\text{LN}}^2}{2}\right)$$

$$O(2\pi) \cong SO(2) \cong \mathbb{R}/2\pi\mathbb{Z}$$










WORLD OF






A 16x16 pixelated grayscale image of the number 1. The number is rendered in a bold, blocky style using various shades of gray and black. The background is white. The number 1 is centered horizontally and vertically within the 16x16 grid.



A pixelated, black and white representation of the number 7. The image is composed of a grid of small squares, some of which are filled with black or gray, creating a blocky, digital appearance of the digit.

A pixelated, black and white representation of the number 20. The digit '2' is on the left, followed by an equals sign, and the digit '0' is on the right. The image has a low-resolution, dithered appearance.

$$S(t) = \sum_{n=0}^{N-1} x[n]$$
















$$H = \begin{pmatrix} -a & -0 & -1 & 0 & 0 & 0 & ar & 0 & r \\ 0 & 0 & 0 & -a & -0 & -1 & 0 & 0 & 0 \\ -c & -b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & -b^2/a & -1 & cr & rb^2/a & r \\ a & 0 & -1 & 0 & 0 & 0 & ar & 0 & -r \\ 0 & 0 & 0 & a & 0 & -1 & 0 & 0 & 0 \\ -c & b^2/a & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & -c & b^2/a & -1 & -rc & rb^2/ar & -r \end{pmatrix}$$

$$\lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X \geq x)}{x},$$







$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x}$$

$$= \lim_{x \rightarrow \infty} - \frac{x/\theta}{x} = 1/\theta.$$





$$= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = \lim_{x \rightarrow \infty} - \frac{\log(1 - \gamma(k, x/\theta)/\Gamma(k))}{x}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(\kappa, x/\theta)/\Gamma(\kappa))}{x}$$

$$\begin{aligned}
 &\stackrel{(a)}{=} \lim_{x \rightarrow \infty} - \frac{\log(\Gamma(k, x/\theta))}{x} = \lim_{x \rightarrow \infty} - \frac{\log(x^{k-1} e^{-x/\theta})}{x}
 \end{aligned}$$

$$= \lim_{x \rightarrow \infty} - \frac{\log(e^{-x/\theta})}{x} = 1/\theta.$$



$$\lim_{x \rightarrow \infty} \frac{\Gamma(s, x)}{x^{s-1} e^{-x}} = 1.$$







$$\begin{aligned}
&= \lim_{x \rightarrow \infty} - \frac{\log \mathbb{P}(X > x)}{x} = - \lim_{x \rightarrow \infty} \frac{\log \operatorname{erfc} \left(\frac{x - \mu}{\sigma \sqrt{2}} \right)}{x} \stackrel{(b)}{\geq} - \lim_{x \rightarrow \infty} \frac{\log \left(\sqrt{\frac{e}{2\pi}} e^{-2 \left(\frac{x - \mu}{\sigma \sqrt{2}} \right)^2} \right)}{x}
\end{aligned}$$

$$\geq -\lim_{x \rightarrow \infty} \frac{\log \left(e^{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2} \right)}{x} = -\lim_{x \rightarrow \infty} \frac{-2 \left(\frac{x-\mu}{\sigma \sqrt{2}} \right)^2}{x} = \frac{1}{\sigma^2} \lim_{x \rightarrow \infty} \frac{x^2 - 2x\mu + \mu^2}{x} = \infty,$$



$$\psi(x) \geq \sqrt{\frac{2e}{\pi}} \frac{\sqrt{\beta - 1}}{\beta} e^{-\beta x^2},$$









































Q1 2024

2023年12月



$$\mathbb{E}[(S_1 + S_2)^2] = \int_0^1 (\cos(2\pi t) + \cos(2\pi t + \pi))^2 dt = \int_0^1 0 dt = 0,$$



1921 + 1921

$$= \int_0^1 \cos^2(2\pi t) dt + \int_0^1 \cos^2(2\pi t + \pi) dt$$

$$= \int_0^1 2 \cos^2(2\pi t) dt = \int_0^1 \cos(4\pi t) dt + 1 = 1.$$





$$E[S_1^2 + S_2^2] = E[S_1^2] + E[S_2^2] = E[S_1^2] + E[S_2^2]$$

1991 + 1991 + 1991

1992-1993

$$\mathbb{E}[S_1 S_2] = \int_0^1 \cos(2\pi t) \cos(2\pi t + \pi) dt = \int_0^1 -\cos^2(2\pi t) dt = -\frac{1}{2} \int_0^1 \cos(4\pi t) dt - \frac{1}{2} = -\frac{1}{2},$$

01

2

02

E

10

2

02

2024 + 2025

2023-04-20

$$\mathbb{E} \phi_1 \phi_2 [\mathbb{E} [S_1^2 + S_2^2]] = \mathbb{E} \phi_1 [\mathbb{E} [S_1^2]] + \mathbb{E} \phi_2 [\mathbb{E} [S_2^2]] + \mathbb{E} \phi_1 \phi_2 [\mathbb{E} [S_1 S_2]]$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_1))^2 dt d\phi_1 + \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_2))^2 dt d\phi_2$$

$$+ \frac{1}{2\pi} \int_0^{2\pi} \int_0^{2\pi} \int_0^1 \cos(2\pi t + \phi_1) \cos(2\pi t + \phi_2) dt d\phi_1 d\phi_2$$

$$= \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_1))^2 dt d\phi_1 + \frac{1}{2\pi} \int_0^{2\pi} \int_0^1 (\cos(2\pi t + \phi_2))^2 dt d\phi_2 = 1/2 + 1/2 = 1.$$

1992



2020-2021





$${}_3F_2(1, 1, 1 + b; 2, 2; x) = \sum_{n=0}^{\infty} \frac{(1)_n (1)_n (1 + b)_n}{(2)_n (2)_n} \frac{x^n}{n!}$$

$$= \sum_{n=0}^{\infty} \frac{(1+b)_n}{(n+1)^2 n!} x^n = \frac{1}{b!} \sum_{n=0}^{\infty} \frac{(n+1)_b}{(n+1)^2} x^n$$

$$\underline{\underline{(a)}} \quad \frac{1}{b!} \sum_{n=0}^{\infty} \frac{\sum_{k=1}^b \begin{bmatrix} b \\ k \end{bmatrix} (n+1)^k}{(n+1)^2} x^n$$

$$\begin{aligned}
 &= \frac{1}{b!} \sum_{k=1}^b \left[\begin{matrix} b \\ k \end{matrix} \right] \sum_{n=0}^{\infty} \frac{x^n}{(n+1)^{2-k}} \stackrel{(b)}{=} \frac{1}{b!} \sum_{k=1}^b \left[\begin{matrix} b \\ k \end{matrix} \right] \frac{\operatorname{Li}_{2-k}(x)}{x},
 \end{aligned}$$



